

Mechanisms underlying firing in healthy and sick human motoneurons

Maria Piotrkiewicz, Parveen N. S. Bawa, Annie Schmied

► **To cite this version:**

Maria Piotrkiewicz, Parveen N. S. Bawa, Annie Schmied. Mechanisms underlying firing in healthy and sick human motoneurons. *Frontiers in Human Neuroscience*, Frontiers, 2015, 9 (174), 10.3389/fn-hum.2015.00174 . hal-01217103

HAL Id: hal-01217103

<https://hal-amu.archives-ouvertes.fr/hal-01217103>

Submitted on 19 Oct 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Mechanisms underlying firing in healthy and sick human motoneurons

Maria Piotrkiewicz^{1*}, Parveen N. S. Bawa^{2*} and Annie Schmied^{3*}

¹ Engineering of Nervous and Muscular System, Natęż Institute of Biocybernetics and Biomedical Engineering, Polish Academy of Sciences, Warsaw, Poland, ² Department of Biomedical Physiology and Kinesiology, Simon Fraser University, Burnaby, BC, Canada, ³ Centre National de la Recherche Scientifique, Plasticité et Pathophysiologie du Mouvement, Institut de Neurosciences de la Timone, University Aix Marseilles, Marseille, France

Keywords: human motoneuron discharge mechanisms, motor unit, afterhyperpolarization and delayed depolarization, double and triple discharges, recruitment and rate coding, axon and peripheral afferents, excitability, motoneuron pathology

In an address to the British Association for the Advancement of Science in Cambridge, Professor Sherrington introduced the terms “motor neurone” and “the final common path,” the latter term implying that all motor commands converge onto the motoneuron which integrates the incoming information and passes the net information to the muscle for contraction (Sherrington, 1904). The relative ease of access of the spinal motoneuron made it feasible to set up techniques for investigating the physiological, biophysical and molecular properties of these neurons. It became the most investigated neuron of the CNS in the twentieth century and the information gained from studies on motoneurons formed the basis for examining the other neurons of the CNS. Since the compound action potential of a muscle unit is strictly related one-to-one to the action potential arriving from the innervating motoneuron, the statistical analysis of muscle unit action potentials provides an investigator with an elegant way to probe the properties of motoneurons in behaving humans. In the following review the terms *motoneuron* and *motor unit* might be used interchangeably. Different aspects of human motoneuron investigations in health and disease are presented in 16 articles of this topic which are summarized below.

An increase in the net excitatory synaptic input to the motoneuron pool results in an increase in the level of muscle contraction by recruitment of additional motor units (MUs) and an increase in firing rates of the already recruited units (Milner-Brown et al., 1973; Henneman et al., 1974). The principle of orderly recruitment of motoneurons by size was originally proposed by Henneman (1957) but was later questioned by other researchers presenting examples of selective, rather than orderly recruitment (e.g., Smith et al., 1980). These controversies are assessed by Bawa et al. (2014), and the opinion unifying the concept of orderly recruitment is presented. In humans, increases in firing rates of motor units have been shown to follow the “onion skin” pattern at lower levels of contraction, meaning that the lower-threshold motor units discharge with higher rates than higher-threshold ones. However, studies performed on the whole range of muscle forces indicated that for higher force levels the motor unit firing rate follows a “reverse onion skin” pattern. Hu et al. (2014) decided to approach this problem using small surface electrodes and step increases in force instead of the “ramp and hold” protocols used by previous authors. They showed that the “onion skin” pattern was preserved until 15% of maximal voluntary contraction, and from their results predict this pattern to be valid for the whole range of muscle forces, which is not supported by the previous published works. However, the reported rate saturation of the MUs discharging with higher rates implies that at the higher forces the “reverse onion skin” pattern may be expected. In another paper, Duchateau and Baudry (2014) show that during ballistic contractions the maximal discharge rates are higher than those observed in ramp contractions. It should be noted, however, that during ballistic contractions one deals with instantaneous rates, while during ramp and hold contractions one refers to tonic firing rates defined as the average

OPEN ACCESS

Edited and reviewed by:

Hauke R. Heekeren,
Freie Universität Berlin, Germany

*Correspondence:

Maria Piotrkiewicz,
masia@ibib.waw.pl;
Parveen N.S. Bawa,
bawa@sfu.ca;
Annie Schmied,
annieschmied@yahoo.fr

Received: 22 January 2015

Accepted: 12 March 2015

Published: 30 March 2015

Citation:

Piotrkiewicz M, Bawa PNS and Schmied A (2015) Mechanisms underlying firing in healthy and sick human motoneurons. *Front. Hum. Neurosci.* 9:174. doi: 10.3389/fnhum.2015.00174

over 1 s. One cannot compare maximal rates during the two patterns of contraction. The authors also suggest that the maximal rate of force development is determined by maximal instantaneous firing rate of the motoneurons confirming earlier work on reduced cat preparations using intracellular current injections (Baldissera et al., 1982). For information of firing rates in older adults, the maximal firing rates have been reported to decline (Duchateau and Baudry, 2014; Kallio et al., 2014).

The high instantaneous firing rates have also been observed at low, slower speeds of muscle contractions. They are generated by some motoneurons, which occasionally fire pairs of closely spaced spikes (doublets); each pair is followed by a prolonged post doublet interval suggested to result from the summation of successive afterhyperpolarizations (AHPs). The “true” doublets (Bawa and Calancie, 1983) should be distinguished from the short interspike intervals fired during ballistic contractions. The former are attributed to the delayed depolarization, an intrinsic property of the motoneuron (Kernell, 1964; Calvin, 1973), while the latter are due to the high rate of rise in synaptic inputs (Baldissera et al., 1982). Doublets have been reported by several authors contributing to the present topic. Piotrkiewicz et al. (2013) observed doublets in the soleus muscle where they have never been documented before. Repetitive doublets, where doublet/postdoublet intervals alternate, were first reported by Bawa and Calancie (1983). Kudina and Andreeva (2010) suggested earlier that repetitive doublets resulted from suprathreshold delayed depolarization supported by plateau potentials. In the present article, Kudina and Andreeva (2013) pose the question whether there are common attributes of motoneurons in the spinal cord which can fire repetitive doublets. They suggest a cranio-caudal gradient of the number of motoneurons that can discharge doublets and more frequent repetitive doublets observed in the cervical motoneurons compared to the motoneurons of the lumbar region. Piotrkiewicz and Kuraszkiwicz (2014) investigated the relationship between duration of motoneuron AHP and confirmed an earlier observation by Kernell (1964) that motoneurons with shorter AHPs discharge doublets more easily. Doublets were also mentioned in Sogaard et al. (2014). Triplet discharges have been reported by Piotrkiewicz and Kuraszkiwicz (2014) and the authors speculate on the possible mechanisms to explain this pattern.

Three papers deal with problems of motoneuron excitability. Various methods used to date to test excitability of human motoneuron pools are reviewed by McNeil et al. (2013). While synaptic inputs to a motoneuron pool recruit motoneurons from small to large, electrical stimulation of a muscle nerve recruits motoneurons in reverse order, from larger to smaller axons. It has been acknowledged for some time that the diameter is not the only factor determining axon excitability. In a mixed nerve, sensory and motor fibers have different biophysical properties making sensory afferents more excitable than motor axons of the same size. Human studies of axon excitability have obvious limitations, thus Lorenz and Jones (2014) explored this problem in a rat model. They have shown that several biophysical parameters differ between axons innervating slow soleus and fast tibialis anterior. These mechanisms may underlie the bimodal distribution of axon excitability observed in the tibialis anterior

by Kudina and Andreeva (2014). Since the axons of larger motoneurons are more excitable, electrical stimulation used for rehabilitation purposes might only recruit and strengthen the larger motor units and let the small units succumb to atrophy. Dean et al. (2014) have proposed a method to recruit smaller low threshold units using specially tailored electrical stimulation of a mixed nerve. The stimulus consists of a high frequency pulse train with current strength subthreshold for eliciting M or H waves. In soleus, stimulation of the posterior tibial nerve with such parameters has been shown to recruit motor units in an orderly fashion from small to large, and at physiological firing rates thus leading to activation and strengthening of small motor units.

Six papers deal with motoneuron discharge related pathology. Sogaard et al. (2014) recorded electromyographic activity bilaterally from upper trapezius muscles. They observed a tonic low level motor unit activity and suggested that such activity, whether it is built into the motor program to stabilize the limbs or results from stress of paying attention during computer work, may be the underlying cause of myalgia observed in computer workers. Garland et al. (2014) reviewed stroke-related changes in motor unit discharge characteristics and suggested that residual motor control strategies may remain after stroke. McNulty et al. (2014) investigated motor unit firing characteristics in different muscles of stroke survivors showing that the effects on peak firing rates and their dynamic range differ not only between joints of the upper and lower limbs but also between muscles of different joints of the same limbs. Furthermore, they have shown that motor units on both paretic and non-paretic sides changed after stroke. Confirming the common observation that firing rates on the affected side are lower than normal, they have also shown that motor units from the unaffected side discharge with firing rates higher than normal. The authors conclude that motor unit properties on both sides should be compared to data from age- and sex-matched healthy subjects.

Neuroscientists have used peripheral inputs to investigate motor output and various properties of motoneurons. Yet, when it comes to voluntary control of movement, the peripheral afferents are generally ignored. In their paper describing extensive single motor unit recordings from a deafferented subject, Schmied et al. (2014) demonstrate the importance of peripheral afferents in the control of motoneuron excitability, variability in firing, synchronization and coherence between different motor units. The viability of motoneurons and the dependence of their excitability on peripheral afferents are discussed by Zijdwind et al. (2014). In incomplete spinal cord injured subjects the mean firing rates decrease compared to those in normal subjects. The authors argue that the decrease in firing rates is not due to changes in motoneuron properties resulting from a decrease or complete elimination of descending inputs. These motoneurons are capable of reaching normal discharge rates during spasms. However, the covariation in firing rates among various motor units mentioned above by Hu et al. (2014) is lacking during spasms, which implies that the weight of different inputs onto different motoneurons of a pool might alter in spinal cord injury. This might also apply to persistent inward currents of the members of a pool.

Motoneurons form the final motor path for all the motor commands; an animal without motoneurons would die. Such is the case in amyotrophic lateral sclerosis (ALS), a deadly disease of unknown etiology, which causes selective degeneration of spinal motoneurons and corticomotoneuronal cells (“lower” and “upper” motoneurons, respectively). De Carvalho et al. (2014) review the properties of the upper and lower motoneurons in ALS, aiming to show how changes in the pattern of motor unit firing could help to delineate the underlying pathophysiological disturbance as the disease progresses. However, some of the few single motor unit studies performed in patients with ALS have

not been covered in this review. The interested reader may find additional information in Schmied et al. (1999), Attarian et al. (2006, 2008).

The articles collected in this exciting Research Topic cover a broad spectrum of human motoneuron research, from their intrinsic properties such as afterhyperpolarization following delayed depolarization to pathological changes in neuromuscular disorders. We hope that this collection will be equally exciting for the potential readers. Our sincere thanks are expressed to all the authors and reviewers who contributed to this important topic of human neurophysiology.

References

- Attarian, S., Vedel, J. P., Pouget, J., and Schmied, A. (2006). Cortical versus spinal dysfunction in amyotrophic lateral sclerosis. *Muscle Nerve* 33, 677–690. doi: 10.1002/mus.20519
- Attarian, S., Vedel, J. P., Pouget, J., and Schmied, A. (2008). Progression of cortical and spinal dysfunctions over time in amyotrophic lateral sclerosis. *Muscle Nerve* 37, 364–375. doi: 10.1002/mus.20942
- Baldissera, F., Campadelli, P., and Piccinelli, L. (1982). Neural encoding of input transients investigated by intracellular injection of ramp currents in cat alpha-motoneurons. *J. Physiol.* 328, 73–86.
- Bawa, P., and Calancie, B. (1983). Repetitive doublets in human flexor carpi radialis muscle. *J. Physiol.* 339, 123–132.
- Bawa, P. N. S., Jones, K. E., and Stein, R. B. (2014). Assessment of size ordered recruitment. *Front. Hum. Neurosci.* 8:532. doi: 10.3389/fnhum.2014.00532
- Calvin, W. H. (1973). “A third mode of repetitive firing: self-regenerative firing due to large delayed depolarization,” in *Control of Posture and Locomotion*, eds R. Stein, K. Pearson, R. Smith, and J. Redford (New York, NY: Plenum Press), 173–178.
- Dean, J., Clair-Augier, J., Lagerquist, O., and Collins, D. (2014). Asynchronous recruitment of low-threshold motor units during repetitive, low-current stimulation of the human tibial nerve. *Front. Hum. Neurosci.* 8:1002. doi: 10.3389/fnhum.2014.01002
- De Carvalho, M., Eisen, A., Krieger, C., and Swash, M. (2014). Motoneuron firing in amyotrophic lateral sclerosis (ALS). *Front. Hum. Neurosci.* 8:719. doi: 10.3389/fnhum.2014.00719
- Duchateau, J., and Baudry, S. (2014). Maximal discharge rate of motor units determines the maximal rate of force development during ballistic contractions in human. *Front. Hum. Neurosci.* 8:234. doi: 10.3389/fnhum.2014.00234
- Garland, S. J., Pollock, C. L., and Ivanova, T. D. (2014). Could motor unit control strategies be partially preserved after stroke? *Front. Hum. Neurosci.* 8:864. doi: 10.3389/fnhum.2014.00864
- Henneman, E. (1957). Relation between size of neurons and their susceptibility to discharge. *Science* 126, 1345–1347.
- Henneman, E., Clamann, H. P., Gillies, J. D., and Skinner, R. D. (1974). Rank order of motoneurons within a pool: law of combination. *J. Neurophysiol.* 37, 1338–1349.
- Hu, X., Rymer, W. Z., and Suresh, N. L. (2014). Control of motor unit firing during step-like increases in voluntary force. *Front. Hum. Neurosci.* 8:721. doi: 10.3389/fnhum.2014.00721
- Kallio, J., Sogaard, K., Avela, J., Komi, P. V., Selänne, H., and Linnamo, V. (2014). Differences between young and elderly in soleus motor unit discharge rate in dynamic movements. *Front. Hum. Neurosci.* 8:773. doi: 10.3389/fnhum.2014.00773
- Kernell, D. (1964). “The delayed depolarization in cat and rat motoneurons,” in *Progress in Brain Research*, eds J. Eccles and J. P. Schade (Amsterdam: Elsevier), 42–52.
- Kudina, L., and Andreeva, R. (2010). Repetitive doublet firing of motor units: evidence for plateau potentials in human motoneurons? *Exp. Brain Res.* 204, 79–90. doi: 10.1007/s00221-010-2298-z
- Kudina, L. P., and Andreeva, R. E. (2013). Delayed depolarization and firing behavior of human motoneurons during voluntary muscle contractions. *Front. Hum. Neurosci.* 7:793. doi: 10.3389/fnhum.2013.00793
- Kudina, L. P., and Andreeva, R. E. (2014). Excitability properties of single human motor axons: are all axons identical? *Front. Cell Neurosci.* 8:85. doi: 10.3389/fncel.2014.00085
- Lorenz, C., and Jones, K. E. (2014). I_H activity is increased in populations of slow versus fast motor axons of the rat. *Front. Hum. Neurosci.* 8:766. doi: 10.3389/fnhum.2014.00766
- McNeil, C. J., Butler, J. E., Taylor, J. L., and Gandevia, S. C. (2013). Testing the excitability of human motoneurons. *Front. Hum. Neurosci.* 7:152. doi: 10.3389/fnhum.2013.00152
- McNulty, P. A., Lin, G. G., and Doust, C. G. (2014). Single motor unit firing rate after stroke is higher on the less-affected side during stable low-level voluntary contractions. *Front. Hum. Neurosci.* 8:518. doi: 10.3389/fnhum.2014.00518
- Milner-Brown, H. S., Stein, R. B., and Yemm, R. (1973). Changes in firing rate of human motor units during linearly changing voluntary contractions. *J. Physiol.* 230, 371–390.
- Piotrkiewicz, M., and Kuraszkiwicz, B. (2014). Double discharges and afterhyperpolarization in human motoneurons. *Front. Hum. Neurosci.* 8:373. doi: 10.3389/fnhum.2014.00373
- Piotrkiewicz, M., Sebik, O., Binboga, E., Mlozniak, D., Kuraszkiwicz, B., and Turker, K. S. (2013). Double discharges in human soleus muscle. *Front. Hum. Neurosci.* 7:843. doi: 10.3389/fnhum.2013.00843
- Schmied, A., Forget, R., and Vedel, J.-P. (2014). Motor unit firing pattern, synchrony and coherence in a deafferented patient. *Front. Hum. Neurosci.* 8:746. doi: 10.3389/fnhum.2014.00746
- Schmied, A., Pouget, J., and Vedel, J. P. (1999). Electromechanical coupling and synchronous firing of single wrist extensor motor units in sporadic amyotrophic lateral sclerosis. *Clin. Neurophysiol.* 110, 960–974.
- Sherrington, C. S. (1904). “Correlation of reflexes and the principle of the common path,” in *British Association for the Advancement of Science, Section I-Physiology* (London: Burlington House), 728–740.
- Smith, J. L., Betts, B., Edgerton, V. R., and Zernicke, R. F. (1980). Rapid ankle extension during paw shakes: selective recruitment of fast ankle extensors. *J. Neurophysiol.* 43, 612–620.
- Sogaard, K., Olsen, H. B., Blangsted, A. K., and Sjogaard, G. (2014). Single motor unit firing behaviour in the right trapezius muscle during rapid movement of right or left index finger. *Front. Hum. Neurosci.* 8:881. doi: 10.3389/fnhum.2014.00881
- Zijdewind, I., Bakels, R., and Thomas, C. K. (2014). Motor unit firing rates during spasms in the muscles of spinal cord injured subjects. *Front. Hum. Neurosci.* 8:922. doi: 10.3389/fnhum.2014.00922

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Piotrkiewicz, Bawa and Schmied. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.