

Evaluation of Serum Lipase as Predictor of Severity of Acute Pancreatitis in Children

Alexandre Fabre, Ophélie Boulogne, Jean Gaudart, E Mas, J.P. Olive,
Jacques Sarles

► **To cite this version:**

Alexandre Fabre, Ophélie Boulogne, Jean Gaudart, E Mas, J.P. Olive, et al.. Evaluation of Serum Lipase as Predictor of Severity of Acute Pancreatitis in Children. Journal of Pediatric Gastroenterology and Nutrition, Lippincott, Williams & Wilkins, 2014, 10.1097/MPG.0000000000000307. hal-01307534

HAL Id: hal-01307534

<https://hal-amu.archives-ouvertes.fr/hal-01307534>

Submitted on 26 Apr 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



You could be reading the full-text of this article now if you...

Become a subscriber

Purchase this article

If you have access to this article through your institution,
you can view this article in

Ovid

Journal of Pediatric Gastroenterology & Nutrition:

April 2014 - Volume 58 - Issue 4 - p e41–e42

doi: 10.1097/MPG.0000000000000307

Letters to the Editor

Evaluation of Serum Lipase as Predictor of Severity of Acute Pancreatitis in Children

Fabre, Alexandre*; Boulogne, Ophélie[†]; Gaudart, Jean[†]; Mas, Emmanuel[†]; Olives, Jean-Pierre[‡]; Sarles, Jacques*

CONTENT NOT FOR REUSE

LETTERS TO THE EDITOR

Preemptive Meso-Rex Bypass for Children With Idiopathic Prehepatic Portal Hypertension: Trick or Treat?

To the Editor: Once upon a time, in an era when the physiopathology of the Hirschsprung disease was still unknown, fixation was on the congenitally dilated colon (1). Patients were followed with conservative means or palliative operations (colostomies or resection of dilated segments). The concept of a surgical cure was introduced much later, only 50 years ago; in a few decades, it changed the life of these children. Contemporary single-step, transperineal, neonatal interventions cure preemptively these children with no hesitation.

Alberti et al (2) proposed a stepwise approach to extrahepatic portal vein obstruction in children. Rex recessus was found patent at portography in 24 children and 13 were proposed for Meso-Rex bypass (MRB); 3 of the 13 MRBs were done preemptively (elective operation in patients with well-controlled esophageal varices) and the authors consider it a violation of their protocol at that time. Although Alberti et al end their discussion with only 1 word about preemptive cure as an opportunity for some children, the latter message is not clearly delivered to the audience.

Beta-blockers, with their unproven value and repeated endoscopic procedures (254 sclerotherapies or bandings in this report), are neither curative nor without risk, although the authors choose these palliative procedures over curative surgery in the vast majority of children with symptoms.

Extrahepatic portal hypertension, caused by portal vein occlusion, is a chronic condition carrying multiple risks and associated with many penalties to the child development and well-being: a cure can be proposed to selected children with a favorable anatomy (patent Rex recessus) by performing an MRB or similar operations. This procedure should be proposed earlier than later in the disease progression, electively and preemptively (3,4).

*Riccardo A. Superina and Jean de Ville de Goyet
†Department of Pediatric Surgery, Children's Memorial Hospital,
Chicago, IL

‡Department of Pediatric Surgery, Bambino Gesù Children's
Hospital, Rome, Italy

REFERENCES

- Geosfeld JL. Hirschsprung disease and allied disorders. In: Holschneider AM, Puri P, eds. *Hirschsprung Disease: A Historical Perspective—1691–2005*. Berlin: Springer-Verlag; 2008:1–12.
- Alberti D, Colusso M, Cheli M, et al. Results of a stepwise approach to extra-hepatic portal vein obstruction in children. *J Pediatr Gastroenterol Nutr* 2013;57:619–26.
- Schweizer B, Enre S, Grossmann R, et al. Expert pediatric opinion on the report of the Bavaria IV consensus workshop on methodology of diagnosis and therapy in portal hypertension. *Pediatr Transplant* 2006;10:887–907.
- Superina R, Schneider B, Enre S, et al. Surgical guidelines for the management of extra hepatic portal vein obstruction. *Pediatr Transplant* 2006;10:908–13.

Authors' Response

To the Editor: Our study reports the outcome of children with extrahepatic portal vein obstruction managed prospectively with the same protocol for 15 years, without any selection. In this protocol, surgery (with priority to meso-portal bypass [MPB]) was indicated only after the failure of medical and endoscopic treatments. Accurate methodology suggested considering our latest cases, which had a preemptive MPB, as violations. With these data, we could certainly not argue in favor of or against different approaches, although the violations tell that we probably agree with the comment made.

We hope that Dr Superina and de Ville will have the chance to produce similar data on a large, unselected population of children presenting with extrahepatic portal vein obstruction and managed prospectively with a protocol considering preemptive MPB as the first step. We look forward to reading such a study that could prove that preemptive MPB is superior to MPB carried out as a second-line treatment (1). Meanwhile, we appreciate the comment and we agree with what, at the moment, is an expert opinion (2).

*Daniele Alberti and †Lorenzo D'Antiga
‡Pediatric Surgery, University of Brescia, Presidio Ospedale dei
Bambini, Brescia
§Paediatric Hepatology, Gastroenterology, and Transplantation,
Hospital Papa Giovanni XXIII, Bergamo, Italy

REFERENCES

- Alberti D, Colusso M, Cheli M, et al. Results of a stepwise approach to extra-hepatic portal vein obstruction in children. *J Pediatr Gastroenterol Nutr* 2013;57:619–26.
- Schneider B, Bosch J, de Franchis R, et al. Portal hypertension in children: expert pediatric opinion on the report of the Bavaria V Consensus Workshop on Methodology of Diagnosis and Therapy in Portal Hypertension. *Pediatr Transplant* 2012;16:426–37.

Evaluation of Serum Lipase as Predictor of Severity of Acute Pancreatitis in Children

To the Editor: The search for an easy-to-use and accurate predictor tool for the measurement of severity in acute pancreatitis is a never-ending quest. In the June 2013 issue of the *Journal of Pediatric Gastroenterology and Nutrition*, Coffey et al (1) suggested that serum lipase $\geq 7 \times$ the upper limit of normal (ULN) is a simple predictor of acute pancreatitis severity. Its sensitivity was 85% and specificity 56%. In 2012, we published a study evaluating different scoring systems for acute pancreatitis in a cohort of 48 children (2). Because we had the lipase level for 44 of them, we could test the usefulness of serum lipase in our cohort. Among these 44 children, 11 were classified as having severe pancreatitis and 33 with its mild form, according to the criteria