

Ticagrelor Improves Peripheral Arterial Function in Acute Coronary Syndrome Patients: Relationship With Adenosine Plasma Level.

Julien Fromonot, Françoise Dignat-George, Pascal Rossi, Giovanna Mottola, Nathalie Kipson, Jean Ruf, L. Bonello, Régis Guieu, F. Paganelli

► **To cite this version:**

Julien Fromonot, Françoise Dignat-George, Pascal Rossi, Giovanna Mottola, Nathalie Kipson, et al.. Ticagrelor Improves Peripheral Arterial Function in Acute Coronary Syndrome Patients: Relationship With Adenosine Plasma Level.. Journal of the American College of Cardiology, Elsevier, 2016, 67 (16), pp.1967-1968. 10.1016/j.jacc.2016.02.023 . hal-01311749

HAL Id: hal-01311749

<https://hal-amu.archives-ouvertes.fr/hal-01311749>

Submitted on 18 Oct 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Ticagrelor Improves Peripheral Arterial Function in Acute Coronary Syndrome Patients

Relationship With Adenosine Plasma Level

Ticagrelor, a P2Y₁₂ receptor antagonist, significantly reduces the incidence of major cardiovascular events in acute coronary syndrome (ACS) compared with standard treatment with clopidogrel (1). It has been suggested that this benefit effect may be accounted for by its pleiotropic properties via a purinergic mechanism. Indeed, ticagrelor increases the endogenous adenosine plasma level (APL) via red blood cell uptake inhibition in primary ACS patients compared with clopidogrel (2). However, the link between these “pleiotropic” properties and the improvement in microvascular dysfunction (MiD) remains poorly investigated. This study aimed to determine whether the increase in APL achieved with ticagrelor might improve endothelial dysfunction in primary ACS patients.


We prospectively randomized 60 primary ACS patients to receive ticagrelor (180-mg loading dose followed by 90 mg twice daily) or clopidogrel (600-mg loading dose followed by 75 mg/day). In addition, all patients received 250 mg of ticagrelor

followed by 75 mg of aspirin daily. Exclusion criteria were New York Heart Association functional class III or higher, cardiac arrest, contraindications to antiplatelet therapy, treatment with a P2Y₁₂-adenosine diphosphate receptor antagonist for <1 month, a platelet count <100 g/l, a history of bleeding diathesis, a history of stroke, recent surgery (preceding month), use of a medication known to interfere with ticagrelor, and bradycardia. Blood samples (3 ml) were collected under basal conditions (just before the primary loading dose) and after 30 days of treatment. APL was measured as previously described using high-pressure liquid chromatography (2).

To assess MiD, we performed digital peripheral artery tonometry at the same time to evaluate the reactive hyperemia index (RHI). Endothelial dysfunction was suspected if the RHI was <1.67. The methodology has been previously described (3). Briefly, each recording consisted of 5 min of baseline measurements, 5 min of occlusion measurements, and then 5 min of post-occlusion measurements (hyperemic period) (3). The RHI was defined as the ratio of the digital pulse volume during reactive hyperemia to that observed at baseline. Adverse cardiovascular events were recorded.

Under basal conditions, all the patients had MiD (RHI <1.67; mean, 1.37 ± 0.12). The APL (Figure 1A) and RHI (Figure 1B) did not differ between the 2 groups of patients. At day 30, the APL increased significantly only in the ticagrelor group (Figure 1A). The APL was more than 2-fold higher in patients taking ticagrelor

FIGURE 1 APL and RHI 30 Days After Treatment



Continuous variables were expressed as mean ± SD or median and interquartile range. The Wilcoxon test was used for intraindividual comparisons. Analysis of variance was used for intergroup comparisons. Pearson's r coefficient of correlation was used for a correlation study. APL (A) or RHI (B) was measured before (basal) and after 30 days of treatment. Correlation between the mean APL was evaluated after 30 days of ticagrelor treatment, the mean APL was evaluated before treatment (DAPL), and the RHI was evaluated after 30 days of ticagrelor treatment. The RHI was evaluated under basal conditions (DRHI). (a) $p < 0.01$ compared with basal conditions. (b) $p < 0.01$ compared with the clopidogrel group. APL = adenosine plasma level; RHI = reactive hyperemia index.

than in patients taking clopidogrel (Figure 1A). After 30 days, the mean RHI increased slightly (mean, \uparrow 15%) in the clopidogrel group and greatly (\uparrow 100%) in the ticagrelor group (Figure 1B). We found a correlation between the increase in the APL and the increase in the RHI in the ticagrelor group. No ischemic or bleeding events were recorded within 1 month. Dyspnea was present in 7 patients in the ticagrelor group and in 5 patients in the clopidogrel group.

Treatment with ticagrelor improves peripheral arterial function compared with clopidogrel. Endothelial function responds quickly to ticagrelor therapy within 30 days, long before the effects on clinical outcomes are seen. The underlying mechanism seems to be the APL increase. It was shown that clopidogrel slightly improved endothelial function (4); however, we found that these effects were not related to an increase in adenosine plasma concentration. It was shown that the ticagrelor dose dependently increased adenosine-mediated coronary blood flow in healthy human subjects (5). These effects occur mostly via the activation of A_{2A} adenosine receptors. Here, we found that ticagrelor induces an increase in the APL that correlates with the increase in the RHI. Because a low RHI is associated with a higher adverse rate during follow-up, our study led to the hypothesis that the adenosine-mediated effects of ticagrelor may explain the accuracy of the PLATO (A Comparison of Ticagrelor (AZD6140) and Clopidogrel in Patients With Acute Coronary Syndrome) mortality data (1).

Finally, our study demonstrates a correlation between the increase in the APL and improvement in the RHI, but whether this increase in the APL has a direct effect on endothelial function remains to be established.

Julien Fromonot, PhD
Françoise Dignat-Georges, PhD
Pascal Rossi, MD, PhD
Giovanna Mottola, PhD
Nathalie Kipson, BSc
Jean Ruf, PhD
Laurent Bonello, MD, PhD
*Régis Guieu, MD, PhD
Franck Paganelli, MD
*UMR MD2
Aix Marseille University
Institute of Biological Research of the
French Army (IRBA)
Boulevard P. Dramard
Marseille
France
E-mail: guieu.regis@orange.fr
<http://dx.doi.org/10.1016/j.jacc.2016.02.023>

Please note: Dr. Bonello has received a research grant from AstraZeneca. All other authors have reported that they have no relationships relevant to the contents of this paper to disclose.

REFERENCES

1. Wallentin L, Becker RC, Budaj A, et al. Ticagrelor versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med* 2009;361:1045-57.
2. Bonello L, Laine M, Kipson N, et al. Ticagrelor increases adenosine plasma concentration in patients with an acute coronary syndrome. *J Am Coll Cardiol* 2014;63:872-7.
3. Rubinshtein R, Kuvin JT, Soffler M, et al. Assessment of endothelial function by non-invasive peripheral arterial tonometry predicts late cardiovascular adverse events. *Eur Heart J* 2010;31:1142-8.
4. Willoughby SR, Luu LJ, Cameron JD, et al. Clopidogrel improves microvascular endothelial function in subjects with stable coronary artery disease. *Heart Lung Circ* 2014;23:534-41.
5. Wittfeldt A, Emanuelsson H, Brandrup-Wognsen G, et al. Ticagrelor enhances adenosine-induced coronary vasodilatory responses in humans. *J Am Coll Cardiol* 2013;61:723-7.