One-pot preparation of 2-(alkyl)arylbenzoselenazoles from the corresponding N-(acetyl)benzoyl-2-iodoanilines via a microwave-assisted methodology

Sébastien Redon, Youssef Kabri, Maxime D. Crozet, Patrice Vanelle

► To cite this version:

Sébastien Redon, Youssef Kabri, Maxime D. Crozet, Patrice Vanelle. One-pot preparation of 2-(alkyl)arylbenzoselenazoles from the corresponding N-(acetyl)benzoyl-2-iodoanilines via a microwave-assisted methodology. *Tetrahedron Letters*, 2014, 55 (36), pp.5052-5054. 10.1016/j.tetlet.2014.07.055 . hal-01373020v2

HAL Id: hal-01373020

<https://hal-amu.archives-ouvertes.fr/hal-01373020v2>

Submitted on 4 Jan 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

One-pot preparation of 2-(alkyl)arylbenzoselenazoles from the corresponding *N*-(acetyl)benzoyl-2-iodoanilines *via* a microwave-assisted methodology.

Sébastien Redon, Youssef Kabri, Maxime D. Crozet and Patrice Vanelle*

Aix-Marseille Université, UMR CNRS 7273, Institut de Chimie Radicalaire (ICR), Laboratoire de Pharmaco-Chimie Radicalaire, Faculté de Pharmacie, 27 Boulevard Jean Moulin, CS 30064, 13385 Marseille Cedex 05, France

ARTICLE INFO

Article history:

Received

Received in revised form

Accepted

Available online

ABSTRACT

We report here the first example of a one-pot synthesis of 2-(alkyl)arylbenzoselenazoles from *N*-(acetyl)benzoyl-2-iodoanilines. The reaction was carried out in the presence of Woollins' reagent under microwave irradiation and resulted in moderate to good yields.

Keywords:

Woollins' reagent

Benzoselenazole

Microwave irradiation

Cyclization

The value of the selenium-containing heterocycles has been demonstrated in medicinal chemistry,¹ offering antiviral,² antihypertensive,³ antifungal⁴ and anticancer⁵ properties. Some 1,3-benzoselenazole derivatives are also useful as precursors of new materials such as cyanine-type dyes.⁶ Compared to their sulfur homologues, selenium compounds exhibit very valuable antioxidant properties.⁷ However, synthetic routes remain poorly explored because of their high cost or low stability in air.

In continuation of our research program on medicinal chemistry⁸ we investigate the synthesis of benzoselenazoles as homologues of benzothiazoles. The few existing synthetic routes it should be notified the condensation of zinc bis(*o*-aminoselenoate) with acid chlorides^{9a} or bis(*o*-aminophenyl) diselenide with aldehydes promoted by sodium metabisulfite.^{9b} Recently an efficient method was developed *via* intermolecular C–Se cross-coupling catalyzed by transition metal giving 2-aminobenzoselenazole from 2-iodo-4-methylphenyl)selenourea.¹⁰ Sashida improved this method *via* a one-pot reaction starting from iodoaniline and isoselenocyanates.¹¹ Kambe reported a copper(I)-catalyzed reaction of 2-bromophenylisocyanide with selenium and heteroatom nucleophiles.¹²

We focused on an existing benzothiazole synthesis *via* an intramolecular nucleophilic aromatic substitution of *o*-halo-thiobenzanilide (INASOB) promoted by base.¹³ We used the well-known Woollins' reagent¹⁴ (WR) to convert the amides into the corresponding selenoamides. We initially explored the reaction of **1a** with 1.2 equiv. of WR under classical heating

procedure (reflux of THF, toluene). Unfortunately, the starting material was entirely recovered without any trace of the expected product (entries 1-2, Table 1). However, this reaction, under reflux of organic base (pyridine), afforded the 2-methylbenzoselenazole **3a** alone with no trace of selenoamide after 16 h (entry 3). Although the yield was low, this encouraging result showed that the selenation/cyclization sequence could be realized under one-pot conditions. With a higher boiling point solvent (mixture of xylenes), a better though still low yield was obtained without any trace of intermediate selenamide **2** (entry 4). With this classical heating procedure, **1a** was still present after reaction and the yield of **3a** remained low, we therefore investigated whether the selenation/cyclization sequence could be promoted under microwave irradiation.¹⁵ Using pyridine as the solvent at 160 °C under microwave irradiation, we monitored the reaction by LCMS. The starting material was completely consumed after 6 h but 2-methylbenzoselenazole **3a** was obtained in moderate yield (entry 5). Variation of the amount of WR (2 equiv) slightly improved the yield to 41% (entry 6). Next, pyridine was replaced by NEt₃ at the same temperature (160 °C). The reaction was finished in 3 h and 2-methylbenzoselenazole **3a** was obtained in higher yield (57%) (entry 7). With other base (2 equiv of Cs₂CO₃ in xylenes),¹³ a low yield (<10%) was observed under microwave irradiation for 3 h (entry 8). Furthermore, cyclization is less efficient with other *N*-acetyl-2-

Table 1. One-pot cyclization of *N*-acetyl-2-haloaniline with Woollins' reagent.

Entry	X	Compounds	W. R. (equiv)	Conditions	Yield % ^a
1	I	1a	1.2	THF, 80 °C, 12h	0
2	I	1a	1.2	Toluene, 110 °C, 16 h	0
3	I	1a	1.2	Pyridine, 115 °C, 16 h	< 10 (3a)
4	I	1a	1.2	Xylenes, 140 °C, 16 h	18 (3a)
5	I	1a	1.2	Pyridine, 160 °C, 6 h	MW 27 (3a)
6	I	1a	2.0	Pyridine, 160 °C, 6 h	MW 41 (3a)
7	I	1a	1.2	NEt ₃ , 160 °C, 3 h	MW 57 (3a)
8	I	1a	1.2	Xylenes, CsCO ₃ (2 equiv), 3 h	MW < 10 (3a)
9	Br	1b	1.2	NEt ₃ , 160 °C, 6 h	MW 35 (2b), 19 (3a)
10	Cl	1c	1.2	NEt ₃ , 160 °C, 3 h	MW 49 (2c), <10 (3a)

^a Isolated yield**Table 2.** Scope of the reaction

Entry	Starting material	R	R ¹	Product	Yield % ^a
1	1a	-CH ₃	-H	3a	57
2	1d	-CH ₂ CH ₂ C ₆ H ₅	-H	3d	53
3	1e	-C ₆ H ₅	-H	3e	59
4	1f	- <i>p</i> MeC ₆ H ₄	-H	3f	50
5	1g	- <i>p</i> ClC ₆ H ₄	-H	3g	49
6	1h	- <i>p</i> CF ₃ C ₆ H ₄	-H	3h	48
7	1i	- <i>p</i> OMeC ₆ H ₄	-H	3i	48
8	1j	-2-furanyl	-H	3j	46
9	1k	-2-thienyl	-H	3k	45
10	1l	-2-pyrrolyl	-H	3l	55
11	1m	-CH ₃	-Br	3m	44
12	1n	-CH ₃	-OCH ₃	3n	21

^a Isolated yield

haloanilines with chlorine or bromine atoms. For example, with *N*-acetyl-2-bromoaniline **1b** and *N*-acetyl-2-chloroaniline **1c**, the reaction gave a mixture of corresponding selenamides **2** and 2-methylbenzoselenazole **3a** (entries 9-10).

To extend the substrate scope of this one-pot selenation/cyclization, various *N*-(acetyl)benzoyl-2-iodoanilines were synthesized¹⁶ and subjected to the microwave protocol. The results obtained under the optimized reaction conditions¹⁷ are listed in Table 2 and show that the reaction is efficient with different alkyl, aryl and heteroaryl derivatives. Different functional groups including electron-withdrawing groups such as chloro and trifluoromethyl and electron-donating groups such as methoxy, furan, thiophene and pyrrole on the benzoyl moiety showed roughly the same efficiency. The electron-withdrawing groups such as carbonyl,^{18a} cyano,^{18b}, amide^{18c} and sulfone^{18d} on the benzoyl moiety were not tested because of their well-known good reactivity with Woollins reagent. The selenation/cyclization process was also applied with electron-withdrawing groups or electron-donating groups such as bromo or methoxy in *para* position of iodo. The corresponding yields are similar for the bromo compound **3m** (entry 11) and lower for the methoxy compound **3n** (entry 12).

In summary, we have developed a novel method of benzoselenazole synthesis with Woollins' reagent under basic conditions and microwave irradiation *via* a one-pot reaction from corresponding *N*-(acetyl)benzoyl-2-iodoanilines. To the best of our knowledge, this work describes for the first time a synthetic pathway in benzoselenazole series under microwave irradiation. Under these experimental conditions, the *N*-acetylchloro or bromoanilines led to selenoamide intermediate as the isolated major product and 2-methylselenazole as the isolated minor product. The scope of the reaction included synthesis of *N*-(acetyl)benzoyl-2-iodoanilines succeeded in moderate to good yields.

Acknowledgments

This work was supported by the CNRS (Centre National de la Recherche Scientifique) and Aix-Marseille Université. The authors thank V. Remusat for NMR spectra recording and the Spectropole team for various analytical measurements.

References and notes

- (a) Muges, G.; du Mont, W.-W.; Sies, H. *Chem. Rev.* **2001**, *101*, 2125–2179. b) Nogueira, C. W.; Zeni, G.; Rocha, J. B. T. *Chem. Rev.* **2004**, *104*, 6255–6285.
- Goldstein, B. M.; Takusagawa, F.; Berman, H. W.; Srivastava, P. C.; Robins, R. K. *J. Am. Chem. Soc.* **1985**, *107*, 1394–1400.
- Grange, R. L.; Ziogas, J.; Angus, J. A.; Schiesser, C. H. *Tetrahedron Lett.* **2007**, *48*, 6301–6303.
- Ryu, C.-K.; Han, J.-Y.; Jung, O.-J.; Lee, S.-K.; Lee, J. Y.; Jeong, S. H. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 679–682.
- Doering, M.; Ba, L. A.; Lilienthal, N.; Nicco, C.; Scherer, C.; Abbas, M.; Zada, A. A. P.; Coriat, R.; Burkholz, T.; Wessjohann, L.; Diederich, M.; Batteux, F.; Herling, M.; Jacob, C. *J. Med. Chem.* **2010**, *53*, 6954–6963.
- For recent examples: (a) Reis, L. V.; Serrano, J. P. C.; Almeida, P.; Santos, P. F. *Synlett* **2002**, *10*, 1617–1620. (b) Santos, P. F.; Reis, L. V.; Almeida, P.; Oliveira, A. S.; Vieira Ferreira, L. F. *J. Photochem. Photobiol. A: Chem.* **2003**, *160*, 159–161. (c) Santos, P. F.; Reis, L. V.; Almeida, P.; Serrano, J. P.; Oliveira, A. S.; Vieira Ferreira, L. F. *J. Photochem. Photobiol. A: Chem.* **2004**, *163*, 267–269. (d) Reis, L. V.; Serrano, J. P.; Almeida, P.; Santos, P. F. *Dyes Pigments* **2009**, *81*, 197–202.
- (a) Sharma, B. K.; Muges, G. *J. Am. Chem. Soc.* **2005**, *127*, 11477–11485. (b) Bhabak, K. P.; Muges, G. *Chem. Eur. J.* **2007**, *13*, 4594–4601. (c) Sharma, B. K.; Muges, G. *Chem. Eur. J.* **2008**, *14*, 10603–10614. (d) Bhabak, K. P.; Muges, G. *Chem. Asian J.* **2009**, *4*, 974–983. (e) Sharma, B. K.; Manna, D.; Minoura, M.; Muges, G. *J. Am. Chem. Soc.* **2010**, *132*, 5364–5374.
- (a) Crozet, M. P.; Archaimbault, G.; Vanelle, P.; Nougier, R. *Tetrahedron Lett.* **1985**, *26*, 5133–5134; (b) Curti, C.; Laget, M.; Ortiz Carle, A.; Gellis, A.; Vanelle, P. *Eur. J. Med. Chem.* **2007**, *42*, 880–884; (c) Cohen, A.; Crozet, M. D.; Rathelot, P.; Vanelle, P. *Green Chem.* **2009**, *11*, 1736–1742; (d) Verhaeghe, P.; Azas, N.; Hutter, S.; Castera-Ducros, C.; Laget, M.; Dumetre, A.; Gasquet, M.; Reboul, J.-P.; Rault, S.; Rathelot, P.; Vanelle, P. *Bioorg. Med. Chem.* **2009**, *17*, 4313–4322; (e) Crozet, M. D.; Botta, C.; Gasquet, M.; Curti, C.; Rémusat, V.; Hutter, S.; Chapelle, O.; Azas, N.; De Méo, M.; Vanelle, P. *Eur. J. Med. Chem.* **2009**, *44*, 653–659.
- (a) Bogert, M. T.; Stull, A. *J. Am. Chem. Soc.* **1927**, *49*, 2011–2016; (b) Mbuyi, M.; Evers, M.; Tihange, G.; Luxen, A.; Christiaens, L. *Tetrahedron Lett.* **1983**, *24*, 5873–5876. (c) Radatz, S., C.; Alves, D.; Schneider, P. H. *Tetrahedron* **2013**, *69*, 1316–1321.
- Ramana, T.; Punniyamurthy T. *Eur. J. Org. Chem.* **2011**, 4756–4759.
- Kaname, M.; Minoura, M.; Sashida, H. *Tetrahedron Lett.* **2011**, *52*, 505–508.
- Fujiwara, S.-i.; Asanuma, Y.; Shin-ike, T.; Kambe N. *J. Org. Chem.* **2007**, *72*, 8087–8090.
- Bernardi, D.; Ba, L. A.; Kirsch G. *Synlett* **2007**, *13*, 2121–2123.
- (a) Hua, G.; Woollins, J. D. *Angew. Chem. Int. Ed.* **2009**, *48*, 1368–1377. (b) Madhumita, M.; Sourav, C.; Parasuraman, J. *Synlett* **2012**, *23*, 2615–2618.
- For an example of efficient thionation under microwave irradiation see: Varma, R. S.; Kumar, D. *Org. Lett.* **1999**, *1*, 697–700.
- The substrates **1** were prepared from the acyl chlorides in toluene.
- Typical experimental procedure for synthesis of benzoselenazoles: In a 5 mL glass vial equipped with a small magnetic stirring bar, the *N*-(2-iodophenyl)-acetamide **1a** (0.57 mmol, 150 mg), Woollins' reagent (0.34 mmol, 180 mg, 1.2 equiv) were mixed with dry triethylamine (3 mL) under nitrogen atmosphere. The vial was tightly sealed with an aluminum/teflon crimp top. The mixture was then irradiated in a Biotage® Initiator microwave reactor for 3 h at 160 °C. After the reaction was complete, the reaction mixture was evaporated under *vacuum*. The residue was purified by flash chromatography on silica gel using petroleum ether/EtOAc (95:5) as the eluent, yielding **3a** (65 mg, 57%). Analysis for compounds **3a**, **3e**, **3f** and **3i** are in agreement with those reported in the literature.^{9b,9c} Data for the new compounds:
2-(2-Phenylethyl)-1,3-benzoselenazole (**3d**): White solid (87 mg), yield = 53%; mp 114–116 °C; *R*_f = 0.35 (ethyl acetate/petroleum ether: 5/95); ¹H NMR (200 MHz, CDCl₃, δ, ppm) 7.93 (d, *J* = 8.2 Hz, 1H), 7.79 (d, *J* = 7.9 Hz, 1H), 7.38 (t, *J* = 7.9 Hz, 1H), 7.12–7.27 (m, 6H), 3.39 (t, *J* = 7.9 Hz, 2H), 3.12 (t, *J* = 7.9 Hz, 2H); ¹³C NMR (50 MHz, CDCl₃, δ, ppm) 174.5, 152.0, 138.0, 136.3, 126.7, 126.6, 124.6, 124.3, 123.2, 123.0, 122.1, 37.2, 33.8. LC-MS (ESI+): *t*_R = 4.02 min; *m/z* [M + H]⁺: 288.15. HRMS for C₁₅H₁₄NSe [M + H]⁺ calcd/found: 288.0291/288.0288.
2-(4-Chlorophenyl)-1,3-benzoselenazole (**3g**): White solid (82 mg), yield = 49%, mp 101 °C; *R*_f = 0.35 (ethyl acetate/petroleum ether: 5/95); ¹H NMR (200 MHz, CDCl₃, δ, ppm) 8.12 (d, *J* = 8.1 Hz, 1H), 7.90–8.00 (m, 3H), 7.42–7.55 (m, 3H), 7.33 (dt, *J* = 7.9, 1.2 Hz, 1H); ¹³C NMR (50 MHz, CDCl₃, δ, ppm) 171.2, 155.5, 138.3, 137.3, 134.6, 129.5, 129.3, 126.7, 125.7, 125.0. HRMS for C₁₃H₉ClNSe [M + H]⁺ calcd/found: 293.9581/293.9584.
2-(4-(Trifluoromethyl)phenyl)-1,3-benzoselenazole (**3h**): White solid (90 mg), yield = 48%, mp 143–145 °C; *R*_f = 0.35 (ethyl acetate/petroleum ether: 5/95); ¹H NMR (200 MHz, CDCl₃, δ, ppm) 8.01–8.06 (m, 3H), 7.95 (d, *J* = 8.0 Hz, 1H), 7.75 (d, *J* = 8.4 Hz, 2H), 7.37 (dt, *J* = 8.2 Hz, 1.2 Hz, 1H), 7.32 (dt, *J* = 8.0 Hz, 1.2 Hz, 1H); ¹³C NMR (50 MHz, CDCl₃, δ, ppm) 170.6, 155.7, 139.3, 138.7, 132.5 (q, *J* = 32.5 Hz), 128.3, 126.8, 126.2 (q, *J* = 3.5 Hz), 126.0, 125.4, 125.1, 123.9 (q, *J* = 27.2 Hz). LC-MS (ESI+): *t*_R = 4.77 min; *m/z* [M + H]⁺: 328.05. HRMS for C₁₄H₉F₃NSe [M + H]⁺ calcd/found: 327.9847/327.9850.
2-(Furan-2-yl)-1,3-benzoselenazole (**3j**): White solid (65 mg), yield = 46%, mp 122–124 °C; *R*_f = 0.46 (ethyl acetate/petroleum ether: 10/90); ¹H NMR (200 MHz, CDCl₃, δ, ppm) 8.11 (d, *J* = 8.1, 0.6 Hz, 1H), 7.87 (d, *J* = 7.8, 0.8 Hz, 1H), 7.62 (d, *J* = 1.3 Hz, 1H), 7.49 (dd, *J* = 7.7, 1.2 Hz, 1H), 7.26–7.35 (m, 2), 6.61 (dd, *J* = 3.5, 1.8 Hz, 1H); ¹³C NMR (50 MHz, CDCl₃, δ, ppm) 160.9, 155.1, 151.0, 145.1, 137.4, 126.7, 125.4, 124.9, 124.7, 113.0, 111.4. LC-MS (ESI+): *t*_R = 3.07 min; *m/z* [M + H]⁺: 250.18. HRMS C₁₁H₈NOSe [M + H]⁺ calcd/found: 249.9766 / 249.9766.
2-(Thiophen-2-yl)-1,3-benzoselenazole (**3k**): Yellow solid (68 mg), yield = 45%, mp 107–109 °C; *R*_f = 0.57 (ethyl acetate/petroleum ether: 10/90); ¹H NMR (200 MHz, CDCl₃, δ, ppm) 8.07 (dd, *J* = 8.1 Hz, 1H), 7.87 (dd, *J* = 7.8 Hz, 1H), 7.60 (d, *J* = 3.7 Hz, 1H), 7.42–7.52 (m, 2H), 7.24–7.34 (m, 1H), 7.12 (d, *J* = 5.1, 1.2 Hz, 1H); ¹³C NMR (50 MHz, CDCl₃) δ: 164.5, 155.1, 140.1, 137.9, 129.9, 129.8, 128.2, 126.7, 125.5, 124.8, 124.6. LC-MS (ESI+): *t*_R = 3.50 min; *m/z* [M + H]⁺: 266.06. HRMS C₁₁H₈NSSe [M + H]⁺ calcd/found: 265.9537/265.9536.

- 2-(1*H*-pyrrol-2-yl)-1,3-benzoselenazole (**3l**): White solid (78 mg), yield = 55%; mp 148-150 °C; R_f = 0.31 (ethyl acetate/petroleum ether: 10/90); ^1H NMR (200 MHz, CDCl_3 , δ , ppm) 10.09 (s, 1H), 7.95 (d, J = 8.2 Hz, 1H), 7.91 (d, J = 8.2 Hz, 1H), 7.48 (dt, J = 6.8, 1.1 Hz, 1H), 7.20-7.25 (m, 1H), 7.00-7.05 (m, 1H), 6.80-6.85 (m, 1H), 6.30-6.35 (m, 1H); ^{13}C NMR (50 MHz, CDCl_3 , δ , ppm) 163.8, 155.0, 136.7, 128.9, 126.5, 124.9, 124.7, 123.3, 122.7, 114.4, 110.9. LC-MS (ESI+): t_R = 3.02 min; m/z $[\text{M} + \text{H}]^+$: 249.22. HRMS $\text{C}_{11}\text{H}_9\text{N}_2\text{Se}$ $[\text{M} + \text{H}]^+$ calcd/found: 248.9926/248.9925.
- 5-Bromo-2-methyl-benzoselenazole (**3m**): White solid (70 mg), yield = 44%; mp 148-150 °C; R_f = 0.25 (ethyl acetate/petroleum ether: 10/90); ^1H NMR (200 MHz, CDCl_3 , δ , ppm) 8.11 (d, J = 1.8 Hz, 1H), 7.70 (d, J = 8.4 Hz, 1H), 7.39 (dd, J = 8.6, 1.8 Hz, 1H), 2.87 (s, 3H); ^{13}C NMR (50 MHz, CDCl_3 , δ , ppm) 173.6, 155.6, 137.7, 128.0, 126.8, 125.8, 119.7, 23.8. LC-MS (ESI+): t_R = 2.96 min; m/z $[\text{M} + \text{H}]^+$: 275.92. HRMS $\text{C}_8\text{H}_7\text{BrNSe}$ $[\text{M} + \text{H}]^+$ calcd / found: 275.8919/275.8918.
- 5-Methoxy-2-methyl-benzoselenazole (**3n**): Orange solid (25 mg), yield = 21%; mp 88-90 °C; R_f = 0.23 (ethyl acetate/petroleum ether: 10/90); ^1H NMR (200 MHz, CDCl_3 , δ , ppm) 7.68 (d, J = 8.5 Hz, 1H), 7.51 (d, J = 2.4 Hz, 1H), 6.92 (dd, J = 9.3, 2.1 Hz, 1H), 3.86 (s, 3H), 2.84 (s, 3H); ^{13}C NMR (50 MHz, CDCl_3 , δ , ppm) 171.3, 157.0, 128.0, 123.0, 114.5, 112.6, 105.2, 53.7, 27.8. LC-MS (ESI+): t_R = 2.34 min; m/z $[\text{M} + \text{H}]^+$: 228.19. HRMS $\text{C}_9\text{H}_{10}\text{NOSe}$ $[\text{M} + \text{H}]^+$ calcd / found: 227.9928/227.9923.
18. (a) Bhattacharyya, P.; Woollins, J. D. *Tetrahedron Lett.* **2001**, *42*, 5949–5951. (b) Hua, G.; Li, Y.; Slawin, A. M. Z.; Woollins, J. D. *Org. Lett.* **2006**, *8*, 5251–5254. (c) Bethke, J.; Karaghiosoff, K.; Wessjohann, L. A. *Tetrahedron Lett.* **2003**, *44*, 6911–6913. (d) Hua, G.; Woollins, J. D. *Tetrahedron Lett.* **2007**, *48*, 3677–3679.

Supplementary Material

Supplementary data associated with this article can be found, in the online version.