



The history of Greenland's ice: Cosmic signature

Pierre-Henri Blard, Guillaume Leduc

► **To cite this version:**

Pierre-Henri Blard, Guillaume Leduc. The history of Greenland's ice: Cosmic signature. Nature, Nature Publishing Group, 2016, 10.1038/540202a . hal-01457254

HAL Id: hal-01457254

<https://hal-amu.archives-ouvertes.fr/hal-01457254>

Submitted on 20 Feb 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The history of Greenland's ice

Global sea levels would rise by several metres if the Greenland Ice Sheet melted completely. Two studies have examined its past behaviour in an effort to evaluate its vulnerability in a warming world — and have come to seemingly conflicting conclusions. Two geochemists and a glaciologist discuss the issues. [SEE LETTERS P.252 & P.256](#)

THE PAPERS IN BRIEF

- Knowledge of the ancient history of ice sheets is needed to inform predictions of their future response to climate change.
- On page 252, Schaefer *et al.*¹ present measurements of beryllium-10 (¹⁰Be) and aluminium-26 (²⁶Al) from bedrock beneath the Greenland Ice Sheet.
- From their data, they propose that Greenland has undergone one or more

episodes of deglaciation during the past 1.1 million years.

- On page 256, Bierman *et al.*² present ¹⁰Be and ²⁶Al data from marine sediment cores collected off the coast of Greenland.
- They conclude that the East Greenland Ice Sheet remained present during the Pleistocene (the epoch that lasted from 2.6 million to 11,700 years ago), but grew and shrank dynamically in response to climate.

Cosmic signature

PIERRE-HENRI BLARD & GUILLAUME LEDUC

It is difficult to find evidence for past waxing and waning of the Greenland Ice Sheet (GrIS), because the glacial debris that can provide a record of ice-sheet dynamics (Fig. 1) is erased each time glaciers advance after a period of deglaciation. Schaefer *et al.* and Bierman *et al.* now report the presence of two rare nuclides (¹⁰Be and ²⁶Al) in Greenland's geological archives. These 'cosmogenic' nuclides are produced by the interaction between atoms in minerals and high-energy cosmic neutrons that probably originated in supernovae, and they provide much-needed evidence about the GrIS's ancient past.

A key characteristic of these isotopes is that they are produced in detectable quantities in only the first few metres below Earth's surface. If they become buried below hundreds of metres of ice, their concentrations decrease through radioactive decay. The half-lives of ¹⁰Be and ²⁶Al are 1.4 million and 0.7 million years, respectively, so the presence of these nuclides in bedrock under ice sheets is direct evidence that one or several ice-free episodes occurred in the recent past (less than 5 million years ago).

Bierman *et al.* report variations in ¹⁰Be and ²⁶Al from marine sediments deposited near the eastern margin of today's GrIS over the past 7.5 million years. The low ¹⁰Be concentrations suggest that the sedimentary material originates from locations that were efficiently

shielded from cosmic rays. The authors therefore conclude that large ice caps probably existed continuously across East Greenland — the source of the sedimentary material — during the past 7.5 million years. They also observe an additional long-term decrease in ¹⁰Be and ²⁶Al concentrations during this period, which indicates a gradual

increase in the volume of the GrIS.

In parallel, Schaefer *et al.* analysed bedrock samples recovered from beneath several kilometres of ice in the centre of the present-day GrIS. These rocks cannot have been transported to their current position from elsewhere. The presence of ¹⁰Be and ²⁶Al in these samples is therefore direct evidence for past deglaciation episodes at this location.

Schaefer and colleagues used the observed concentrations and ratios of ¹⁰Be and ²⁶Al in a numerical model to explore plausible glaciation–deglaciation scenarios. They firmly conclude that the sampled bedrock must have been ice-free for at least one episode of about 280,000 years during the past 1.1 million years. Other glaciation–deglaciation scenarios are more plausible, however, such as several shorter disappearances (several thousand years) of the GrIS during the warmest interglacial periods of the Pleistocene³.

The deglaciation proposed by Schaefer



NICK COBBING/GREENPEACE

Figure 1 | Ice-sheet dynamics. These icebergs calved from the Helheim Glacier, an outlet glacier at the periphery of the Greenland Ice Sheet. Such glaciers are responding rapidly to climate change.

et al. seems at odds with Bierman and colleagues' conclusions. But small ice caps might have persisted on the East Greenland highlands during such deglaciated episodes, and could be the origin of the ^{10}Be -poor material observed in Bierman and co-workers' marine sediments. Indeed, modelling experiments⁴ estimate that ice caps 1,000 metres thick might persist on the south-eastern Greenland highlands in the future even if 95% of the GrIS disappears (Fig. 2).

Could the GrIS undergo similar deglaciations in the future? Average global warming of just 2 °C compared with preindustrial temperatures could be enough to melt more than 75% of the GrIS³. If this high sensitivity to warming is confirmed, then the current worst-case scenarios⁵ for future sea-level rise associated with anthropogenic global warming will need to be revised.

Pierre-Henri Blard is at the Centre de Recherches Pétrographiques et Géo-chimiques, CNRS, Université de Lorraine, 54500 Vandoeuvre-lès-Nancy, France. **Guillaume Leduc** is at the Centre Européen de Recherche et d'Enseignement des Géosciences de l'Environnement, (Aix-Marseille University, CNRS, IRD), Europôle Méditerranéen de l'Arbois, 13545 Aix en Provence, France. e-mails: blard@crpg.cnrs-nancy.fr; leduc@cerege.fr