

MUS-2, a novel variant of the chromosome-encoded β -lactamase MUS-1, from *Myroides odoratimimus*

Charbel Al Bayssari, S. Kumar Gupta, Fouad Dabboussi, M Hamze,
Jean-Marc Rolain

► **To cite this version:**

Charbel Al Bayssari, S. Kumar Gupta, Fouad Dabboussi, M Hamze, Jean-Marc Rolain. MUS-2, a novel variant of the chromosome-encoded β -lactamase MUS-1, from *Myroides odoratimimus*. *New Microbes and New Infections*, Wiley Online Library 2015, 7, pp.67-71. 10.1016/j.nmni.2015.06.007 . hal-01772724

HAL Id: hal-01772724

<https://hal-amu.archives-ouvertes.fr/hal-01772724>

Submitted on 20 Apr 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



MUS-2, a novel variant of the chromosome-encoded β -lactamase MUS-I, from *Myroides odoratimimus*

C. Al-Bayssari^{1,2}, S. Kumar Gupta², F. Dabboussi¹, M. Hamze¹ and J.-M. Rolain²

1) Laboratoire Microbiologie Santé et Environnement (LMSE), Centre Azm pour la Recherche en Biotechnologie et ses Applications, Ecole Doctorale de Sciences et de Technologie, Université Libanaise, Tripoli, Libya and 2) Aix-Marseille Université, Unité de Recherche sur les Maladies Infectieuses et Tropicales Emergentes (URMITE), IHU Méditerranée Infection, Faculté de Médecine et de Pharmacie, Marseille, France

Abstract

The aim of the present study was to investigate the molecular mechanism of carbapenem resistance of three imipenem-resistant isolates of *Myroides odoratimimus* recovered from two livestock farms of cows and pigeons by rectal swab in Lebanon in January 2014. Investigation of imipenem resistance of these isolates using the modified Hodge test, the EDTA test, the modified CarbaNP test and the matrix-assisted laser desorption/ionization time-of-flight mass spectrometry Ultraflex assay showed a carbapenemase activity due to the presence of a chromosome-encoded β -lactamase MUS, verified by PCR. However amplification and sequencing of this chromosomal gene showed a novel variant of it designated MUS-2 by the curators of the Lahey database of β -lactamases (<http://www.lahey.org/Studies/webt.asp>). Cloning of the *bla*_{MUS-2} was performed, followed by protein expression in *Escherichia coli* TOP 10. Pulsed-field gel electrophoresis clearly showed that the three isolates belonged to the same clone. This study reports a novel variant of the chromosome-encoded *bla*_{MUS-1} associated with carbapenem resistance in *Myroides odoratimimus* and shows that animals may represent a reservoir of bacteria harbouring several variants of resistance genes.

New Microbes and New Infections © 2015 The Authors. Published by Elsevier Ltd on behalf of European Society of Clinical Microbiology and Infectious Diseases.

Keywords: Animals, *bla*_{MUS-2}, carbapenem-resistance, metallo- β -lactamase, *Myroides odoratimimus*

Original Submission: 8 April 2015; **Accepted:** 22 June 2015

Available online 27 June 2015

Corresponding author: J.-M. Rolain, Aix-Marseille Université, Unité de Recherche sur les Maladies Infectieuses et Tropicales Emergentes (URMITE), UM63, CNRS 7278, IRD 198, Inserm 1095, IHU Méditerranée Infection, Faculté de Médecine et de Pharmacie, 27 Bd Jean Moulin, 13005 Marseille, France
E-mail: jean-marc.rolain@univ-amu.fr

Introduction

Myroides species, previously named as *Flavobacterium odoratum*, are aerobic Gram-negative bacteria that can be found in environmental sources such as soil and water, but are not residents of the normal human microflora [1]. The genus *Myroides* comprises two species, *Myroides odoratus* and *Myroides odoratimimus* [2]. Taking into consideration their role as causative agents in human disease, both *M. odoratimimus* and *M. odoratus*

are seen rarely as opportunistic pathogens, causing infections in severely immunocompromised patients and rarely, in immunocompetent hosts [1,3]. *Myroides odoratimimus* strains have been found to be responsible for urinary tract infections [4,5], cellulitis [6,7] in immunocompromised patients, and also for septic shock, pneumonia and soft-tissue infections [1,8] in immunocompetent hosts. In addition, three outbreaks of *M. odoratimimus*, two of urinary tract infections and the third of central venous catheter-associated bloodstream infections due to contaminated water, have been reported [2,4].

Antibiotic resistance patterns of *Myroides* strains show variable susceptibility to β -lactams [5], with a conserved decreased susceptibility to cephalosporins and carbapenems [9]. This decreased susceptibility is the result of the production of a chromosome-encoded β -lactamase MUS-I. This enzyme is a member of the subclass BI of metallo- β -lactamases and is distantly related to other metalloenzymes, being most closely

related to IND-1 from *Chryseobacterium indologenes* (42% amino acid identity). However, phylogenetic analysis showed that MUS-1 belongs to the same phylogenetic lineage of subclass B1 enzymes that groups the subclass B1 β -lactamases of *Flavobacterium* species [9]. Here we report a novel variant, MUS-2, of the chromosome-encoded β -lactamase MUS-1 from *M. odoratimimus* isolated from livestock animals in Lebanon.

Materials and methods

Bacterial isolates

Three imipenem-resistant *M. odoratimimus* strains (32, 35a and 104b) were isolated from two livestock farms of cows and pigeons by rectal swabs (32: farm 1, cow; 35a: farm 1, pigeon; 104b: farm 2, cow) in January 2014. These strains were collected during a study that aimed to detect carbapenemase-encoding genes in animals in Lebanon [10]. They were isolated on MacConkey agar supplemented by ertapenem (1 mg/L), from alive and non-sick animals without contact with any other animals. Strains were sub-cultured on trypticase soy agar plates at 37°C for 18–24 h and identification was confirmed by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF/MS) (Microflex™; Bruker Daltonics, Bremen, Germany) with FLEXCONTROL software (Bruker Daltonics).

Antibiotic susceptibility testing

Antibiotic susceptibility testing was determined on Müller–Hinton agar by standard disc diffusion method as recommended by the European Committee on Antimicrobial Susceptibility Testing (EUCAST; www.eucastr.org). Seventeen antibiotics were tested, including ticarcillin, ticarcillin-clavulanic acid, piperacillin, piperacillin-tazobactam, ceftazidime, cefotaxime, cefepime, aztreonam, amikacin, tobramycin, gentamicin, ciprofloxacin, rifampicin, ertapenem, meropenem, imipenem and colistin (Bio-Rad, Marnes-la-Coquette, France). The MIC for imipenem was determined using the Etest method (bioMérieux, La Balmes les Grottes, France) and the result was interpreted according to the EUCAST breakpoint for *Enterobacteriaceae* (susceptible if MIC \leq 2 mg/L)

Phenotypic and molecular detection of carbapenemases

Isolates were screened for carbapenemase production using the modified Hodge test, the EDTA, the Modified CarbaNP test as previously described [11–13] and also by MALDI-TOF MS Ultraflex assay [14]. Screening for class A, B and D carbapenemases was carried out using PCR according to previous protocols and previously described primers including for *bla*_{VIM}, *bla*_{IMP}, *bla*_{NDM-1}, *bla*_{KPC} and all the *bla*_{OXA} variants [15–20]. In

addition, amplification of the chromosomal MUS gene was performed using primers designed in this study (MUS-F/R 5'-CGTTAGGCACACCAGAAGAA-3'/3'-TCCACACCATTCT-GAGCGTA-5'). These primers were designed outside the region containing the *bla*_{MUS-1} located on the chromosome of a random *M. odoratimimus* genome deposited in GenBank. PCR products were purified and sequenced using the Big dye terminator chemistry on an ABI 3730 automated sequencer (Applied Biosystems, Foster City, CA, USA). The sequences obtained were analysed using BLASTN and BLASTP against the NCBI database (www.ncbi.nlm.nih.gov).

Cloning of the MUS-2 gene

The full length of the *bla*_{MUS-2} from *M. odoratimimus* was amplified using *Herculase II Fusion Enzyme* with *dNTPs Combo* (Agilent Technologies, Santa Clara, CA, USA) with primer MUS_F/MUS_R. The gene was then cloned into the high-copy-number plasmid pCR-BluntII-TOPO using the Zero Blunt TOPO PCR cloning kit (Invitrogen, Carlsbad, CA, USA). The resulting plasmid pTOPO-MUS-2 was transformed into TOP10 Electrocomp™ *Escherichia coli*. Transformants were selected on Luria–Bertani agar supplemented with 100 mg/L of kanamycin; the presence of the insert was confirmed by PCR. The antibiotic resistance profile of the TOP10 *E. coli* containing pTOPO-MUS vector was then determined.

Molecular strain typing

The three isolates were typed by the pulsed-field gel electrophoresis (PFGE) method as previously described [4]. The DNA was digested with *Sma*I restriction enzyme and the fragments of DNA were separated using a GenePath system (Bio-Rad). The running conditions were 6 V/cm at 14°C for 20 h.

Results

During a previous study that was aiming to detect carbapenemase encoding genes in animals in livestock farms in Lebanon [10], three isolates of *M. odoratimimus* were isolated and were studied separately. These isolates were obtained after culture of animal faeces in MacConkey agar supplemented with ertapenem (1 mg/L) and were identified by MALDI-TOF MS as *M. odoratimimus* (score: 2.439). The results of antibiotic susceptibility testing revealed that the isolates were resistant to almost all antibiotics, including β -lactams, aminoglycosides and colistin but remain susceptible to sulfamethoxazole-trimethoprim, rifampicin and fluoroquinolones (Table 1). In addition, all the isolates showed resistance to carbapenems with MIC for imipenem at 8 mg/L. The isolates were then analysed via the modified Hodge test, the MALDI-TOF MS Ultraflex

TABLE 1. Antimicrobial susceptibility testing of the three *Myroides* isolates

	TIC	PRL	TIM	TPZ	CAZ	FEP	CTX	ETP	MEM	IPM	TOB	AMK	GEN	CIP	ATM	RA	SXT	CT
MO_32	R	R	R	R	R	R	R	R	R	R	R	R	R	I	R	S	S	R
MO_35a	R	R	R	R	R	R	R	R	R	R	R	R	R	I	R	S	S	R
MO_104b	R	R	R	R	R	R	R	R	R	R	R	R	R	I	R	S	S	R

MO, *Myroides odoratimimus*; TIC, ticarcillin; PRL, piperacillin; TIM, ticarcillin-clavulanic acid; TPZ, piperacillin-Tazobactam; CAZ, ceftazidime; FEP, cefepime; CTX, cefotaxime; ETP, ertapenem; MEM, meropenem; IPM, imipenem; TOB, tobramycin; AMK, amikacin; GEN, gentamicin; CIP, ciprofloxacin; ATM, aztreonam; RA, rifampicin; SXT, sulfamethoxazole-trimethoprim; CT, colistin; R, resistant; I, intermediate; S, susceptible.




FIG. 1. Amino acid sequence alignment between the MUS-1 and MUS-2 proteins.

assay, the EDTA-disc test and the modified CarbaNP test and they were all positive on the modified Hodge test, EDTA-disc and modified CarbaNP test positive. In addition, the MALDI-TOF MS Ultraflex assay was positive for all of them—suggesting a carbapenemase production, more specifically a metallo-β-lactamase as the EDTA test was positive. Screening for carbapenemase-encoding genes by PCR using carbapenemase gene primers showed a negative result for *bla_{VIM}*, *bla_{IMP}*, *bla_{NDM-1}*, *bla_{KPC}* and all the *bla_{OXA}* variants. However the chromosomal MUS gene was amplified using primers MUS-F/R.

Sequencing of the chromosomal MUS gene of the three isolates and nucleotide alignment between the *bla_{MUS-1}* and the new *bla_{MUS-2}* sequence showed some differences. The *bla_{MUS-2}* differed from *bla_{MUS-1}* due to substitution in nucleotides at positions 69 (A → G), 525 (A → G), 171 (C → T), 177 (C → T), 397 (C → T), 671 (C → T), 179 (T → C), 514 (T → C) and 675 (C → G), resulting in a different protein MUS-2 that differ from MUS-1 protein due to substitution of amino acid at position 60 (Valine → Alanine), 133 (Histidine → Tyrosine) and 224 (Threonine → Isoleucine) (Fig. 1). The percentage of homology between the two amino acid sequences was 98.78.

The new enzyme variant was designated MUS-2 by the curators of the Lahey database of β-lactamases (<http://www.lahey.org/Studies/webt.asp>) and deposited in GenBank under accession number KP658209.

Cloning of the *Bla_{MUS-2}* successfully gave plasmids harbouring this gene; these plasmids were transformed into TOPI0 Electrocomp™ *E. coli*. The presence of the insert was verified by PCR. MIC values of β-lactams for the *E. coli* TOPI0 strain, which harbours recombinant plasmid pTOPO-MUS-2, showed

that the bacteria was resistant to amoxicillin and ticarcillin and had a reduced susceptibility to piperacillin, in addition it showed an increased resistance to extended-spectrum cephalosporins and carbapenems by at least four-fold of MIC (Table 2).

Finally, PFGE analysis showed that the three strains of *M. odoratimimus* were genetically clonal since their PFGE profiles were similar (Fig. 2).


Discussion

To date, infections caused by *M. odoratimimus* in immunocompromised and immunocompetent hosts had been rarely

TABLE 2. MICs of β-lactams for *Myroides odoratimimus* 35a, TOPI0 *Escherichia coli* and TOPI0 *E. coli* pTOPO-MUS-2

Substrate ^a	MIC (µg/mL)		
	<i>M. odoratimimus</i> 35a	TOPI0 <i>E. coli</i> pTOPO-MUS-2	TOPI0 <i>E. coli</i>
Amoxicillin	32	128	2
Amoxicillin-CLA	32	128	2
Ticarcillin	256	256	2
Piperacillin	128	8	2
Piperacillin-TZB	24	8	1
Cephalotin	128	8	2
Cefoxitin	32	4	2
Moxalactam	32	0.064	0.19
Cefotaxime	32	0.094	0.047
Cefuroxime	128	6	1
Ceftazidime	128	0.38	0.064
Cefepime	32	0.064	0.064
Cefpirone	32	0.064	0.064
Aztreonam	64	0.064	0.064
Imipenem	8	0.38	0.064
Meropenem	4	0.38	0.064

^aCLA, clavulanic acid at 2 µg/mL; TZB, tazobactam at 4 µg/mL.



Line 1: *M. odoratimimus* 32
 Line 2: *M. odoratimimus* 35a
 Line 3: *M. odoratimimus* 104b
 MM: Molecular marker

FIG. 2. Pulsed-field gel electrophoresis of *Smal* DNA restriction digests of the three *Myroides odoratimimus* isolates.

reported [1,2,4–8]. The fact that this bacterium might cause infection in immunocompetent hosts is a real health problem because it is resistant to β -lactams including carbapenems, aminoglycoside but also to colistin. Biochemical detection of β -lactamases in clinical strains of *Myroides* spp. (formerly *F. odoratum*) was performed in 1985 by Sato et al. [21], but the molecular characteristics of β -lactamase were not reported. We would have to wait until 2002 when Mammeri et al. biochemically and genetically characterized β -lactamases expressed in *M. odoratimimus* and *M. odoratus* [9]. In fact, *M. odoratimimus* species produce chromosomal MUS-I β -lactamase, which belongs to the subclass B1 of Ambler classification, and their antibiotic resistance patterns show a constant decreased susceptibility to imipenem [9]. The β -lactamase produced by the three *M. odoratimimus* isolates in our study shared 98.78% amino acid identity with the known MUS-I protein. At the present time, each bacterial species of the *Flavobacteriaceae* family that has been investigated for β -lactamase characterization produces a subclass B1 metalloenzyme [9]. Our study confirms a previous study performed on *M. odoratimimus* species; however, here we report a new variant of this enzyme in Lebanon. PFGE analysis showed that these isolates belong to the same clone because they had the same PFGE profile. This confirms that contamination between farms has occurred and precautions should be established to limit the emergence of this clone in livestock farms because that could be a source of human infection.

Taking into consideration the biochemical criteria established by Rasmussen and Bush to classify metallo- β -lactamases [22], MUS-I and most probably MUS-2 belong to functional subgroup 3a, as their catalytic efficiencies for penicillins are at least 60% of that for imipenem [9]. However, these data are discrepant with those from Rasmussen and Bush [22], who

mentioned that a metallo- β -lactamase produced by *F. odoratum* strain belongs to subgroup 3b, which groups true carbapenem-hydrolysing β -lactamases. Regarding this conflict, it is very difficult to estimate the exact role of either MUS-1 or MUS-2 in the intrinsic β -lactam resistance of *M. odoratimimus* because the metalloenzymes expressed in *E. coli* give much lower levels of resistance to β -lactams than those seen in the original producers as seen in the TOP10 *E. coli* after cloning.

Finally, it remains unknown if these bacteria could be a source of such a variety of metalloenzymes. The most serious finding is that animals could be a reservoir for such genes and are a source of new and/or emerging multidrug-resistant bacteria and due to possible links with animals, humans may have high risk factors for colonization/infection with such bacteria.

In conclusion, we report a novel variant of the chromosome-encoded β -lactamase MUS-I that has been designated MUS-2 from *M. odoratimimus* isolates from livestock animals in Lebanon.

GenBank accession number

The full sequence of the Bla_{MUS-2} described in this study is deposited in the GenBank database with accession no. KP658209.

Funding

This work was supported by AZM research centre for biotechnology and its application, Lebanese University, the National Council for Scientific Research, Lebanon, the IHU Méditerranée Infection and the French CNRS.

Transparency declarations

None to declare; all authors have read and approved the manuscript.

Conflict of Interest

None declared.

Acknowledgements

The authors thank Linda Hadjadj, Said Azza, Taha Abdo and Maryam Yehya for technical assistance.

References

- [1] Maraki S, Sarchianaki E, Barbagadakis S. *Myroides odoratimimus* soft tissue infection in an immunocompetent child following a pig bite: case report and literature review. *Braz J Infect Dis* 2012;16:390–2.
- [2] Ktari S, Mnif B, Koubaa M, et al. Nosocomial outbreak of *Myroides odoratimimus* urinary tract infection in a Tunisian hospital. *J Hosp Infect* 2012;80:77–81.
- [3] Schröttner P, Rudolph WW, Eing BR, Bertram S, Gunzer F. Comparison of VITEK2, MALDI-TOF MS, and 16S rDNA sequencing for identification of *Myroides odoratus* and *Myroides odoratimimus*. *Diagn Microbiol Infect Dis* 2014;79:155–9.
- [4] Yagci A, Cerikcioglu N, Kaufmann ME, et al. Molecular typing of *Myroides odoratimimus* (*Flavobacterium odoratum*) urinary tract infections in a Turkish hospital. *Eur J Clin Microbiol Infect Dis* 2000;19:731–2.
- [5] Holmes B, Snell JJ, Lapage SP. *Flavobacterium odoratum*: a species resistant to a wide range of antimicrobial agents. *J Clin Pathol* 1979;32:73–7.
- [6] Bachmeyer C, Entressengle H, Khosrotehrani K, et al. Cellulitis due to *Myroides odoratimimus* in a patient with alcoholic cirrhosis. *Clin Exp Dermatol* 2008;33:97–8.
- [7] Bachman KH, Sewell DL, Strausbaugh LJ. Recurrent cellulitis and bacteremia caused by *Flavobacterium odoratum*. *Clin Infect Dis* 1996;22:1112–3.
- [8] Benedetti P, Rattu M, Pavan G, Sefton A, Pellizzer G. Septic shock, pneumonia, and soft tissue infection due to *Myroides odoratimimus*: report of a case and review of *Myroides* infections. *Infection* 2011;39:161–5.
- [9] Mammeri H, Bellais S, Nordmann P. Chromosome-encoded β -lactamases TUS-I and MUS-I from *Myroides odoratus* and *Myroides odoratimimus* (formerly *Flavobacterium odoratum*), new members of the lineage of molecular subclass B1 metalloenzymes. *Antimicrob Agents Chemother* 2002;46:3561–7.
- [10] Al Bayssari C, Dabboussi F, Hamze M, Rolain JM. Emergence of carbapenemase-producing *Pseudomonas aeruginosa* and *Acinetobacter baumannii* in livestock animals in Lebanon. *J Antimicrob Chemother* 2015;70:950–1.
- [11] Lee K, Lim YS, Yong D, Yum JH, Chong Y. Evaluation of the Hodge test and the imipenem-EDTA double-disk synergy test for differentiating metallo- β -lactamase-producing isolates of *Pseudomonas* spp. and *Acinetobacter* spp. *J Clin Microbiol* 2003;41:4623–9.
- [12] Dortet L, Poirel L, Errera C, Nordmann P. CarbAcineto NP test for rapid detection of carbapenemase-producing *Acinetobacter* spp. *J Clin Microbiol* 2014;52:2359–64.
- [13] Dortet L, Poirel L, Nordmann P. Rapid identification of carbapenemase types in *Enterobacteriaceae* and *Pseudomonas* spp. by using a biochemical test. *Antimicrob Agents Chemother* 2012;56:6437–40.
- [14] Kempf M, Bakour S, Flaudrops C, et al. Rapid detection of carbapenem resistance in *Acinetobacter baumannii* using matrix-assisted laser desorption ionization-time of flight mass spectrometry. *PLoS One* 2012;7:e31676.
- [15] Mesli E, Berrazeg M, Drissi M, Bekkhoucha SN, Rolain JM. Prevalence of carbapenemase-encoding genes including New Delhi metallo- β -lactamase in *Acinetobacter* species, Algeria. *Int J Infect Dis* 2013;17:e739–43.
- [16] Poirel L, Naas T, Nordmann P. Diversity, epidemiology, and genetics of class D β -lactamases. *Antimicrob Agents Chemother* 2010;54:24–38.
- [17] Ruiz M, Marti S, Fernandez-Cuenca F, Pascual A, Vila J. High prevalence of carbapenem-hydrolysing oxacillinases in epidemiologically related and unrelated *Acinetobacter baumannii* clinical isolates in Spain. *Clin Microbiol Infect* 2007;13:1192–8.
- [18] El-Ageery SM, Al-Hazmi SS. Microbiological and molecular detection of VIM-I metallo β lactamase-producing *Acinetobacter baumannii*. *Eur Rev Med Pharmacol Sci* 2014;18:965–70.
- [19] Kusradze I, Diene SM, Goderdzishvili M, Rolain JM. Molecular detection of OXA carbapenemase genes in multidrug-resistant *Acinetobacter baumannii* isolates from Iraq and Georgia. *Int J Antimicrob Agents* 2011;38:164–8.
- [20] Amudhan SM, Sekar U, Arunagiri K, Sekar B. OXA β -lactamase-mediated carbapenem resistance in *Acinetobacter baumannii*. *Indian J Med Microbiol* 2011;29:269–74.
- [21] Sato K, Fujii T, Okamoto R, Inoue M, Mitsuhashi S. Biochemical properties of β -lactamase produced by *Flavobacterium odoratum*. *Antimicrob Agents Chemother* 1985;27:612–4.
- [22] Rasmussen BA, Bush K. Carbapenem-hydrolyzing β -lactamases. *Antimicrob Agents Chemother* 1997;41:223–32.