

Prospective association between consumption frequency of organic food and body weight change, risk of overweight or obesity: Results from the NutriNet-Santé Study

Emmanuelle Kesse-Guyot^{1*}, Julia Baudry¹, Karen E Assmann¹, Pilar Galan¹, Serge Hercberg^{1,2}, Denis Lairon³

¹Sorbonne Paris Cité Epidemiology and Statistics Research Center (CRESS), Inserm U1153, Inra U1125, Cnam, Paris 13 University, Nutritional Epidemiology Research Team (EREN), F-93017 Bobigny, France ;

²Département de Santé Publique, Hôpital Avicenne, F-93017 Bobigny, France ;

³Aix Marseille Université, NORT ; Insem, UMR S 1062 ; Inra 1260, F-13385 Marseille, France

***Correspondence:** Email: e.kesse@eren.smbh.univ-paris13.fr

Equipe de Recherche en Epidémiologie Nutritionnelle (EREN)

SMBH Université Paris 13

74 rue Marcel Cachin, 93017 Bobigny, France

Running title: Organic food consumption and corpulence

Keywords: organic food, obesity, prospective cohort study, weight gain

Abbreviations: BMI: body mass index, OS: organic score, mPNNS-GS, PCA: Principal component analysis

Abstract

A lower body mass index (BMI) has been reported among consumers of organic foods but this relationship has never been examined in a prospective design. Our aim was to prospectively investigate the association between the frequency of organic foods and weight change. We analyzed data from 62,224 participants of the NutriNet-Santé cohort (78% women, mean age=45y) with information on consumption frequency of organic foods, dietary intake, and repeated anthropometric data. For 16 products, participants reported their consumption frequency of labeled organic foods (never, occasionally, most of the time). An organic score (OS) with a maximum of 32 points was computed. The associations of the OS (modeled as quartiles (Q)) with change in BMI during follow-up (on average 3·1 y), and with the risk of overweight and obesity were estimated ANCOVA and multivariable logistic regression. A lower BMI increase was observed across quartiles of the OS (mean difference (95%CI) Q4 versus Q1= - 0·16 (-0·32, -0·01). An increase in OS was associated with a lower risk of overweight and obesity (among non-overweight and non-obese participants at inclusion): odds ratios (95%CI) Q4 versus Q1 were 0·77 (0·68, 0·86) and 0·69 (0·58, 0·82), respectively. Concerning obesity risk, the association was stronger among participants with higher adherence to nutritional guidelines. This study supports a strong protective role of consumption frequency of organic foods concerning the risk of overweight and obesity that depends on overall dietary quality. Upon confirmation, these results may contribute to fine-tune nutritional guidelines by accounting for farming practices in food production.

Introduction

The global obesity epidemic, involving multifaceted origins, is a major public health issue ⁽¹⁾.

Overweight and obesity contribute to the global burden of chronic diseases ⁽²⁾ - especially

type 2 diabetes and ischemic cardiovascular diseases ⁽³⁾, along with cancers ^(4,5).

Among the factors involved in the etiology of obesity, excess caloric consumption, sedentary

lifestyle and genetic susceptibility are well-recognized, but there is a growing concern for the

identification of novel factors involved such as gut microbiota ⁽⁶⁾ or environmental chemicals

⁽⁷⁻¹⁰⁾.

Among the different dietary factors, organic food is of major interest since it presents multiple

features that could potentially protect from weight gain and obesity. Notably, compared to

conventional food, organic food has been suggested to present better nutritional values

concerning fatty acids profiles and specific micronutrients ⁽¹¹⁻¹³⁾. Nonetheless, the possible

implications at an individual level in terms of daily nutrient intake are unknown due to the

lack of food composition tables accounting for farming practices. A small number of clinical

studies have been conducted that compared specific nutritional biomarkers according to the

type of diet (organic or conventional). However, due to short study durations, these studies

were not well-equipped to provide reliable results, and findings were inconsistent ⁽¹⁴⁾.

Organic foods are also characterized by a markedly low level or an absence of pesticide

residues, as repeatedly reported in food residue analyses ^(11,14,15) and in experimental studies

showing that adopting an organic diet leads to a drastic reduction in pesticide residues and

urine metabolites in children and adults ⁽¹⁶⁻²⁰⁾. For most pesticide families (organochlorines

(now banned in the EU but still persistent), organophosphates, and pyrethrynoïds), a large

number of molecules have been recognized as endocrine disruptors ⁽²¹⁾, leading to possible

metabolic disorders ⁽²²⁾. Indeed, a higher exposure to some of these compounds has been

associated with a higher risk for obesity or type 2 diabetes in humans ⁽⁹⁾.

In this context, consumption of organic foods might contribute to the management of weight gain and of obesity risk. In a recent cross-sectional analysis based on the NutriNet-Santé cohort, we showed that participants identified as regular consumers of organic food, compared to non-consumers, presented reduced odds of being overweight or obese (-36% and -62% in men and -42% and -48% in women, respectively) ⁽²³⁾. Comparable findings were reported in the German National Nutrition Survey II (NVS II), a nationwide food consumption study conducted among 13,074 adults: German buyers of organic food exhibited healthier lifestyles compared with non-buyers, and presented lower body weight ⁽²⁴⁾. To the best of our knowledge, no epidemiologic study has yet investigated the prospective relation between consumption frequency of organic foods and the risk of overweight and obesity. The main objective of the present study was thus to investigate the longitudinal association between the frequency consumption of organic foods and change in body mass index, the risk of overweight, and the risk of obesity in a very large adult cohort. Given our previous finding that organic food consumers show a higher level of adherence to nutritional recommendations ⁽²³⁾ - especially a higher consumption of fruit and vegetables and a lower consumption of animal products - a secondary objective was to explore a potential modifying effect of the nutritional quality of the diet on the investigated associations.

Material and methods

Study population

The NutriNet-Santé is a web-based prospective observational cohort that was launched in France in May 2009. The objectives, design and methodology have been described elsewhere ⁽²⁵⁾. The study was conducted according to the guidelines laid down in the Declaration of Helsinki and was approved by the Institutional Review Board of the French Institute for Health and Medical Research (IRB Inserm no. 0000388FWA00005831) and the “Commission

Nationale de l'Informatique et des Libertés” (CNIL no. 908450 and no. 909216). All subjects signed an electronic informed consent.

Data collection and computation

Volunteers filled in self-administrated questionnaires using a dedicated website at baseline and during follow-up on an approximately monthly basis. The baseline questionnaires, inquired sociodemographic data and lifestyles, health status, physical activity, anthropometrics and diet. These questionnaires were first pilot-tested and then compared to traditional assessment methods or objectively validated ⁽²⁶⁻²⁸⁾.

Consumption frequency of organic products data

Two months after enrollment, volunteers were asked to provide information on their consumption frequency of 16 labeled organic products (fruit, vegetables, soya, dairy products, meat and fish, eggs, grains and legumes, bread and cereals, flour, vegetable oils and condiments, ready-to-eat meals, coffee/tea/herbal tea, wine, biscuits/chocolate/sugar/marmalade, other foods, dietary supplements). Initially, the collection of these data was related to research questions focused on reasons for non-consumptions. Consumption frequencies were presented in 8 modalities: (1) most of the time; 2) occasionally; 3) never (“too expensive”); 4) never (“product not available”); 5) never (“I’m not interested in organic products”); 6) never (“I avoid such products”); 7) never (“for no specific reason”); and 8) “I don’t know”). For each product, we allocated two and one points to the “most of the time” and “occasionally” modalities, respectively (and 0 otherwise) - since the objective of the present study was to focus on the level of frequency, and not on reasons for non-consumption. The 16 dietary components were summed up to provide an **organic score** (ranging from 0 to 32).

Anthropometric data

At enrollment and yearly after, participants were asked to report weight and height assessed during a medical or occupational health examination by a physician, or from self-measurement guided by standardized procedures (on flat surface, lightly dressed, and without shoes). Self-reported anthropometric data have been shown to present an elevated concordance with clinical assessment ⁽²⁶⁾.

Body mass index (kg/m²) was calculated as weight divided by the square of height. Subjects were classified as underweight or normal weight (BMI<25), overweight (including obesity; BMI≥25), or obese (BMI≥30) according to the World Health Organization (WHO) reference values ⁽¹⁾.

Dietary data and physical activity

At baseline, quantitative dietary intakes were assessed using three 24-hour records (24HR), randomly allocated over a two-week period, including two week-days and one weekend day, using a validated method ^(27,28). Participants reported all foods and beverages consumed at each eating occasion. Portion sizes were estimated with the help of photographs, derived from a previously validated picture booklet ⁽²⁹⁾ or directly entered as grams, volumes or purchased units. Since alcohol is only episodically consumed by most individuals, alcohol intake was calculated using either the 24HR or a frequency questionnaire for those identified as abstainers in the three 24HR days. Moreover, since fish and seafood are infrequently consumed by many individuals, the weekly consumption of this food group was assessed by a specific frequency question. Individual daily mean food consumption was calculated from the three 24HR, weighted for the type of day (week or weekend day). Nutrient intakes were calculated using the NutriNet-Santé composition table ⁽³⁰⁾. Under-reporters were identified and excluded using the validated method developed by Black ⁽³¹⁾. To assess nutritional diet quality, a modified version of the validated PNNS-GS (without physical activity) was computed. This modified score, the mPNNS-GS, reflects adherence to the official French

nutritional recommendations⁽³²⁾. The score includes 12 components: eight refer to food-serving recommendations (fruit and vegetables; starchy foods; whole grain products; dairy products; meat, eggs and fish; fish and seafood; vegetable fat; water vs soda) and four refer to moderation in consumption (added fat; salt; sweets; alcohol). Moreover, points are deducted for overconsumption of salt, added sugars, or when energy intake exceeds the estimated energy needs by more than 5%.

In order to also account for *a posteriori* dietary patterns, we performed a principal component analysis (PCA), on 31 aggregated food groups. Dietary patterns obtained by PCA are independent linear combinations of the 31 food group consumptions, maximizing the explained variance. Two dietary patterns were retained based on Cattell's Scree plots and the interpretability of the factors. **Supplementary table S1** presents all factor loading coefficients (corresponding to the correlations between the different food groups and the two dietary patterns) > 0.3 . For each participant, the individual pattern score was calculated by summing the intake of the 31 food groups, weighted by their factor loading.

Covariates

At baseline, self-administered questionnaires were used to collect data including age, gender, formal education ($\leq$ high school diploma, high school, post-secondary graduate), occupation (managerial staff, intermediate profession, employee/manual worker, retired, unemployed, never employed/homemaker and self-employed), marital status (cohabiting or single), income, number of children and smoking status (never, former and current). Income per household unit was calculated by dividing the household's total monthly income by the number of consumption units (CU), using the following coefficients: 1 CU for the first adult in the household, 0.5 CU for all other household members aged 14 or older and 0.3 CU for children under 14⁽³³⁾. The following categories of monthly income were used: $<1,200$, 1,200-1,800, 1,800-2,700 and $>2,700$ euros per household unit.

Physical activity was assessed by the International Physical Activity Questionnaire (IPAQ)

⁽³⁴⁾. Metabolic equivalents (MET) measured in minutes per week were computed. The

recommended IPAQ categories of physical activity were used: low (<30 min of brisk

walking/day), moderate (30 - <60 min/day of brisk walking /day or equivalent) and high (≥ 60

min of brisk walking /day or equivalent).

The inquired baseline health data included use of medication and self-reported history of

diseases (cancer, cardiovascular diseases, hypertension, dyslipidemia and diabetes).

Statistical analysis

For the present study, we used data from volunteers who were included before June 2014 and

initially aged 18-74 y. Among them, we selected those 1) who completed the organic

questionnaire, 2) with baseline anthropometric data and at least one measurement during

follow-up, 3) with available data to compute PNNS-GS and 4) who were not identified as

energy underreporters (**Figure 1**). Follow-up anthropometric data were collected until June

2015.

The participants included (N=62,224) into our analyses were compared to those excluded,

using Mann-Whitney-U tests and Chi square tests.

Baseline characteristics were presented by quartiles of the organic score. Values represent

means ($\pm$ SD) or percentages, and P-values were calculated using linear contrast tests (for

continuous variables) or Chi square trend tests (for categorical variables).

The association of the organic score with BMI change was assessed by covariance analysis

(ANCOVA), modeling change in BMI as a percentage of the baseline value. Mean differences

(95% confidence intervals) were presented across quartiles of the organic score. Three

different models were performed. The first model was adjusted for baseline age and gender.

The second model was further adjusted for year and month of inclusion, follow-up duration,

occupation, marital status, education, monthly income per household unit, baseline use of

185 dietary supplements, the mPNNS-GS, PCA-extracted dietary patterns scores, energy intake,
186 physical activity and smoking status. The final model was further adjusted for history of
187 diseases (cancer, cardiovascular diseases, diabetes, hypertension and dyslipidemia).
188 In a second set of analyses, we estimated odds ratios (OR) and 95% confidence intervals (CI)
189 for becoming overweight or obese after exclusion of overweight and obese subjects at
190 baseline, respectively (leading to new study samples of n= 43,301 and n=56,806). Three
191 multivariate logistic regression models were computed with similar covariables to those used
192 in the covariance analysis.

193 A set of supplementary analyses was performed for the obesity risk outcome. First, stratified
194 analyses were conducted according to physical activity level/day brisk walking versus ≥ 30
195 min/day brisk walking or equivalent), use of dietary supplements (yes versus no), tobacco
196 status (never and former smokers versus current smokers), education level, and level of
197 adherence to nutritional guidelines (using tertiles of the mPNNS-GS). Second, we used an
198 alternative method of accounting for potential confounder bias: adjustment for a ‘propensity
199 score’ that contains information on potential confounders in a combined manner⁽³⁵⁾. To
200 obtain the propensity score, a multinomial logistic regression model was used to estimate the
201 predicted probability of organic food consumption (using quartiles of the 16-point organic
202 score) as a function of a wide range of factors (sociodemographic, health characteristics, food
203 group consumptions). Finally, we used inverse probability weighting to correct the estimates
204 for potential selection bias⁽³⁶⁾. All tests of statistical significance were two-sided and the type
205 I error was set at 5%. Statistical analyses were performed using SAS® software (version 9.3,
206 SAS Institute Inc, Cary, NC, USA).

207 **Results**

208 *Comparison of included and excluded participants*

Compared with excluded subjects, included subjects (N=62,224) were older, had a higher education level and income, were less often smokers and less physically active. They also presented a lower BMI and a higher nutritional quality of the diet (all P-values<0.05, data not tabulated).

Baseline Characteristics of the sample

Baseline characteristics across quartiles of organic score are shown in **Table 1**. Higher levels of the organic score were related to higher proportions of women, participants who were cohabiting, former smokers, physically active participants, individuals reaching a post-secondary educational level, and participants with a high income or occupational level. A positive association was also observed with age, follow-up duration and the mPNNS-GS (reflecting the nutritional quality of the diet), while there was a negative association with energy intake, alcohol consumption and BMI.

Components of the organic score across quartiles of the organic score are shown in **Table 2**.

In the first quartile (Q1), participants mostly reported no consumption of any organic products. In the 4th quartile (Q4), participants reported more frequently consuming organic products, especially products of the following food groups: eggs, starchy food, vegetables, vegetable oil, fruits and flour.

Then mean follow-up time in our study sample was 3.12 (SD=1.37) years. Results on the prospective association between the organic score and change in BMI over time are presented in **Table 3**. In the second model, higher organic score levels were related to a substantially lower increase in BMI over time (mean difference Q4 versus Q1= -0.15; confidence interval= -0.31, -0.01, P for trend =0.05). After further adjustment for history of chronic diseases (third model), an even stronger association was observed (mean difference Q4 versus Q1= -0.16; confidence interval= -0.32, -0.01, P for trend =0.04).

Results on the prospective association between the organic score and the risk of overweight and obesity are presented in **Table 4**. In the fully-adjusted model, accounting for sociodemographic data, lifestyles and history of chronic diseases, a linear decrease in the risk of overweight was observed across quartiles of the organic score, with risk reduction of 23% in Q4 compared to Q1. Findings concerning the risk of obesity were similar, with a risk reduction of 31% in Q4 compared to Q1.

Stratified analyses are presented in **Figure 2** and in **Supplemental Table S2**. We observed that the association between the organic score and the risk of obesity was stronger among participants with a higher nutritional quality of the diet. Overall, in the different stratified analyses, the association between the organic score and the risk of obesity was observed in each sub-group, except for dietary supplement users and participants with an intermediate education level. Accounting for selection bias via inverse probability weighting did not substantially modify our findings (data not shown). Models with additional adjustment for a propensity score are presented in **Supplemental Table S3**. Here, the investigated associations were attenuated but remained statistically significant.

Discussion

The present results show, for the first time, a strong negative association between the consumption frequency of organic foods and BMI change over time, as well as a marked reduction of the risk of overweight and obesity.

In stratified analyses, significant associations were observed in almost all investigated subgroups, except for dietary supplement users and participants with an intermediate level of education. Importantly, both participants with a low level of physical activity and participants with a low education level presented a significantly lower risk of obesity with increasing organic food consumption. It is noteworthy that the nutritional quality of the diet, estimated by using an a priori dietary index reflecting adherence to the French nutritional guidelines⁽³²⁾,

appears to be a key effect modifier. Indeed, the strongest associations were observed among participants presenting a high nutritional quality of the diet.

Consumption frequency of organic foods and the risk of obesity or overweight

We prospectively observed a markedly lower risk of obesity among subjects with a high consumption frequency of organic foods. Previous investigations of data from the NutriNet-Santé cohort have revealed that regular consumers of organic foods presented a healthier diet and healthier lifestyle characteristics (related to physical activity and tobacco use) as compared to irregular consumers or non-consumers. Moreover, regular consumers of organic foods in the NutriNet-Santé study presented specificities with respect to chronic disease history^(23,37). After accounting for these potential confounders, the findings of the present study were partially attenuated but the association remained strong and highly significant, with a reduction in the risk of obesity of 37% after a 3·1 y follow-up.

A similar association was observed for overweight, although the strength of the association was smaller.

Comparison of our results with the findings of other studies

No previous longitudinal study has investigated the association between organic food consumption frequency and weight change or the risk of overweight and obesity, but a potential beneficial link between body weight and organic food consumption or purchase has been documented in several cross-sectional studies^(23,24,38). For instance, a previous investigation of NutriNet-Santé data⁽²³⁾ has shown that regular organic food consumers showed a markedly lower probability of overweight (excluding obesity) and obesity compared to non-consumers (-36% and -62% in men and -42% and -48% in women, respectively). In addition, in the German National Nutrition Survey II⁽²⁴⁾, as compared to non-buyers of organic food, buyers of organic food presented lower proportions of overweight (35·5% versus 39·2%) and obesity (17·9% versus 22·5%). Our findings are also

concordant with a small Italian clinical study (including 100 healthy males and 50 males suffering from chronic kidney disease, CDK) that reported a statistically significant reduction in weight among CDK patients after introducing an organic diet for a 2-week period (85.17 ± 13.97 kg at baseline versus 79.52 ± 10.41 kg after the 2-week intervention, $p < 0.05$)⁽³⁹⁾. Overall, the currently available cross-sectional or longitudinal surveys consistently reported an association between higher organic food consumption and a lower BMI. The association with a lower increase in BMI over time that we observed in the present study is of particular interest as it supports a possible role of the organic-based diet in weight management among all subjects, beyond the risk of overweight or obesity among initially normal weight (or underweight) individuals.

Modulating effect of the nutritional quality of the diet

There is growing evidence supporting the observation that consumers of organic food present a nutritionally healthier diet as well as other beneficial lifestyles such as lower alcohol consumption, no smoking and a higher physical activity level^(23,24,40,41). Thus, we hypothesized that the link between consumption of organic foods and body weight might be modulated by the overall nutritional quality of the diet. Indeed, we found that the strongest reduction in obesity risk (related to a higher consumption frequency of organic foods) was observed among participants with a healthier diet. Conversely, among participants with a less healthy diet (susceptible to promote weight gain), the association between consumption frequency of organic food and obesity risk was of smaller magnitude. It is well known that unhealthy eating habits play a key role in the etiology of obesity⁽³⁾. Therefore, the mode of production of the foods consumed may be of secondary importance in these subjects. It is also possible that their dietary patterns, which are low in fruit and vegetables, lead to less contaminated diets.

Several hypotheses can be proposed to explain the differences in the risk of obesity between organic food consumers and non-consumers who present a healthy diet. Because we adjusted and then stratified for the global nutritional quality of the diet (using the mPNNS-GS score) in our models, it is not probable that our results were biased by the fact that organic food consumers present healthier diets. Other possible explanations include differences in organic food and conventional food with respect to various nutritional compounds.

First, the results of a number of studies argue for a higher concentration of polyunsaturated fatty acids (especially n-3 fatty acids) in organic dairy and meat products; and of antioxidants (especially vitamin C and phenolic compounds) in organic plant foods⁽¹¹⁻¹³⁾. While observed differences in nutrient content can vary by about 10-68% at the food level (not accounted for in our study), it is possible that the overall variations of nutrient intake in an organic diet is sufficient to affect weight management^(42,43). However, this remains to be further evaluated.

Another hypothesis is related to the fact that individuals with a higher adherence to the French nutritional guidelines tend to consume more plant-based foods. It is well known that plant foods are frequently contaminated by various pesticide residues (about 45% of the tested samples in Europe)⁽⁴⁴⁾ because they are heavily sprayed with pesticides during conventional agricultural production and storage. This hypothesis is in line with findings of human surveys that have related obesity and type 2 diabetes to pesticide exposure^(7,9,10,45,46). Thus, unlike the consumption of pesticide-free or only slightly contaminated plant products^(11,14,15), a high consumption of conventionally grown plant foods may be related to adverse health effects related to higher pesticide exposure.

Indeed, replacing conventional food by organic food has repeatedly been shown to drastically reduce the level of organophosphate residues in human urines⁽¹⁶⁻²⁰⁾. A specific example of a potential adverse health effect of contaminated fruit and vegetables is that high consumers of conventional or contaminated fruits and vegetables presented a particularly low semen quality

^(47,48). This reinforces the concept that different dietary profiles (with various degrees of intake of contaminated food) lead to different levels of exposure to “obesogen” chemicals ⁽⁹⁾, but this hypothesis needs to be investigated in future biomonitoring-based studies that compare organic and conventional diets with various dietary profiles.

Potential mechanistic pathways explaining the association between organic food consumption and body weight

Our findings may be interpreted in light of the reduced exposure to pesticides among organic food consumers. Pesticides (prohibited in organic farming when they are synthetic) often present endocrine-disrupting properties that cause developmental and reproductive abnormalities via the modification of signaling processes ⁽⁴⁶⁾. Besides, new scientific research argues for a role of pesticides in metabolic disruption, ⁽²²⁾ leading to obesity and type 2 diabetes ⁽⁹⁾. Mechanistic pathways depend on the type of pesticides. Previous studies have shown an alteration of glucose and lipid metabolism by organochlorines ⁽⁴⁹⁾. Moreover, organochlorines have been shown to affect the control of adipogenesis through an alteration of glucose transport and glycolysis, an alteration of mitochondrial activity, and of fatty acid oxidation ⁽⁴⁹⁾. Organophosphates have been shown to alter carbohydrate and lipid metabolism by disrupting glucose homeostasis ⁽⁴⁹⁾. In addition, some pesticides have been shown to affect the regulation of eating behavior and the differentiation of adipocytes ⁽⁵⁰⁾.

Further studies are necessary to better evaluate metabolic disruption and the “obesogen” capacity of endocrine-disrupting chemicals contained in conventional foods.

Strengths and limitations

The main limitation of our study is that it is based on self-reported weight and height data. However, these data have been shown to have a good concordance with data from clinical assessments in a validation study ⁽²⁶⁾. This validation study showed high intraclass correlation coefficients, ranging from 0.94 for height to 0.99 for weight, and the concordance for BMI

classification was 93% (sensitivity=88% and specificity=99%). Second, the generalizability of our findings is limited since participants were volunteers involved in a long-term cohort focused on nutrition and health. The individuals included in our study sample are thus likely to be particularly health conscious. A final limitation pertains to the difficulty to disentangle the role of overall dietary patterns from the role of organic food consumption, despite of the extensive adjustment and stratification made. Since the design of our study is observational, residual confounding cannot be ruled out. In particular, it is likely that unmeasured or only indirectly measured factors, including genetic factors, ethnicity, environmental (e.g. food or built environment) or psychological factors (e.g. occupational stress) may modify the association between organic food consumption and obesity.

Our study also presents important strengths. First, the rich and accurately collected data permitted to account for a broad range of potential confounders, including lifestyles and health outcomes. Moreover, the very large sample size of our study enabled us to conduct statistically powerful stratified analyses. Another important strength is the prospective design of our analysis that implies a high level of evidence. Finally, the availability of accurate dietary data allowed us to adjust for the nutritional quality of the diet, using a validated dietary index.

Conclusion

This study, based on data collected in a very large prospective cohort, is the first to support a prospective relation between consumption frequency of organic foods and body weight change, as well as a strong negative association with the risk of overweight and obesity. The overall nutritional quality of the diet may exert a modulating effect in these relationships, with a stronger effect observed among those presenting a healthy plant-based diet. Further studies, especially studies based on quantitative organic consumption data taking into account a diversity of dietary profiles (plant based, western etc.), are needed to confirm these results. If

confirmed, these findings are of major interest from a public health point of view, as they reinforce the need to fine-tune nutritional guidelines by accounting for the mode of food production.

Acknowledgment

The authors Nathalie Arnault, Stephen Besseau, Laurent Bourhis, Julien Allègre, Than Duong Van, Younes Esseddik, Cédric Agaesse, Claudia Chahine, Paul Flanzy, Mac Rakotondrazafy and Fabien Szabo for their technical contribution to the NutriNet-Santé study.

Financial support

The NutriNet-Santé study is supported by the French Ministry of Health (DGS), the French Institute for Health Surveillance (InVS), the National Institute for Prevention and Health Education (INPES), the Foundation for Medical Research (FRM), the National Institute for Health and Medical Research (INSERM), the National Institute for Agricultural Research (INRA), the National Conservatory of Arts and Crafts (CNAM), and the University of Paris 13.

Conflict of Interest

None of the authors declares any conflict of interest.

Authorship

EKG: conducted the literature review and drafted the manuscript; EKG: performed analyses; EKG, JB, KEA, PG, SH and DL: were involved in the interpretation of results and critically reviewed the manuscript; and SH, PG and EKG: were responsible for the development of the design and the protocol of the study. All authors read and approved the final manuscript.

Reference List

1. WHO (2000) *Obesity: preventing and managing the global epidemic.*, vol. 894: World Health Organization Technical Report Series.
2. WHO (2009) *Global Health Risks*. Geneva: WHO Technical Report.
3. WHO Europe (2007) *The challenge of obesity in the WHO European Region*.
4. Latino-Martel P, Cottet V, Druet-Pecollo N *et al.* (2016) Alcoholic beverages, obesity, physical activity and other nutritional factors, and cancer risk: A review of the evidence. *Crit Rev Oncol Hematol* **99**, 308-323.
5. American Institute for Cancer Research / World Cancer Research Fund (2007) *Food, Nutrition, Physical activity, and the Prevention of Cancer: a global perspective*. Washington, DC: American Institute for Cancer Research.
6. Hansen TH, Gobel RJ, Hansen T *et al.* (2015) The gut microbiome in cardio-metabolic health. *Genome Med* **7**, 33.
7. Lee DH, Porta M, Jacobs DR, Jr. *et al.* (2014) Chlorinated persistent organic pollutants, obesity, and type 2 diabetes. *Endocr Rev* **35**, 557-601.
8. Snedeker SM & Hay AG (2012) Do interactions between gut ecology and environmental chemicals contribute to obesity and diabetes? *Environ Health Perspect* **120**, 332-339.
9. Thayer KA, Heindel JJ, Bucher JR *et al.* (2012) Role of environmental chemicals in diabetes and obesity: a National Toxicology Program workshop review. *Environ Health Perspect* **120**, 779-789.
10. Lubrano C, Genovesi G, Specchia P *et al.* (2013) Obesity and metabolic comorbidities: environmental diseases? *Oxid Med Cell Longev* **2013**, 640673.
11. Baranski M, Srednicka-Tober D, Volakakis N *et al.* (2014) Higher antioxidant and lower cadmium concentrations and lower incidence of pesticide residues in organically grown crops: a systematic literature review and meta-analyses. *Br J Nutr* **112**, 794-811.
12. Srednicka-Tober D, Baranski M, Seal C *et al.* (2016) Composition differences between organic and conventional meat: a systematic literature review and meta-analysis. *Br J Nutr*, 1-18.
13. Srednicka-Tober D, Baranski M, Seal CJ *et al.* (2016) Higher PUFA and n-3 PUFA, conjugated linoleic acid, alpha-tocopherol and iron, but lower iodine and selenium concentrations in organic milk: a systematic literature review and meta- and redundancy analyses. *Br J Nutr*, 1-18.
14. Smith-Spangler C, Brandeau ML, Hunter GE *et al.* (2012) Are organic foods safer or healthier than conventional alternatives?: a systematic review. *Ann Intern Med* **157**, 348-366.
15. Lairon D (2010) Nutritional quality and safety of organic food: a review. *Agron Sustain Dev* **30**, 33-41.

- 440 16. Curl CL, Fenske RA & Elgethun K (2003) Organophosphorus pesticide exposure of urban and
441 suburban preschool children with organic and conventional diets. *Environ Health*
442 *Perspect* **111**, 377-382.
- 443 17. Lu C, Toepel K, Irish R *et al.* (2006) Organic diets significantly lower children's dietary
444 exposure to organophosphorus pesticides. *Environ Health Perspect* **114**, 260-263.
- 445 18. Bradman A, Quiros-Alcala L, Castorina R *et al.* (2015) Effect of Organic Diet Intervention on
446 Pesticide Exposures in Young Children Living in Low-Income Urban and Agricultural
447 Communities. *Environ Health Perspect* **123**, 1086-1093.
- 448 19. Oates L, Cohen M, Braun L *et al.* (2014) Reduction in urinary organophosphate pesticide
449 metabolites in adults after a week-long organic diet. *Environ Res* **132**, 105-111.
- 450 20. Curl CL, Beresford SA, Fenske RA *et al.* (2015) Estimating pesticide exposure from dietary
451 intake and organic food choices: the Multi-Ethnic Study of Atherosclerosis (MESA).
452 *Environ Health Perspect* **123**, 475-483.
- 453 21. Mnif W, Hassine AI, Bouaziz A *et al.* (2011) Effect of endocrine disruptor pesticides: a review.
454 *Int J Environ Res Public Health* **8**, 2265-2303.
- 455 22. Casals-Casas C & Desvergne B (2011) Endocrine disruptors: from endocrine to metabolic
456 disruption. *Annu Rev Physiol* **73**, 135-162.
- 457 23. Kesse-Guyot E, Peneau S, Mejean C *et al.* (2013) Profiles of organic food consumers in a large
458 sample of French adults: results from the Nutrinet-Sante cohort study. *PLoS One* **8**,
459 e76998.
- 460 24. Eisinger-Watzl M, Wittig F, Heuer T *et al.* (2015) Customers Purchasing Organic Food - Do
461 They Live Healthier? Results of the German National Nutrition Survey II. *Eur J Nutr*
462 *Food Saf* **5**, 59-71.
- 463 25. Hercberg S, Castetbon K, Czernichow S *et al.* (2010) The Nutrinet-Sante Study: a web-based
464 prospective study on the relationship between nutrition and health and determinants of
465 dietary patterns and nutritional status. *BMC Public Health* **10**, 242.
- 466 26. Lassale C, Peneau S, Touvier M *et al.* (2013) Validity of web-based self-reported weight and
467 height: results of the Nutrinet-Sante study. *J Med Internet Res* **15**, e152.
- 468 27. Lassale C, Castetbon K, Laporte F *et al.* (2015) Validation of a Web-based, self-administered,
469 non-consecutive-day dietary record tool against urinary biomarkers. *Br J Nutr* **113**, 953-
470 962.
- 471 28. Lassale C, Castetbon K, Laporte F *et al.* (2015) Correlations between Fruit, Vegetables, Fish,
472 Vitamins, and Fatty Acids Estimated by Web-Based Nonconsecutive Dietary Records
473 and Respective Biomarkers of Nutritional Status. *J Acad Nutr Diet*.
- 474 29. Le Moullec N, Deheeger M, Preziosi P *et al.* (1996) Validation du manuel photos utilisé pour
475 l'enquête alimentaire de l'étude SU.VI.MAX. *Cahier de Nutrition et de Diététique* **31**,
476 158-164.
- 477 30. NutriNet-Santé coordination (2013) *Table de composition des aliments - Etude NutriNet-Santé*.
478 Paris: Economica.

- 479 31. Black AE (2000) Critical evaluation of energy intake using the Goldberg cut-off for energy
480 intake:basal metabolic rate. A practical guide to its calculation, use and limitations. *Int J*
481 *Obes Relat Metab Disord* **24**, 1119-1130.
- 482 32. Estaquio C, Kesse-Guyot E, Deschamps V *et al.* (2009) Adherence to the French Programme
483 National Nutrition Sante Guideline Score is associated with better nutrient intake and
484 nutritional status. *J Am Diet Assoc* **109**, 1031-1041.
- 485 33. INSEE (Institut National de la Statistique et des Etudes Economiques) website (2012).
486 <http://www.insee.fr/en/methodes/>.
- 487 34. Hagstromer M, Oja P & Sjostrom M (2006) The International Physical Activity Questionnaire
488 (IPAQ): a study of concurrent and construct validity. *Public Health Nutr* **9**, 755-762.
- 489 35. Ali MS, Groenwold RH & Klungel OH (2016) Best (but oft-forgotten) practices: propensity
490 score methods in clinical nutrition research. *Am J Clin Nutr* **104**, 247-258.
- 491 36. Seaman SR & White IR (2011) Review of inverse probability weighting for dealing with
492 missing data. *Stat Methods Med Res*.
- 493 37. Baudry J, Mejean C, Peneau S *et al.* (2015) Health and dietary traits of organic food consumers:
494 results from the NutriNet-Sante study. *Br J Nutr*, 1-10.
- 495 38. Torjusen H, Brantsaeter AL, Haugen M *et al.* (2010) Characteristics associated with organic
496 food consumption during pregnancy; data from a large cohort of pregnant women in
497 Norway. *BMC Public Health* **10**, 775.
- 498 39. De Lorenzo A, Noce A, Bigioni M *et al.* (2010) The effects of Italian Mediterranean organic
499 diet (IMOD) on health status. *Curr Pharm Des* **16**, 814-824.
- 500 40. Oates L, Cohen M & Braun L (2012) Characteristics and consumption patterns of Australian
501 organic consumers. *J Sci Food Agric* **92**, 2782-2787.
- 502 41. Petersen SB, Rasmussen MA, Strom M *et al.* (2013) Sociodemographic characteristics and food
503 habits of organic consumers--a study from the Danish National Birth Cohort. *Public*
504 *Health Nutr* **16**, 1810-1819.
- 505 42. Lorente-Cebrian S, Costa AG, Navas-Carretero S *et al.* (2013) Role of omega-3 fatty acids in
506 obesity, metabolic syndrome, and cardiovascular diseases: a review of the evidence. *J*
507 *Physiol Biochem* **69**, 633-651.
- 508 43. Meydani M & Hasan ST (2010) Dietary polyphenols and obesity. *Nutrients* **2**, 737-751.
- 509 44. EFSA (European Food Safety Authority) (2015) The 2013 European Union Report on Pesticide
510 Residues in Food. *EFSA J* **13**.
- 511 45. Lee DH, Steffes MW, Sjodin A *et al.* (2011) Low dose organochlorine pesticides and
512 polychlorinated biphenyls predict obesity, dyslipidemia, and insulin resistance among
513 people free of diabetes. *PLoS One* **6**, e15977.
- 514 46. Heindel JJ, Newbold R & Schug TT (2015) Endocrine disruptors and obesity. *Nat Rev*
515 *Endocrinol* **11**, 653-661.
- 516 47. Chiu YH, Afeiche MC, Gaskins AJ *et al.* (2015) Fruit and vegetable intake and their pesticide
517 residues in relation to semen quality among men from a fertility clinic. *Hum Reprod* **30**,
518 1342-1351.

- 519 48. Juhler RK, Larsen SB, Meyer O *et al.* (1999) Human semen quality in relation to dietary
520 pesticide exposure and organic diet. *Arch Environ Contam Toxicol* **37**, 415-423.
- 521 49. Androutsopoulos VP, Hernandez AF, Liesivuori J *et al.* (2013) A mechanistic overview of
522 health associated effects of low levels of organochlorine and organophosphorous
523 pesticides. *Toxicology* **307**, 89-94.
- 524 50. Mostafalou S & Abdollahi M (2013) Pesticides and human chronic diseases: evidences,
525 mechanisms, and perspectives. *Toxicol Appl Pharmacol* **268**, 157-177.
526
527

533 **Table 2 Components of the organic score, across organic score quartiles, Nutrinet-Santé**
 534 **study, N=62,224***

	Q1	Q2	Q3	Q4	P [†]
Fruits	0·06 (0·25)	0·57 (0·52)	0·95 (0·41)	1·49 (0·53)	<0·0001
Vegetables	0·07 (0·27)	0·59 (0·55)	0·97 (0·48)	1·51 (0·54)	<0·0001
Rice, pasta, other cereals	0·06 (0·23)	0·42 (0·51)	0·90 (0·50)	1·62 (0·52)	<0·0001
Breads	0·05 (0·22)	0·36 (0·51)	0·82 (0·53)	1·44 (0·59)	<0·0001
Flour	0·04 (0·20)	0·23 (0·45)	0·67 (0·62)	1·47 (0·65)	<0·0001
Vegetable oils	0·04 (0·22)	0·27 (0·51)	0·74 (0·65)	1·58 (0·58)	<0·0001
Dairy products	0·05 (0·23)	0·40 (0·54)	0·86 (0·58)	1·42 (0·65)	<0·0001
Meats & seafoods	0·02 (0·14)	0·19 (0·41)	0·50 (0·54)	0·92 (0·66)	<0·0001
Eggs	0·13 (0·36)	0·70 (0·70)	1·19 (0·70)	1·72 (0·54)	<0·0001
Soja products	0·06 (0·24)	0·20 (0·44)	0·40 (0·59)	0·88 (0·76)	<0·0001
Sweet products	0·03 (0·19)	0·33 (0·48)	0·76 (0·50)	1·38 (0·58)	<0·0001
Tea, coffee, herbal tea	0·04 (0·19)	0·27 (0·47)	0·66 (0·59)	1·33 (0·67)	<0·0001
Wines	0·03 (0·17)	0·14 (0·35)	0·29 (0·47)	0·65 (0·64)	<0·0001
Ready-to-use dishes	0·01 (0·08)	0·06 (0·24)	0·19 (0·39)	0·39 (0·54)	<0·0001
Dietary supplements	0·03 (0·18)	0·13 (0·36)	0·25 (0·49)	0·56 (0·68)	<0·0001
Other dietary items	0·01 (0·12)	0·09 (0·30)	0·24 (0·45)	0·80 (0·80)	<0·0001

535 Q ; Quartile

536 *Values are means (SD) of sub-score given that non-consumption, occasional, regular
 537 consumption are coded 0, 1, 2, respectively.

538 [†]P for linear trend

Table 3 Association between organic score in quartile and BMI change over time, Nutrinet-Santé study, 2009-2015, N=62,224*

model	Q1	Q2	Q3	Q4	P for trend
Model 1 [†]	0·00 (ref)	0·01 (-0·13, 0·15)	-0·06 (-0·20, 0·08)	-0·34 (-0·49, -0·20)	<0·0001
Model 2 [‡]	0·00 (ref)	0·03 (-0·11, 0·17)	0·01 (-0·13, 0·16)	-0·15 (-0·31, -0·00)	0·05
Model 3 [§]	0·00 (ref)	0·03 (-0·12, 0·17)	0·01 (-0·14, 0·15)	-0·16 (-0·32, -0·01)	0·04

Q; Quartile

*Values are mean differences (95% confident intervals). A negative value (-0·xx) indicates that the observed increase (expressed as a percentage of the initial anthropometric marker) was lower of 0·xx in the respective quartile than in quartile 1 (ref).

[†]Model 1 is adjusted for age and gender.

[‡]Model 2 is model 1 further adjusted for month and year of inclusion, duration of follow-up, occupation, marital status, education, monthly income per unit, dietary supplement use, mPNNS-GS, PCA-extracted dietary patterns scores, energy intake, physical activity and tobacco status.

[§]Model 3 is model 2 further adjusted for the history of chronic diseases.

Table 4 Prospective association between quartiles of the organic score and overweight or obesity risk, Nutrinet-Santé study, 2009-2015^{*}.

Model	Q1	Q2	Q3	Q4	P for trend
Overweight (N=3,259/40,042)					
Model 1 [†]	1·00	0·94 (0·86- 1·04)	0·83 (0·75- 0·91)	0·62 (0·56- 0·69)	<·0001
Model 2 [‡]	1·00	1·00 (0·91, 1·10)	0·93 (0·84, 1·02)	0·75 (0·67, 0·84)	<·0001
Model 3 [§]	1·00	1·00 (0·91, 1·10)	0·93 (0·84, 1·03)	0·77 (0·68, 0·86)	<·0001
Obesity (1,337/55,469)					
Model 1 [†]	1·00	0·87 (0·75, 1·01)	0·79 (0·68, 0·91)	0·52 (0·45, 0·61)	<·0001
Model 2 [‡]	1·00	0·93 (0·80, 1·08)	0·90 (0·78, 1·04)	0·66 (0·55, 0·78)	<·0001
Model 3 [§]	1·00	0·94 (0·81, 1·09)	0·92 (0·79, 1·06)	0·69 (0·58, 0·82)	0·0001

^{*}Overweight (including obesity) and obesity analyses were performed among participants who were not overweight or obese at inclusion, respectively. Values are odds ratios (95% confident intervals), using 1st quartile Q1 as the reference.

[†]Model 1 is adjusted for age and gender.

[‡]Model 2 is model 1 further adjusted for month and year of inclusion, delay of follow-up, occupation, marital status, education, monthly income per unit, dietary supplement use, mPNNS-GS, PCA-extracted dietary patterns scores, energy intake physical activity and tobacco status.

[§]Model 3 is model 2 further adjusted for the history of chronic diseases.

562 **Supplemental Table S1: Factor loading matrix, NutriNet-Santé study***

Variable	Factor1	Factor2
Vegetables	0·65	0·14
Fruit	0·45	0·06
Broth	0·43	0·17
Coffee, tea, tisane	0·43	0·01
Soda	-0·42	0·06
Croissants	-0·37	0·03
Nuts	0·36	0·04
Fat and sweet products	-0·35	0·07
Dried fruits	0·33	-0·03
Snacks	-0·28	0·23
Soya products	0·28	-0·16
Meat and meat products	-0·28	0·39
Seafood	0·27	-0·02
Dairy desserts	-0·25	0·04
Bread and breakfast cereals	0·22	0·54
Cakes and pastries	-0·18	0·15
Alcoholic beverages	-0·18	0·34
Margarine	0·16	0·12
Flour	0·13	0·06
Eggs	0·12	0·02
Animal Fat	0·09	0·41
Dairy products	0·07	-0·09
Cheese	-0·05	0·53
Potatoes	0·05	0·41
Vegetable oils	0·03	0·28
Starchy food	0·03	0·07
Meal substitutes	0·02	-0·16
Sauces	-0·01	0·23
Sweet products	-0·01	0·27
Water and non-sweet bevarages	0·01	0·14
Wine	0·01	0·42

563 *Absolute values < 0·3 are not displayed in the table.

564 **Supplemental Table S2 Prospective association between quartiles of the organic score**
 565 **and obesity risk, stratified analyses, Nutrinet-Santé study, 2009-2015^{*,†}.**

	Q1	Q2	Q3	Q4	P for trend
Education					
< high school diploma	1·00	1·21 (0·91, 1·61)	0·88 (0·65, 1·19)	0·73 (0·52, 1·04)	0·04
High school diploma	1·00	0·80 (0·56, 1·14)	0·97 (0·69, 1·35)	0·82 (0·55, 1·20)	0·48
Post-secondary graduate	1·00	0·95 (0·78, 1·15)	0·92 (0·75, 1·12)	0·69 (0·55, 0·87)	0·003
Physical activity					
<30 min/day brisk walking	1·00	0·68 (0·51, 0·90)	0·76 (0·58, 1·01)	0·53 (0·38, 0·74)	0·001
≥30 min/day brisk walking or equivalent	1·00	0·97 (0·80, 1·18)	0·90 (0·73, 1·10)	0·74 (0·59, 0·92)	0·01
Smoking status					
Never or former smokers	1·00	0·95 (0·81, 1·11)	0·90 (0·76, 1·06)	0·72 (0·59, 0·86)	0·001
Current smokers	1·00	0·74 (0·51, 1·06)	0·95 (0·65, 1·38)	0·55 (0·36, 0·86)	0·04
Dietary supplement users					
yes	1·00	1·16 (0·94, 1·45)	1·05 (0·84, 1·32)	0·79 (0·61, 1·03)	0·11
no	1·00	0·92 (0·75, 1·12)	0·75 (0·60, 0·95)	0·67 (0·52, 0·85)	0·0003

566 ^{*}Obesity analyses were performed among participants who were not obese at inclusion
 567 (N=56,806). Values are odds ratios (95% confident intervals), using 1st quartile Q1 as the
 568 reference.

569 [†]Models are adjusted for age, gender, month and year of inclusion, delay of follow-up,
 570 occupation, marital status, education, monthly income per unit, dietary supplement use,
 571 mPNNs-GS, PCA-extracted dietary patterns scores, energy intake physical activity, tobacco
 572 status and for the history of chronic diseases.

573 **Supplemental Table S3 Prospective association between quartiles of the organic score**
 574 **and obesity risk, sensitive analysis, Nutrinet-Santé study, 2009-2015^{*}.**

Model	Q1	Q2	Q3	Q4	P for trend
Model 1 [†]	1·00	0·87 (0·81, 0·94)	0·79 (0·73, 0·85)	0·52 (0·48, 0·57)	<·0001
Model 2 [‡]	1·00	0·93 (0·87, 1·01)	0·91 (0·84, 0·98)	0·67 (0·61, 0·72)	<·0001
Model 3 [§]	1·00	0·95 (0·88, 1·02)	0·92 (0·86, 0·99)	0·70 (0·64, 0·76)	<·0001


575 ^{*}Obesity analyses were performed among participants who were not obese at inclusion
 576 (N=56,806). Values are odds ratios (95% confident intervals), using 1st quartile Q1 as the
 577 reference.

578 [†]Model 1 is adjusted for age and gender and propensity score.

579 [‡]Model 2 is model 1 further adjusted for month and year of inclusion, delay of follow-up,
 580 occupation, marital status, education, monthly income per unit, dietary supplement use,
 581 mPNNS-GS, PCA-extracted dietary patterns scores, energy intake physical activity and
 582 tobacco status.

583 [§]Model 3 is model 2 further adjusted for the history of chronic diseases.

584 **Figure 1. Participants of the NutriNet-Santé selected for the present analyses, 2009–2015**



585

586

587 **Figure 2. Prospective association between organic score in quartile and risk of obesity**
 588 **stratified according the nutritional quality of the diet, Nutrinet-Santé, 2009-2015***




Figure 2. Prospective association between organic score in quartile and risk of obesity stratified according the nutritional quality of the diet, Nutrinet-Santé, 2009-2015*

*Values are odds ratios (95% confident intervals) using the 1st quartile (Q1) as the reference, adjusted for age, gender, month and year of inclusion, delay of follow-up, occupation, marital status, education, monthly income per unit, dietary supplement use, mPNNS-GS, PCA-extracted dietary patterns scores energy intake, physical activity, tobacco status and history of chronic diseases.

589

590 *Values are odds ratios (95% confident intervals) using the 1st quartile (Q1) as the reference, adjusted for age,
 591 gender, month and year of inclusion, delay of follow-up, occupation, marital status, education, monthly income
 592 per unit, dietary supplement use, mPNNS-GS, PCA-extracted dietary patterns scores energy intake, physical
 593 activity, tobacco status and history of chronic diseases.

528 **Table 1 Baseline characteristics of the sample across organic score quartiles, Nutrinet-Santé study, 2009-2014, N=62,224***

	Q1	Q2	Q3	Q4	P [†]
N	15,245	16,249	15,807	14,923	
Organic score max=32	0·73 (0·82)	4·96 (1·41)	10·36 (1·68)	19·15 (4·21)	<0·0001
Female (%)	73·45	77·94	78·71	81·63	<0·0001
Age (years)	44·16 (15·34)	44·36 (14·83)	45·62 (14·29)	46·72 (13·29)	<0·0001
Follow-up duration (days)	1140·34 (504·64)	1157·01 (500·06)	1144·88 (501·80)	1118·12 (499·69)	<0·0001
Education (%)					<0·0001
Unidentified	0·68	0·66	0·58	0·82	
< high school diploma	23·68	20·09	17·21	15·22	
High school diploma	19·12	17·01	15·44	13·99	
Post-secondary graduate	56·52	62·24	66·76	69·97	
Monthly income per unit household unit in € (%)					<0·0001
Missing	12·06	11·50	10·62	9·35	
900-1200	19·48	16·13	13·26	12·36	
1200-1800	26·76	25·15	22·67	23·09	
1800-2700	21·66	23·72	24·89	26·20	
>2700	20·05	23·50	28·56	29·00	

Occupational categories (%)					<0.0001
Non employed	5.60	5.35	5.18	5.84	
Retired	21.66	20.35	21.40	20.31	
Employee/Manual worker	23.22	19.82	16.67	14.29	
Intermediate profession	16.20	17.44	17.42	18.53	
Managerial staff	17.82	21.47	25.48	28.26	
Never employed	13.93	13.92	12.17	10.43	
Craftsman, shopkeeper, business owner, farmer	1.58	1.66	1.68	2.34	
Dietary supplement use (%)	35.52	45.55	51.58	62.13	<0.0001
Cohabiting (%)	80.89	81.88	83.22	86.32	<0.0001
Tobacco status (%)					<0.0001
Former smoker	33.05	33.09	35.51	37.69	
Current smoker	15.29	15.31	14.46	13.12	
Never smoker	51.66	51.60	50.03	49.19	
Physical activity (%)					<0.0001
Missing	15.12	14.36	12.85	11.96	
Low	15.12	14.36	12.85	11.96	
Medium	27.65	27.92	30.04	31.9	

High	33·67	37·15	38·33	39·5	
Energy intake (kcal/d)	1942·90 (514·75)	1914·76 (487·92)	1908·50 (491·42)	1894·24 (474·50)	<0·0001
% Carbohydrates	42·59 (6·89)	42·93 (6·81)	43·05 (6·94)	43·43 (7·11)	<0·0001
% Lipids	39·22 (6·61)	38·95 (6·46)	38·96 (6·58)	39·45 (6·72)	0·0017
% Proteins	17·87 (3·95)	17·80 (3·92)	17·67 (4·01)	16·81 (3·92)	<0·0001
Alcohol consumption (g/d)	8·65 (14·11)	8·27 (13·16)	8·22 (12·23)	7·63 (11·20)	<0·0001
mPNNS-GS	7·60 (1·62)	7·87 (1·62)	8·15 (1·60)	8·44 (1·58)	<0·0001
Body mass index (kg/m ²) at baseline	24·50 (4·87)	23·96 (4·56)	23·71 (4·31)	23·00 (3·90)	<0·0001
Obesity [‡] (%) at baseline	11·79	9·42	7·96	5·58	<0·0001
Body mass index (kg/m ²) at follow-up	24·72 (4·92)	24·17 (4·62)	23·91 (4·40)	23·13 (4·00)	<0·0001
Obesity [‡] (%) at follow-up	12·62	10·04	8·78	6·06	<0·0001

529 Abbreviations: Q; Quartile

530 *All variables were assessed at baseline, except when listed as “at follow-up”

531 [†]P for linear contrast

532 [‡]Body mass index ≥ 30 kg/m²