



HAL
open science

Non-coding RNAs Potentially Controlling Cell Cycle in the Model *Caulobacter crescentus*: A Bioinformatic Approach

Wanassa Beroual, M. Brili, Emanuele G Biondi

► **To cite this version:**

Wanassa Beroual, M. Brili, Emanuele G Biondi. Non-coding RNAs Potentially Controlling Cell Cycle in the Model *Caulobacter crescentus*: A Bioinformatic Approach. *Frontiers in Genetics*, 2018, 9, 10.3389/fgene.2018.00164 . hal-01916083

HAL Id: hal-01916083

<https://amu.hal.science/hal-01916083>

Submitted on 8 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Non-coding RNAs Potentially Controlling Cell Cycle in the Model *Caulobacter crescentus*: A Bioinformatic Approach

Wanassa Beroual, Emanuele G Biondi

► **To cite this version:**

Wanassa Beroual, Emanuele G Biondi. Non-coding RNAs Potentially Controlling Cell Cycle in the Model *Caulobacter crescentus*: A Bioinformatic Approach. *Frontiers in Genetics*, Frontiers, 2018, 9, <10.3389/fgene.2018.00164>. <hal-01916083>

HAL Id: hal-01916083

<https://hal-amu.archives-ouvertes.fr/hal-01916083>

Submitted on 8 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Non-coding RNAs Potentially Controlling Cell Cycle in the Model *Caulobacter crescentus*: A Bioinformatic Approach

Wanassa Beroual¹, Matteo Brilli² and Emanuele G. Biondi^{1*}

¹ Laboratoire de Chimie Bactérienne, Centre National de la Recherche Scientifique, Aix Marseille University, Marseille, France,

² Department of Biosciences, Pediatric Clinical Research Center "Romeo ed Enrica Invernizzi," University of Milan, Milan, Italy

Caulobacter crescentus represents a remarkable model system to investigate global regulatory programs in bacteria. In particular, several decades of intensive study have revealed that its cell cycle is controlled by a cascade of master regulators, such as DnaA, GcrA, CcrM, and CtrA, that are responsible for the activation of functions required to progress through DNA replication, cell division and morphogenesis of polar structures (flagellum and stalk). In order to accomplish this task, several post-translational (phosphorylation and proteolysis) and transcriptional mechanisms are involved. Surprisingly, the role of non-coding RNAs (ncRNAs) in regulating the cell cycle has not been investigated. Here we describe a bioinformatic analysis that revealed that ncRNAs may well play a crucial role regulating cell cycle in *C. crescentus*. We used available prediction tools to understand which target genes may be regulated by ncRNAs in this bacterium. Furthermore, we predicted whether ncRNAs with a cell cycle regulated expression profile may be directly regulated by DnaA, GcrA, and CtrA, at the onset, during or end of the S-phase/swarmer cell, or if any of them has CcrM methylation sites in the promoter region. Our analysis suggests the existence of a potentially very important network of ncRNAs regulated by or regulating well-known cell cycle genes in *C. crescentus*. Our hypothesis is that ncRNAs are intimately connected to the known regulatory network, playing a crucial modulatory role in cell cycle progression.

Keywords: *Caulobacter crescentus*, Cell cycle regulation, ncRNAs, DNA replication, Two-component regulatory systems (TCS)

INTRODUCTION

In the last two decades, bacterial non-coding RNAs (ncRNAs) and small RNAs in particular (sRNAs, 50–400 nts), have emerged as central regulators of important cellular processes (Dutta and Srivastava, 2018). Most sRNAs are post-transcriptional regulators positively or negatively affecting the translation and/or stability of their targets. As a result, sRNAs have been shown to play key roles in the adaptive response to the environment and to stress conditions, in particular. In *Enterobacteria*, most of the proficient sRNAs/target pairs requires the RNA chaperone Hfq that both stabilizes them and facilitates the RNA duplex formation; however, this role of Hfq in sRNA control of gene expression is not conserved in all bacteria that possess an Hfq homolog (Vogel and Luisi, 2011). While sRNAs are best characterized in enterobacteria (*Escherichia coli* and

OPEN ACCESS

Edited by:

Alessio Mengoni,
Università Degli Studi di Firenze, Italy

Reviewed by:

Francesco Russo,
University of Copenhagen, Denmark
Jean-Yves Matroule,
University of Namur, Belgium

*Correspondence:

Emanuele G. Biondi
ebiondi@imm.cnrs.fr

Citation:

Beroual W, Brilli M and Biondi EG
(2018) Non-coding RNAs Potentially
Controlling Cell Cycle in the Model
Caulobacter crescentus: A
Bioinformatic Approach.
Front. Genet. 9:164.
doi: 10.3389/fgene.2018.00164

in stalk cells and fire a single round of DNA replication thanks to the protein DnaA, which binds the unique origin of replication and activates the DNA polymerase complex (Collier, 2012; Felletti et al., 2018). Besides controlling the replication of the chromosome, DnaA controls the transcription of the gene encoding GcrA, which in turn controls many essential genes during S-phase, including *ctrA* (Holtzendorff et al., 2004; Hottes et al., 2005; Collier et al., 2006). This latter gene codes for a response regulator that activates cell division and the expression of genes essential for cell differentiation, such as those for flagellum/pilum assembly, chemotaxis, stalk biogenesis and many others (Reisenauer et al., 1999; Jones et al., 2001; Laub et al., 2002; Biondi et al., 2006). DnaA, GcrA and CtrA constitute an essential transcriptional cascade that requires multiple regulatory levels in order to ensure the correct timing of ensuing events. For example, GcrA activity depends on methylation of the chromosome by the methyl-transferase CcrM (Fioravanti et al., 2013; Murray et al., 2013; Mohapatra et al., 2014; Haakonsen et al., 2015); CtrA's activity is regulated by phosphorylation (Biondi et al., 2006) and inactivated by a ClpXP-dependent degradation (Ryan et al., 2002, 2004; Joshi et al., 2015). Among well-known regulatory layers, surprisingly, the activity and the role of ncRNAs in *C. crescentus* has been poorly investigated, and still, they represent ideal candidates for such regulations, as they provide dynamic patterns not easily achievable with transcriptional regulation only. In 2008, a paper entitled "Small non-coding RNAs in *Caulobacter crescentus*" described 27 ncRNAs in this organism (Landt et al., 2008). Unfortunately only few of them appeared to be involved in important functions and, besides a work in 2010 describing CrfA (Landt et al., 2010), an sRNA involved in adaptation to carbon starvation, no other ncRNA identified in 2008 was further characterized. More recently another sRNA was characterized in *C. crescentus* and named GsrN (Tien et al., 2017). This ncRNA is involved in multiple stresses response factor directly controlled by the general stress sigma factor, σ^T . Finally new recent approaches using RNAseq and post-genomic techniques expanded the *plethora* of ncRNA candidates in this species to over 100 (Zhou et al., 2015).

This observation prompted us to start a systematic investigation of this new ncRNA world aiming to identify global regulators and ncRNAs possibly involved in the cell cycle in *C. crescentus*. We applied several bioinformatics tools in order to understand what kind of functions may be controlled by ncRNAs and we focused on uncharacterized ncRNAs with a changing expression level during cell cycle. We predicted their targets in the *C. crescentus* genome and we integrate this information with motif scanning and ChIP-Seq data.

METHODS

Prediction of Targets of ncRNAs

Sequences of ncRNAs were retrieved by the annotation of genes by previous results (Schrader et al., 2014; Zhou et al., 2015). PredatorRNA (Eggenhofer et al., 2011). was used using the genome NC_011916, as deposited in the PredatorRNA

website, and prediction was performed as default. TargetRNA2 (Kery et al., 2014) was performed using the following settings: NTs before start codon 80, NTs after start codon 20, Seed length 7, sRNA conservation and accessibility true, sRNA window size 13, mRNA structural accessibility true, Interaction region 20, Filter size 1,000, *P*-value threshold 0.5 (predictions were considered significant only with a *P* < 0.05). Finally CopraRNA (Wright et al., 2014) was used with default settings.

Secondary structure prediction of ncRNAs was performed using mFold (Zuker, 2003) and RNAfold (Gruber et al., 2008) using default settings.

Prediction of DnaA, CcrM, and CtrA Consensus Sequences

Predictions of DnaA, CcrM, and CtrA potential controls were performed as previously described (Brilli et al., 2010). Position Weight Matrices (PWM) modeling the CtrA and DnaA binding sites were used to scan the entire genome by calculating a measure of similarity for each genome position with the formula:

$$S_{i:i+L} = \frac{1}{L} \sum_{n=i}^{i+L} (2 + \log_2 f_{nx}).$$

From Schneider et al. (1986). Basically, for each position *n* in a genomic window of length *L*, where *L* is also the length of the transcription factor binding motif, we sum the logarithm base 2 of the frequency of nucleotide *x* at position *n* of the motif and then we average over all nucleotides.

As this score is continuous, there is the need to establish a threshold, that can be quite arbitrary. In this context, we normalize all the scores with respect to the maximum score attainable by the PWM under analysis, and we only retain scores that are at least 60% of the maximum.

ChIPseq data of GcrA were analyzed as previously described (Fioravanti et al., 2013) using the new annotation of ncRNAs (Zhou et al., 2015).

RESULTS

Selection of Cell Cycle Regulated ncRNAs

Previous works based on total RNA sequencing and 5'-RACE have identified a list of predicted ncRNAs expressed in culture conditions (Schrader et al., 2014; Zhou et al., 2015). Based on this experimental analysis we recovered all sequences previously identified and we initially separated ncRNA candidates based on previous annotation and their dynamic regulation during cell cycle. Out of 199, identified as ncRNAs expressed in rich or poor media, respectively PYE and M2G, 88 were further characterized for their Transcriptional Start Site (TSS), with few of them having multiple transcription start sites (Zhou et al., 2015). Among those ncRNAs, 43 were related to the translational machinery (tRNAs or ribosome-related), therefore we excluded them in the following analyses, as well as the TmRNA (Keiler and Shapiro, 2003; Cheng and Keiler, 2009; Russell and Keiler, 2009). The bibliographical data mining

TABLE 1 | ncRNAs in *Caulobacter crescentus*.

CCNA	Multiple TSS	Cell cycle phase G1, S, G2	TSS location	strand	0 min	20 min	40 min	60 min	80 min	100 min	120 min	140 min	Annotation based on Zhou et al. (2015)
CCNA_R0004	/	/	210993	-	5.5	3.9	8.0	4.8	12.8	2.2	3.7	4.2	Cell cycle regulated
CCNA_R0009	/	/	708218	+	105.7	136.7	42.0	88.7	64.2	92.4	93.0	58.9	Small non-coding RNA
CCNA_R0016	/	/	844332	+	32.2	19.0	43.0	63.0	59.6	28.2	61.4	46.6	Small non-coding RNA
CCNA_R0019	/	/	920752	-	11.3	2.9	1.0	1.3	1.0	8.4	1.8	4.2	Stationary phase
CCNA_R0025	3	/	1176006	+	214.0	13.2	262.0	360.4	275.7	10.1	356.7	551.2	Small non-coding RNA
CCNA_R0040	/	/	1645202	+	7.8	7.9	16.0	14.3	7.7	20.0	17.9	19.4	Small non-coding RNA
CCNA_R0063	/	/	2889177	+	261.6	112.8	312.0	266.9	192.6	215.9	341.6	207.5	Minimal medium
CCNA_R0111	/	/	622892	+	0.8	1.4	5.0	0.4	0.0	2.7	0.5	3.9	Small non-coding RNA
CCNA_R0112	/	/	626656	+	121.1	143.6	101.0	118.7	109.4	66.8	139.2	74.8	Small non-coding RNA
CCNA_R0134	/	/	1294595	+	12.7	42.8	67.0	28.7	64.7	25.8	70.1	70.2	Small non-coding RNA
CCNA_R0137	/	/	1387892	-	43.3	45.3	21.0	43.5	20.0	68.8	39.8	63.5	Small non-coding RNA
CCNA_R0141	/	/	1607996	-	7.4	0.2	1.0	7.4	0.0	1.0	3.7	7.8	Small non-coding RNA
CCNA_R0142	/	/	1682820	+	13.3	16.8	4.0	6.5	3.1	5.3	8.2	15.9	Small non-coding RNA
CCNA_R0145	/	/	1911195	+	5.3	10.1	1.0	6.5	10.8	13.8	10.5	11.3	Small non-coding RNA
CCNA_R0147	/	/	1986625	+	129.1	31.7	77.0	50.4	66.2	8.2	27.9	14.5	Small non-coding RNA
CCNA_R0170	/	/	2778348	+	15.8	15.7	18.0	30.9	23.6	18.1	27.0	21.2	Small non-coding RNA
CCNA_R0171	/	/	2800063	+	0.4	5.8	0.0	7.0	8.2	3.4	5.5	3.9	Small non-coding RNA
CCNA_R0173	/	/	2967588	-	9.8	16.6	0.0	15.6	10.8	29.0	10.5	13.1	Small non-coding RNA
CCNA_R0174	/	/	2977449	+	31.6	40.9	20.0	41.7	12.3	35.9	38.9	25.4	Small non-coding RNA
CCNA_R0182	/	/	3288242	+	94.8	93.8	130.0	201.3	120.2	118.0	200.1	115.7	Small non-coding RNA
CCNA_R0183	/	/	3294320	+	17.9	1.0	1.0	10.0	2.6	0.0	1.8	2.5	Small non-coding RNA
CCNA_R0188	/	/	3462949	+	117.2	86.2	40.0	104.3	70.3	47.3	76.0	104.8	Small non-coding RNA
CCNA_R0193	/	/	3699499	+	214.6	72.0	134.0	130.4	118.6	75.3	132.8	121.0	Small non-coding RNA
CCNA_R0018	/	G1	920425	+	86.2	24.6	10.0	48.3	49.8	18.8	32.1	25.4	Rich medium
CCNA_R0051	/	G1	2404309	+	281.3	162.6	148.0	140.8	102.7	203.6	141.9	91.7	Small non-coding RNA
CCNA_R0095	/	G1	4950	-	99.3	6.2	18.0	36.1	22.6	13.5	19.2	12.7	Small non-coding RNA
CCNA_R0123	/	G1	921719	-	9.2	4.6	1.0	3.9	2.6	1.4	0.5	4.2	Small non-coding RNA
CCNA_R0176	/	G1	3091482	-	231.0	41.2	8.0	3.9	2.6	10.1	23.8	16.6	Small non-coding RNA
CCNA_R0199	/	G1	3987216	+	89.0	15.4	14.0	28.3	21.1	16.6	23.4	52.2	Small non-coding RNA
CCNA_R0025	1	G1-G2	1176026	+	139.7	81.3	59.0	83.9	86.8	49.9	131.0	150.3	Small non-coding RNA
CCNA_R0025	2	G1-G2	1176040	-	500.6	58.7	3.0	12.2	42.1	56.9	254.1	392.0	Small non-coding RNA
CCNA_R0164	/	G1-G2	2608458	-	34.3	26.2	10.0	12.2	22.6	46.8	40.8	26.8	Small non-coding RNA
CCNA_R0050	/	G1-S	2397735	+	105.5	700.9	385.0	306.0	167.9	99.2	103.9	64.2	Cell cycle regulated
CCNA_R0117	/	G1-S	772628	+	30.6	84.5	70.0	110.0	49.8	49.7	62.7	73.4	Small non-coding RNA
CCNA_R0104	/	G2	362212	-	4.5	0.2	0.0	4.8	7.7	6.0	17.4	22.6	Small non-coding RNA
CCNA_R0165	/	G2	2672357	-	25.2	10.8	2.0	10.9	18.5	42.2	78.8	43.8	Small non-coding RNA
CCNA_R0093	/	S	3781500	-	8141.7	16080.3	26189.0	16986.4	21895.1	19086.7	20603.8	15762.8	Minimal medium
CCNA_R0094	/	S	182	-	55.0	15.4	2.0	46.1	181.3	314.6	176.3	118.9	Small non-coding RNA
CCNA_R0116	/	S	757263	-	11.1	276.7	480.0	793.3	448.8	325.5	180.9	190.9	Small non-coding RNA
CCNA_R0124	/	S	938932	+	98.9	180.7	268.0	390.4	297.3	309.6	220.7	290.8	Small non-coding RNA
CCNA_R0126	/	S	1081336	-	1.4	5.5	6.0	15.6	10.3	12.8	9.6	4.9	Small non-coding RNA
CCNA_R0139	/	S	1523958	+	5.7	16.9	25.0	19.6	22.6	16.6	27.9	18.7	Small non-coding RNA
CCNA_R0152	/	S	2086425	+	447.7	526.2	416.0	1091.9	547.4	439.4	580.6	777.0	Small non-coding RNA
CCNA_R0159	/	S	2434068	+	1.0	12.3	15.0	25.6	14.9	16.9	18.3	15.2	Small non-coding RNA
CCNA_R0163	/	S	2560042	-	4.9	84.7	108.0	134.3	78.0	56.0	30.2	55.0	Small non-coding RNA
CCNA_R0172	/	S	2940670	-	51.7	71.5	210.0	159.5	181.3	146.0	173.5	109.0	Small non-coding RNA

Coordinates of the Transcriptional Start Site (TSS) and classification of the cell cycle regulated phase are listed.

therefore allowed us to select 44 ncRNAs for further analysis. Among them, as previously described in the introduction, two ncRNAs were already characterized and named CrfA (Landt et al., 2008, 2010) and GsrN (Tien et al., 2017). In conclusion 42 ncRNAs were the object of this study (Figure 1 and Table 1).

By studying the expression patterns of the selected ncRNA in synchronized cells, we classified those 42 ncRNAs based on their expression levels during cell cycle. Specifically, we identified 23 ncRNAs whose expression changes during the cell cycle: 6 are expressed in G1, 2 at the onset of the S phase (noted as G1-S), CCNA_R0025 and CCNA_R0164 are expressed in G1 and G2 phases, while 10 are expressed in S-phase (noted as S) and 2 are toward the end of the cell cycle (G2). The latter genes may be reflecting the accumulation in a specific cell type, as we will discuss later. The observation that some ncRNAs genes are cell cycle regulated, may suggest a putative function associated to regulation of functions that are required at a specific phase (Laub et al., 2000).

Prediction of Target Genes Regulated by ncRNAs

Although experimental validation is absolutely required to identify targets of ncRNAs, predictive tools may be useful to suggest candidate target genes and therefore give a glance about the functions regulated by a set of sRNAs. This is particularly valid in case of a systematic analysis as the experimental validation/identification of targets is not easily scalable. In order to reconstitute the whole ncRNAs network connected to cell cycle, we used three different available tools, mainly RNAPredator (Eggenhofer et al., 2011) but then as confirmation also CopraRNA (Backofen et al., 2014; Wright et al., 2014) and TargetRNA2 (Kery et al., 2014). Each analysis provides specific features and it is able to predict classes of targets. By using PredatorRNA we predicted the first 100 targets that were retained for the analysis (Supplementary Table 1).

Genes coding for factors playing a role in cell cycle regulation were suggested as probable targets of some of the ncRNAs (Table 2). In particular we considered genes annotated as of cell cycle/cell division regulators. In order to evaluate the predictions, we performed an additional analysis based on the following consideration: regions that are complementary to the target RNA should be located within open regions (loops) of the ncRNA molecule and not to paired regions (stems) to ensure accessibility. Therefore, we aimed to understand *in silico* how those putative small RNAs were structured and whether targets were interacting with the same loop regions. We used RNAfold (Gruber et al., 2008) and mFold (Zuker, 2003) in order to identify loop regions that would be more accessible for the interaction with target genes.

We asked whether among cell cycle-regulated ncRNAs target genes were more likely being cell cycle regulators with respect to non-cell cycle regulated ncRNAs. Among the 22 ncRNAs that showed dynamic expression during the cell cycle, we found 13

TABLE 2 | Prediction based on PredatorRNA, confirmed by Copra and/or TargetRNA2 (regions targeted by ncRNAs are defined in Supplementary Table 1).

CCNA	Cell cycle phase G1, S, G2	Functions:	Significant genes (genes for "flagellum" are not listed):
CCNA_R0004	/	Pilus, flagellum, cell division	<i>CpaE, maf</i>
CCNA_R0009	/	Flagellum	
CCNA_R0016	/	Cell cycle	<i>hdaA</i>
CCNA_R0019	/	Flagellum	
CCNA_R0025	/	Flagellum, ppGpp	<i>spoT</i>
CCNA_R0040	/	Cell division	<i>ftsK</i>
CCNA_R0063	/	Cell division	<i>ftsH</i>
CCNA_R0111	/	Flagellum, pilus, cell division	<i>cpaE, cpaF, ftsK</i>
CCNA_R0112	/	/	
CCNA_R0134	/	Chromosome partitioning, cell division	<i>mipZ, ftsY</i>
CCNA_R0137	/	/	
CCNA_R0141	/	Cell division, cell cycle	<i>ftsH, tipN</i>
CCNA_R0142	/	/	
CCNA_R0145	/	Cell cycle, flagellum	<i>hdaA, divL</i>
CCNA_R0147	/	Cell division	<i>ftsE</i>
CCNA_R0170	/	Cell cycle, flagellum	<i>hdaA</i>
CCNA_R0171	/	Cell cycle, cell division, pilus	<i>divJ, cpdR, mraZ, cpaD</i>
CCNA_R0173	/	Cell cycle	<i>podJ</i>
CCNA_R0174	/	Flagellum	
CCNA_R0182	/	/	
CCNA_R0183	/	Cell division, flagellum	<i>ftsI</i>
CCNA_R0188	/	Pilus, cell cycle	<i>cpaE, divL, hdaA</i>
CCNA_R0193	/	Flagellum	
CCNA_R0018	G1	Flagellum, cell division	<i>ftsW</i>
CCNA_R0051	G1	Flagellum	
CCNA_R0095	G1	Pilus, flagellum, cell cycle, cell division, Chromosome partitioning	<i>parA, cckA, ftsH, cpaE</i>
CCNA_R0123	G1	Cell cycle, flagellum	<i>dnaA</i>
CCNA_R0176	G1	Cell cycle	<i>dnaA, divJ</i>
CCNA_R0199	G1	Stalk	<i>shkA</i>
CCNA_R0025	G1-G2	Flagellum, ppGpp	<i>spoT</i>
CCNA_R0025	G1-G2	Flagellum, ppGpp	<i>spoT</i>
CCNA_R0164	G1-G2	Flagellum	
CCNA_R0050	G1-S	Cell cycle, flagellum	<i>divJ, pleD, tipN</i>
CCNA_R0117	G1-S	Flagellum	
CCNA_R0104	G2	Cell cycle	<i>dnaA, cenR</i>
CCNA_R0165	G2	Cell cycle, flagellum, ppGpp	<i>popZ, spoT</i>
CCNA_R0093	S	Cell cycle	<i>divL</i>
CCNA_R0094	S	Cell cycle	<i>dnaA</i>
CCNA_R0116	S	Stalk	<i>shkA</i>
CCNA_R0124	S	Cell cycle	<i>hdaA</i>
CCNA_R0126	S	/	
CCNA_R0139	S	Pilus, Flagellum, cell cycle	<i>cpaF, divL, tacA, pleD</i>
CCNA_R0152	S	/	
CCNA_R0159	S	Cell cycle	<i>dnaA</i>
CCNA_R0163	S	Cell cycle	<i>kidO</i>
CCNA_R0172	S	Cell cycle	<i>chpT</i>

TABLE 3 | Prediction of DnaA and CtrA binding sites, GcrA ChIPseq peaks and CcrM methylation sites.

Name	TSS position		region* = TSS – 100; TSS + 50			DnaA	GcrA peaks (ChIPseq) (Fioravanti et al., 2013)
			CtrA Full	CtrA half	CcrM		
CCNA_R0004	210993	–	0	1	0	0	0
CCNA_R0009	708218	+	0	4	0	0	1
CCNA_R0016	844332	+	0	0	0	1	0
CCNA_R0018	920425	+	0	0	0	0	0
CCNA_R0019	920752	–	0	0	0	0	1
CCNA_R0025	1176006	+	1	2	0	0	0
CCNA_R0025	1176026	+	1	2	0	0	0
CCNA_R0025	1176040	–	0	1	1	0	0
CCNA_R0040	1645202	+	0	0	2	0	0
CCNA_R0050	2397735	+	0	0	0	0	0
CCNA_R0051	2404309	+	0	1	0	0	0
CCNA_R0063	2889177	+	0	0	0	0	0
CCNA_R0093	3781500	–	0	0	0	0	1
CCNA_R0094	182	–	2	4	1	0	0
CCNA_R0095	4950	–	0	0	0	0	0
CCNA_R0104	362212	–	0	2	0	0	0
CCNA_R0111	622892	+	0	0	0	0	0
CCNA_R0112	626656	+	0	1	1	0	0
CCNA_R0116	757263	–	0	1	3	0	1
CCNA_R0117	772628	+	0	0	1	0	0
CCNA_R0123	921719	–	0	0	0	0	0
CCNA_R0124	938932	+	0	3	0	0	0
CCNA_R0126	1081336	–	0	0	1	0	0
CCNA_R0134	1294595	+	0	0	0	0	0
CCNA_R0137	1387892	–	0	1	0	0	0
CCNA_R0139	1523958	+	0	0	0	1	0
CCNA_R0141	1607996	–	0	0	0	0	0
CCNA_R0142	1682820	+	0	0	0	0	0
CCNA_R0145	1911195	+	0	0	1	0	0
CCNA_R0147	1986625	+	0	0	0	0	0
CCNA_R0152	2086425	+	0	1	2	0	0
CCNA_R0159	2434068	+	0	0	2	1	0
CCNA_R0163	2560042	–	0	1	2	0	1
CCNA_R0164	2608458	–	0	0	0	0	0
CCNA_R0165	2672357	–	0	1	2	0	0
CCNA_R0170	2778348	+	0	0	2	0	0
CCNA_R0171	2800063	+	2	4	0	0	0
CCNA_R0172	2940670	–	0	0	1	1	0
CCNA_R0173	2967588	–	0	1	0	1	0
CCNA_R0174	2977449	+	0	0	0	0	1
CCNA_R0176	3091482	–	0	0	1	0	0
CCNA_R0182	3288242	+	0	0	0	0	0
CCNA_R0183	3294320	+	0	0	0	0	0
CCNA_R0188	3462949	+	0	0	0	0	0
CCNA_R0193	3699499	+	0	0	0	1	0
CCNA_R0199	3987216	+	0	2	1	2	0

*Opposite for genes annotated on the minus strand.




FIGURE 2 | Network of ncRNAs and master regulators of cell cycle. Integrated view of control on selected ncRNAs by DnaA, CtrA, CcrM, and GcrA and of ncRNA on transcripts. Violet squares represent ncRNAs. CtrA, DnaA, CcrM, and GcrA (Upper Case) represent proteins, while the same factors (lower case) represent targets of ncRNAs. For CtrA blue arrows represent “half sites” while green arrows represent “full sites” (whenever a full site is present, we don't indicate half sites).

out of 22 directly acting on cell cycle regulators (ca 60%), while only 7 out of 23 (30%) of the remaining ncRNAs, not showing cell cycle regulation, have targets with a role in the cell cycle. This observation suggests that ncRNAs may be indeed cell cycle-regulated as they act on functions that need to be activated only at specific phases of cell cycle progression. But can we identify whether ncRNAs are regulated by one of the master regulators of the cell cycle?

Presence of DnaA and CtrA Boxes or Methylation Sites Upstream Cell Cycle-Regulated ncRNAs

In order to understand how ncRNAs are regulated by the cell cycle we explored the presence of known binding sites in their promoter regions. Specifically promoters were scanned for the presence of DnaA, CcrM (GAnTC) or CtrA putative DNA binding sites (both “half” and “full” binding sites) (Ouimet and Marczyński, 2000; Brilli et al., 2010). We also analyzed the presence of GcrA binding sites using previously published Chromatin Immunoprecipitation-deep sequencing (ChIP-seq) data (Fioravanti et al., 2013; Murray et al., 2013). Those four regulators represent the core transcriptional machinery responsible for cell cycle progression and for the coordinated expression of all the cell cycle-dependent functions. The analysis

revealed that many of those ncRNAs are potentially regulated by master regulators of cell cycle (Table 3). In particular ncRNAs expressed in S-phase appear to possess binding sites of master regulators, justifying their cell regulated expression. Although the expression pattern may be the result of combinatorial regulation, this enrichment of cell cycle regulators binding motifs is intriguing and asks for experimental validation.

In order to understand the full picture of interconnection between ncRNAs and master regulators, we integrated the data of Table 2 (cell cycle targets of ncRNAs) with those in Table 3 (master regulators regulating ncRNAs) and we represent the regulations as a network (Figure 2). This network represents ncRNAs that are potentially regulated by master regulator of cell cycle and that are eventually connected to cell cycle regulator genes.

CONCLUSIONS AND PERSPECTIVES

Our comprehensive analysis of *C. crescentus* ncRNAs has revealed that many of those factors potentially play important roles during the cell cycle. In particular, our predictions show that many of them target the UTRs of *dnaA* or *hdaA*. Initiation of DNA replication is a fundamental event during the cell cycle and for this reason, it likely justifies multiple regulation levels.

DnaA has been shown to be subject to multiple regulatory layers (Felletti et al., 2018), including a ncRNA named SsrA, or tmRNA, a ncRNA interacting with ribosomes to regulate protein degradation (Keiler and Shapiro, 2003). Here we indicate for the first time several ncRNAs able to potentially regulate *dnaA*.

However the extent of ncRNAs regulation on the cell cycle is even vaster (Figure 2). Clearly, CtrA and CcrM seem to play a major role in coordinating the expression of many genes, while *dnaA* seems to be a major target of at least four ncRNAs. Several ncRNAs seem to be controlled by two master regulators and R0116 is controlled by CcrM, GcrA, and CtrA, suggesting an important role for this ncRNA.

Although this analysis is based on predictions and every connection must be validated experimentally, the amplitude of all connections revealed by this bioinformatic predictions strongly suggests that ncRNAs are indeed playing a major role in cell cycle regulation of *C. crescentus*. Next years will reveal whether this preliminary analysis grasped this important role.

REFERENCES

- Backofen, R., Amman, F., Costa, F., Findeiß, S., Richter, A. S., and Stadler, P. F. (2014). Bioinformatics of prokaryotic RNAs. *RNA Biol.* 11, 470–483. doi: 10.4161/rna.28647
- Barquist, L., and Vogel, J. (2015). Accelerating discovery and functional analysis of small RNAs with new technologies. *Annu. Rev. Genet.* 49, 367–394. doi: 10.1146/annurev-genet-112414-054804
- Biondi, E. G., Reisinger, S. J., Skerker, J. M., Arif, M., Perchuk, B. S., Ryan, K. R., et al. (2006). Regulation of the bacterial cell cycle by an integrated genetic circuit. *Nature* 444, 899–904. doi: 10.1038/nature05321
- Brilli, M., Fondi, M., Fani, R., Mengoni, A., Ferri, L., Bazzicalupo, M., et al. (2010). The diversity and evolution of cell cycle regulation in alpha-proteobacteria: a comparative genomic analysis. *BMC Syst. Biol.* 4:52. doi: 10.1186/1752-0509-4-52
- Cheng, L., and Keiler, K. C. (2009). Correct timing of *dnaA* transcription and initiation of DNA replication requires trans translation. *J. Bacteriol.* 191, 4268–4275. doi: 10.1128/JB.00362-09
- Collier, J. (2012). Regulation of chromosomal replication in *Caulobacter crescentus*. *Plasmid* 67, 76–87. doi: 10.1016/j.plasmid.2011.12.007
- Collier, J., Murray, S. R., and Shapiro, L. (2006). DnaA couples DNA replication and the expression of two cell cycle master regulators. *EMBO J.* 25, 346–356. doi: 10.1038/sj.emboj.7600927
- Dutta, T., and Srivastava, S. (2018). Small RNA-mediated regulation in bacteria: a growing palette of diverse mechanisms. *Gene.* 656, 60–72. doi: 10.1016/j.gene.2018.02.068
- Eggenhofer, F., Tafer, H., Stadler, P. F., and Hofacker, I. L. (2011). RNApredator: fast accessibility-based prediction of sRNA targets. *Nucleic Acids Res.* 39, W149–W154. doi: 10.1093/nar/gkr467
- Felletti, M., Omnus, D. J., and Jonas, K. (2018). Regulation of the replication initiator DnaA in *Caulobacter crescentus*. *Biochim. Biophys. Acta* S1874–9399(17)30295-X. doi: 10.1016/j.bbagr.2018.01.004
- Fioravanti, A., Fumeaux, C., Mohapatra, S. S., Bompard, C., Brilli, M., Frandi, A., et al. (2013). DNA binding of the cell cycle transcriptional regulator GcrA depends on N6-adenosine methylation in *Caulobacter crescentus* and other *Alphaproteobacteria*. *PLoS Genet.* 9:e1003541. doi: 10.1371/journal.pgen.1003541
- Gruber, A. R., Lorenz, R., Bernhart, S. H., Neuböck, R., and Hofacker, I. L. (2008). The Vienna RNA websuite. *Nucleic Acids Res.* 36, W70–W74. doi: 10.1093/nar/gkn188

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

ACKNOWLEDGMENTS

We would like to thank Nadia Benzaina and Maxence Vincent for their continuous enthusiasm and support. We would like to thank also José Ignacio Jiménez Zurdo, Marta Robledo Garrido, Maude Guillier, Eric Massé and David Lalaouna for long constructive discussions on ncRNAs.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fgene.2018.00164/full#supplementary-material>

Supplementary Table 1 | Results of predictions by RNA predator (see Materials and Methods).

- Haakonsen, D. L., Yuan, A. H., and Laub, M. T. (2015). The bacterial cell cycle regulator GcrA is a $\sigma 70$ cofactor that drives gene expression from a subset of methylated promoters. *Genes Dev.* 29, 2272–2286. doi: 10.1101/gad.270660.115
- Holtzendorff, J., Hung, D., Brende, P., Reisenauer, A., Viollier, P. H., McAdams, H. H., et al. (2004). Oscillating global regulators control the genetic circuit driving a bacterial cell cycle. *Science* 304, 983–987. doi: 10.1126/science.1095191
- Hottes, A. K., Shapiro, L., and McAdams, H. H. (2005). DnaA coordinates replication initiation and cell cycle transcription in *Caulobacter crescentus*. *Mol. Microbiol.* 58, 1340–1353. doi: 10.1111/j.1365-2958.2005.04912.x
- Jones, S. E., Ferguson, N. L., and Alley, M. R. (2001). New members of the *ctrA* regulon: the major chemotaxis operon in *Caulobacter* is CtrA dependent. *Microbiol. Read. Engl.* 147, 949–958. doi: 10.1099/00221287-147-4-949
- Joshi, K. K., Bergé, M., Radhakrishnan, S. K., Viollier, P. H., and Chien, P. (2015). An adaptor hierarchy regulates proteolysis during a bacterial cell cycle. *Cell* 163, 419–431. doi: 10.1016/j.cell.2015.09.030
- Keiler, K. C., and Shapiro, L. (2003). TmRNA is required for correct timing of DNA replication in *Caulobacter crescentus*. *J. Bacteriol.* 185, 573–580. doi: 10.1128/JB.185.2.573-580.2003
- Kery, M. B., Feldman, M., Livny, J., and Tjaden, B. (2014). TargetRNA2: identifying targets of small regulatory RNAs in bacteria. *Nucleic Acids Res.* 42, W124–W129. doi: 10.1093/nar/gku317
- Landt, S. G., Abeliuk, E., McGrath, P. T., Lesley, J. A., McAdams, H. H., and Shapiro, L. (2008). Small non-coding RNAs in *Caulobacter crescentus*. *Mol. Microbiol.* 68, 600–614. doi: 10.1111/j.1365-2958.2008.06172.x
- Landt, S. G., Lesley, J. A., Britos, L., and Shapiro, L. (2010). CrfA, a small noncoding RNA regulator of adaptation to carbon starvation in *Caulobacter crescentus*. *J. Bacteriol.* 192, 4763–4775. doi: 10.1128/JB.00343-10
- Lasker, K., Mann, T. H., and Shapiro, L. (2016). An intracellular compass spatially coordinates cell cycle modules in *Caulobacter crescentus*. *Curr. Opin. Microbiol.* 33, 131–139. doi: 10.1016/j.mib.2016.06.007
- Laub, M. T., Chen, S. L., Shapiro, L., and McAdams, H. H. (2002). Genes directly controlled by CtrA, a master regulator of the *Caulobacter* cell cycle. *Proc. Natl. Acad. Sci. U.S.A.* 99, 4632–4637. doi: 10.1073/pnas.062065699
- Laub, M. T., McAdams, H. H., Feldblyum, T., Fraser, C. M., and Shapiro, L. (2000). Global analysis of the genetic network controlling a bacterial cell cycle. *Science* 290, 2144–2148. doi: 10.1126/science.290.5499.2144
- Mohapatra, S. S., Fioravanti, A., and Biondi, E. G. (2014). DNA methylation in *Caulobacter* and other *Alphaproteobacteria* during cell cycle progression. *Trends Microbiol.* 22, 528–535. doi: 10.1016/j.tim.2014.05.003

- Murray, S. M., Panis, G., Fumeaux, C., Viollier, P. H., and Howard, M. (2013). Computational and genetic reduction of a cell cycle to its simplest, primordial components. *PLoS Biol.* 11:e1001749. doi: 10.1371/journal.pbio.1001749
- Ouimet, M. C., and Marczyński, G. T. (2000). Analysis of a cell-cycle promoter bound by a response regulator. *J. Mol. Biol.* 302, 761–775. doi: 10.1006/jmbi.2000.4500
- Reisenauer, A., Quon, K., and Shapiro, L. (1999). The CtrA response regulator mediates temporal control of gene expression during the *Caulobacter* cell cycle. *J. Bacteriol.* 181, 2430–2439.
- Russell, J. H., and Keiler, K. C. (2009). Subcellular localization of a bacterial regulatory RNA. *Proc. Natl. Acad. Sci. U.S.A.* 106, 16405–16409. doi: 10.1073/pnas.0904904106
- Ryan, K. R., Huntwork, S., and Shapiro, L. (2004). Recruitment of a cytoplasmic response regulator to the cell pole is linked to its cell cycle-regulated proteolysis. *Proc. Natl. Acad. Sci. U.S.A.* 101, 7415–7420. doi: 10.1073/pnas.0402153101
- Ryan, K. R., Judd, E. M., and Shapiro, L. (2002). The CtrA response regulator essential for *Caulobacter crescentus* cell-cycle progression requires a bipartite degradation signal for temporally controlled proteolysis. *J. Mol. Biol.* 324, 443–455. doi: 10.1016/S0022-2836(02)01042-2
- Schneider, T. D., Stormo, G. D., Gold, L., and Ehrenfeucht, A. (1986). Information content of binding sites on nucleotide sequences. *J. Mol. Biol.* 188, 415–431.
- Schrader, J. M., and Shapiro, L. (2015). Synchronization of *Caulobacter crescentus* for investigation of the bacterial cell cycle. *J. Vis. Exp.* doi: 10.3791/52633
- Schrader, J. M., Zhou, B., Li, G.-W., Lasker, K., Childers, W. S., Williams, B., et al. (2014). The coding and noncoding architecture of the *Caulobacter crescentus* genome. *PLoS Genet.* 10:e1004463. doi: 10.1371/journal.pgen.1004463
- Tien, M. Z., Fiebig, A., and Crosson, S. (2017). Gene network analysis identifies a central post-transcriptional regulator of cellular stress survival. *bioRxiv*, 212902. doi: 10.1101/212902
- Vogel, J., and Luisi, B. F. (2011). Hfq and its constellation of RNA. *Nat. Rev. Microbiol.* 9, 578–589. doi: 10.1038/nrmicro2615
- Wright, P. R., Georg, J., Mann, M., Sorescu, D. A., Richter, A. S., Lott, S., et al. (2014). CopraRNA and IntaRNA: predicting small RNA targets, networks and interaction domains. *Nucleic Acids Res.* 42, W119–W123. doi: 10.1093/nar/gku359
- Zhou, B., Schrader, J. M., Kalogeraki, V. S., Abeliuk, E., Dinh, C. B., Pham, J. Q., et al. (2015). The global regulatory architecture of transcription during the *Caulobacter* cell cycle. *PLoS Genet.* 11:e1004831. doi: 10.1371/journal.pgen.1004831
- Zuker, M. (2003). Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Res.* 31, 3406–3415. doi: 10.1093/nar/gkg595

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Beroual, Brilli and Biondi. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.