

NCR3 polymorphism, haematological parameters, and severe malaria in Senegalese patients

Alassane Thiam, Sabrina Baaklini, Babacar Mbengue, Samia Nisar, Maryam Diarra, Sandrine Marquet, Mouhamadou Mansour Fall, Michel Sanka, Fatou Thiam, Rokhaya Ndiaye Diallo, et al.

► **To cite this version:**

Alassane Thiam, Sabrina Baaklini, Babacar Mbengue, Samia Nisar, Maryam Diarra, et al.. NCR3 polymorphism, haematological parameters, and severe malaria in Senegalese patients. PeerJ, PeerJ, 2018, 6, pp.e6048. 10.7717/peerj.6048 . hal-01947567

HAL Id: hal-01947567

<https://hal-amu.archives-ouvertes.fr/hal-01947567>

Submitted on 7 Dec 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





NCR3 polymorphism, haematological parameters, and severe malaria in Senegalese patients

Alassane Thiam¹, Sabrina Baaklini², Babacar Mbengue³, Samia Nisar², Maryam Diarra⁴, Sandrine Marquet², Mouhamadou Mansour Fall⁵, Michel Sanka², Fatou Thiam¹, Rokhaya Ndiaye Diallo¹, Magali Torres², Alioune Dieye^{1,3} and Pascal Rihet²

¹Unité d'Immunogénétique, Institut Pasteur de Dakar, Dakar, Senegal

²Aix Marseille Univ, INSERM, TAGC, Marseille, France

³Service d'Immunologie, University Cheikh Anta Diop of Dakar, Dakar, Senegal

⁴G4 Biostatistique, Institut Pasteur de Dakar, Dakar, Sénégal

⁵Service de Réanimation, Hôpital Principal de Dakar, Dakar, Senegal

ABSTRACT

Background. Host factors, including host genetic variation, have been shown to influence the outcome of *Plasmodium falciparum* infection. Genome-wide linkage studies have mapped mild malaria resistance genes on chromosome 6p21, whereas *NCR3-412* polymorphism ([rs2736191](#)) lying within this region was found to be associated with mild malaria.

Methods. Blood samples were taken from 188 *Plasmodium falciparum* malaria patients (76 mild malaria patients, 85 cerebral malaria patients, and 27 severe non-cerebral malaria patients). *NCR3-412* ([rs2736191](#)) was analysed by sequencing, and haematological parameters were measured. Finally, their association with clinical phenotypes was assessed.

Results. We evidenced an association of thrombocytopenia with both cerebral malaria and severe non-cerebral malaria, and of an association of high leukocyte count with cerebral malaria. Additionally, we found no association of *NCR3-412* with either cerebral malaria, severe non-cerebral malaria, or severe malaria after grouping cerebral malaria and severe non-cerebral malaria patients.

Conclusions. Our results suggest that *NCR3* genetic variation has no effect, or only a small effect on the occurrence of severe malaria, although it has been strongly associated with mild malaria. We discuss the biological meaning of these results. Besides, we confirmed the association of thrombocytopenia and high leukocyte count with severe malaria phenotypes.

Subjects Genetics, Epidemiology, Infectious Diseases

Keywords Genetic association, Mild malaria, Severe malaria, *Plasmodium falciparum*, Host factors

BACKGROUND

Plasmodium falciparum malaria remains a major public health concern causing 212 million malaria cases occurred in 2015 ([WHO, 2016](#)). The most severe forms of malaria caused by

Submitted 2 July 2018
Accepted 31 October 2018
Published 3 December 2018

Corresponding author
Pascal Rihet,
pascal.rihet@univ-amu.fr

Academic editor
Katie Ewer

Additional Information and
Declarations can be found on
page 10

DOI [10.7717/peerj.6048](https://doi.org/10.7717/peerj.6048)

© Copyright
2018 Thiam et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

P. falciparum are cerebral malaria, severe anaemia, and respiratory distress (Marsh & Snow, 1999). According to the WHO report, the major complications account for 429,000 deaths annually, essentially in children, pregnant woman and immuno-suppressed individuals. It should be noticed, however, that approximately 2% of the clinical cases are severe (Greenwood, 1997), while most of malarial infection cases remain asymptomatic.

Host genetic factors have been shown to influence the outcome of infection (Hernandez-Valladares, Rihet & Iraqi, 2014; Marquet, 2017). Genome scans have identified several human loci linked or associated with parasitaemia, mild malaria (Brisebarre et al., 2014; Milet et al., 2010; Sakuntabhai et al., 2008), or severe malaria (Band et al., 2013; Jallow et al., 2009; Malaria Genomic Epidemiology et al., 2015; Timmann et al., 2012). Genome scans that have been performed in mice have also identified some loci, with some corresponding to those genetically linked to parasitemia (Char 3 and Char 8) or cerebral malaria (Cm5c) in humans (Hernandez-Valladares, Rihet & Iraqi, 2014). In addition, many candidate gene studies provided evidence of an association between human genetic variants and malaria phenotypes (Driss et al., 2011; Hernandez-Valladares, Rihet & Iraqi, 2014; Kwiatkowski, 2005; Marquet, 2017). Some genes have been found to be associated with both mild and severe forms of clinical malaria. These include *HBB*, *TNF* and *NOS2A* (Kwiatkowski, 2005; Marquet, 2017). Interestingly, *TNF* is located within chromosome 6p21.3 linked to parasitemia and mild malaria in humans, and the corresponding chromosomal region in mice is genetically linked to mild malaria and cerebral malaria (Burt et al., 1999; Hernandez-Valladares et al., 2004; Ohno & Nishimura, 2004). Moreover, other genetic variants within chromosome 6p21.3 have been associated with severe malaria (Diakite et al., 2009).

NCR3 that is also located within chromosome 6p21.3 has been shown to be associated with mild malaria in Burkina Faso (Delahaye et al., 2007) and in the Republic of Congo (Baaklini et al., 2017). Both population-based and family-based studies showed that *NCR3-412C* carriers had more frequent mild malaria attacks than *NCR3-412GG* individuals. In addition, Baaklini et al. (2017) showed that *NCR3-412* (rs2736191) alters both the binding of transcription factors to the DNA and the promoter activity. Moreover, Mavoungou et al. (2007) reported that NKp30 encoded by *NCR3* is involved in the recognition of *P. falciparum*-infected red blood cells by natural killer (NK) cells. It might be possible that *NCR3-412* alters the transcriptional level of *NCR3* resulting in an effect on the amount of NKp30 at the NK cell surface and their response to infected red blood cells. To our knowledge, there was, however, no *NCR3* association study with severe malaria.

Here we considered *NCR3* as a candidate gene for severe malaria, and analysed its polymorphism in *P. falciparum*-infected individuals. We genotyped *NCR3-412* (rs2736191) in DNA samples from patients including 76 mild malaria, 85 cerebral malaria, and 27 severe non-cerebral malaria subjects. We report the analysis of *NCR3-412* (rs2736191) and several haematological parameters association with severe malaria phenotypes.

METHODS

Patients and phenotypes

The study was performed between 2012 and 2015 in two Senegalese hospitals including one in Dakar and the other in Tambacounda at the Southeast part of the country. Two categories

of patients with clinical malaria namely, severe malaria (SM) and mild malaria (MM) patients were enrolled. All the clinical cases were defined according to the WHO criteria (WHO, 2000). For all patients, the presence of *P. falciparum* infection was determined by an immunoassay enabling the detection of pfHRP2 (Standard diagnostics-Abbott Inc, Chicago, IL, USA). *P. falciparum* parasitemia was assessed by microscopic examination of thin and thick smears, prior anti-malarial treatment. Clinical histories, data and basic demographic information, ethnic group, and study questionnaires were recorded at baseline for each subject. In particular, the affiliation to an ethnic group was provided by the patients or their parents. At the day of admission, biological data including parasite density, haematology and biochemistry parameters were determined at the hospitals' medical laboratories and recorded. Briefly, thick and thin blood films were stained with 10% Giemsa solution for 15 min to measure parasitaemia. Parasite determination and measurement were established by two independent readings. The parasitaemia was defined as the number of parasitized erythrocytes observed per μl of blood. *P. falciparum* was the only *Plasmodium* species detected. Samples were taken at the admission before any treatment. An informed consent was obtained from each participant and/or their relatives prior to inclusion, after giving them written or verbal information in their native language. The protocols were approved by the investigator's institutions, the National Ethical Committee and the Ministry of Health of Senegal.

For severe malaria, two groups of patients were defined according to the neurological dysfunction, namely cerebral malaria cases (CM) and non CM (NCM) cases. CM cases were defined on the basis of a deep coma, an unpurposeful response or no response to a painful stimulus by Glasgow score <9 , a microscopically diagnosed *P. falciparum* infection, without other clinically cause of impaired consciousness, such as hypoglycemia, meningitis, and encephalitis according to WHO criteria (Saissy, Rouvin & Koulmann, 2003). NCM cases were defined as severe malaria (SM) without neurological symptoms such as impaired consciousness, convulsions and long-term neurological deficits. NCM patients presented isolated symptoms of SM such as severe anaemia, hypoglycaemia, respiratory distress, or hypoxia. The NCM group consisted of 27 patients, whereas the CM group consisted of 85 patients. The patients recruited in each hospital were managed by the medical staff. The treatment protocol was based on the Senegalese national recommendations, which are intramuscular quinine 20 mg/kg followed by 20 mg/kg every 8 h. Patients were examined every 4 h for the first 24 h and every 6 h thereafter.

Regarding MM, a total of 76 patients who were treated at the outpatient clinic of the hospital were initially enrolled. Of these, 76 patients had fever with *P. falciparum* parasitaemia of $<25,000$ parasites/ μL of blood, with no evidence of impaired consciousness or seizures before and at the time of enrolment. MM patients were followed up for two weeks after their enrolment, and there was no case of impaired consciousness or seizures. Blood samples from MM patients were obtained on the day of hospital admission.

DNA extraction, genome amplification, and genotyping

Blood Mini Kit (Qiagen, Hilden, Germany) was used for DNA extraction according to the manufacturer's instructions (Koukouvikila-Koussounda et al., 2013). The total DNA samples

were then amplified by IllustraGenomiPhi V2 DNA Amplification kit (GE Healthcare Life Sciences, Velizy-Villacoublay, France) using the following protocol: 1 μ l containing approximately 10 ng of genomic DNA was mixed with 9 μ l of Sample Buffer. The mixture was heated to 95°C for 3 min to denature genomic DNA and then cooled down to 4 °C or incubated on ice. For DNA amplification, a master mix solution containing 9 μ l of Reaction Buffer and 1 μ l of Phi29 DNA polymerase enzyme mix was added directly in the sample. Sample were then incubated at 30 °C for 1.5 h and heated to 65 °C for 10 min to inactivate the enzyme. The amplification product was diluted up to 1 ml in total volume with pure water to reach a DNA concentration of about 4–7 ng/ μ l and was stored at –20 °C until used.

The primer pair to amplify DNA fragments containing *NCR3-412* ([rs2736191](#)) was designed as described ([Baaklini et al., 2017](#)); PCR amplification of DNA fragments was performed with forward (5'-GATGGGTCTGGTACTGGTG-3') and reverse (5'-GGGATCTGAGCAGTGAGGTC-3') primers, respectively. For Hemoglobin S ([rs334](#)) and hemoglobin C ([rs33930165](#)), the forward primer was 5'-CTGAGGGTTGAAGTCCAAC-3', and the reverse one was 5'-CAGCATCAGGAGTGGACAG-3'. The PCR products were submitted to electrophoresis on 1.5% agarose gel electrophoresis to verify the product size and were purified using a PCR DNA Purification Kit (QIAGEN). The purified PCR products were sequenced using the Sanger method (GATC Biotech, Konstanz, Germany).

Statistical analyses

Hardy-Weinberg equilibrium was tested and allelic frequencies were calculated, as described ([Rodriguez, Gaunt & Day, 2009](#)). Other statistical tests were performed by using R or the SPSS software (SPSS, Boulogne, France). To assess the association of quantitative biological phenotypes and clinical phenotypes, we used the Kruskal-Wallis test and the Mann-whitney test. To estimate the association of [rs2736191](#) polymorphism with malaria phenotypes, we conducted a chi-square test and a logistic regression analysis, and we evaluated the odds ratios (OR) and their 95% confidence intervals (CIs). MM group was considered the control group, whereas the case group consisted of CM patients, NCM patients, or severe malaria patients after grouping CM and NCM patients. The distribution of MM patient genotypes was compared with the one of CM, NCM, or severe malaria patient genotypes. All the tests were two-sided. G*power software was used to compute statistical power for the genetic association study.

RESULTS

Association between biological parameters and severe malaria

[Table 1](#) shows the characteristics of the patients. There was no effect of gender (Chi-square = 2.4, $df = 2$, $P = 0.3$). CM patients had a significantly higher risk of dying compared to NCM ($P = 0.04$) or MM patients ($P = 2.0 \cdot 10^{-6}$) by using a Fisher exact test. Moreover, there were significant differences between CM, NCM, and MM patients on the basis of the Kruskal-Wallis test for age ($P = 0.009$), erythrocyte count ($P = 0.019$), haematocrit ($P = 3.3 \cdot 10^{-6}$), haemoglobin concentration ($P = 2.4 \cdot 10^{-6}$), leucocyte count ($P = 0.006$), and platelet count ($P = 7.9 \cdot 10^{-4}$).

Table 1 Patient characteristics. Characteristics include age, gender, ethnicity, haematological parameters, and beta-globin genetic variation.

	Cerebral malaria (N = 85)	Non cerebral malaria (N = 27)	Mild malaria (N = 76)
Gender			
Female	28	10	34
Male	57	17	42
Survival			
Dead	19	1	0
Alive	66	26	76
Ethnic group^a			
Bambara	8	3	5
Diola	5	3	2
Peulh	22	14	30
Serrere	8	0	9
Soninke	3	1	8
Wolof	28	4	20
Others	11	2	2
Age^b			
Median	22	8	27
(25th and 75th percentile)	(12–41)	(3–33)	(13–41)
Haematological data^b			
<i>-Red blood cells</i> ($\times 10^6/\mu\text{L}$)			
N	68	23	66
Median	3.8	2.8	4.0
(25th and 75th percentile)	(3.0–4.3)	(2.4–4.1)	(3.5–4.6)
<i>-Haematocrit</i> (%)			
N	67	23	66
Median	30.6	20.0	35.0
(25th and 75th percentile)	(25.8–34.4)	(16.9–27.5)	(29.7–38.7)
<i>-Haemoglobin</i> (g/dL)			
N	80	26	66
Median	10.4	7.1	11.1
(25th and 75th percentile)	(8.4–11.9)	(5.3–9.4)	(9.8–12.5)
<i>-Leucocytes</i> ($\times 10^3/\mu\text{L}$)			
N	77	24	66
Median (25th and 75th percentile)	11.0 (6.5–14.0)	7.9 (5.4–12.9)	7.9 (5.0–10.6)
<i>-Platelets</i> ($\times 10^3/\mu\text{L}$)			
N	66	26	66
Median	87.5	100.5	183.0
(25th and 75th percentile)	(40.0–168.3)	(58.5–165.0)	(83.0–237.8)
<i>-Parasitemia</i> (pRBC ^c / μL)			
N	37	16	51
Median	560	2,800	6,545
(25th and 75th percentile)	(115–5,983)	(592–6,716)	(620–57,840)

(continued on next page)

Table 1 (continued)

	Cerebral malaria (N = 85)	Non cerebral malaria (N = 27)	Mild malaria (N = 76)
HBB genetic variation^d			
-Hb S carrier, n (%)	3 (3.6)	1 (3.7)	5 (6.7)
-Hb C carrier, n (%)	0 (0.0)	2 (7.4)	1 (1.3)

Notes.

^aEthnic groups are shown. Other ethnic groups that are Bassari, Cap-vert, Mandinge, and Soce groups were rare ($n \leq 5$). The ethnic group was unknown for two individuals.

^bMedian and (25th and 75th percentile) are shown.

^cParasitized red blood cells.

^dTwo individuals have not been genotyped.

NCM patients were younger than CM and MM patients ($P < 0.006$) by using the Mann–Whitney method, whereas there was no difference between CM and MM for age. Erythrocyte count ($P = 0.01$), haematocrit ($P = 4.9 \times 10^{-6}$), and haemoglobin concentration ($P = 1.4 \times 10^{-6}$) of NCM patients were lower than those of MM patients. NCM patients showed a lower haematocrit ($P = 1.1 \times 10^{-4}$) and a lower haemoglobin concentration ($P = 3.4 \times 10^{-5}$) compared to CM patients. Moreover, we detected a reduction of platelet count ($P = 0.0003$) and an increase of leucocyte count ($P = 0.0015$) in CM patients compared to the ones of MM patients, as shown in Fig. 1. There was no significant difference between CM and NCM for leucocyte count ($P = 0.14$) and for platelet count ($P = 0.64$). Furthermore, we found a reduction of platelet count in NCM patients compared to the ones of MM patients ($P = 0.018$), whereas there was no difference for the leucocyte count ($P = 0.44$) between these two groups.

Influence of ethnic groups and genetic factors on severe malaria

Table 1 shows the distribution of the ethnic groups according to the clinical phenotypes. Peul and Wolof groups were the most represented ethnic groups among the patients. These groups differed in the distribution of their clinical phenotype (Chi-square = 6.4, $df = 2$, $P = 0.04$), whereas there was a trend when comparing Peul to all other ethnic groups (Chi-square = 5.6, $df = 2$, $P = 0.06$). More specifically, there was a trend in favour of a better protection from CM (OR = 0.576, 95% confidence interval 0.296–1.120), although not significant ($P > 0.05$), on the basis of the comparison of MM with CM patients. In the same way, there was a trend in favour of a lower parasitaemia mean in Peul patients, although not significant ($P > 0.05$); there was, nevertheless, a lower parasitaemia variance in Peul patients ($P = 0.003$). Haemoglobin S and haemoglobin C carriers were rare in CM, NCM, and MM patients, as shown in Table 1. There was no significant association between haemoglobin genotypes and clinical phenotypes.

Table 2 shows the distribution of *NCR3* genotype by clinical status. There were 123 GG, 52 GC, and 8 CC individuals. *NCR3-412* was in Hardy-Weinberg equilibrium ($P = 0.77$). *NCR3-412G* allele frequency was 0.81, whereas *NCR3-412C* allele frequency was 0.19. We clustered GC and CC individuals for further analyses.

We tested the association of the *NCR3-412C* allele carriage with severe malaria by using the Chi-square test. To this aim, we compared the distribution of MM patient genotypes with the one of severe malaria patient genotypes. There was no association with CM (Chi-square = 0.9, $df = 1$, $P = 0.34$), NCM (Chi-square = 0.09, $df = 1$, $P = 0.77$), or severe malaria after grouping CM and NCM patients (Chi-square = 0.77, $df = 1$, $P = 0.38$).




Figure 1 Haematological characteristics of mild malaria (MM), cerebral malaria (CM), and noncerebral malaria (NCM) patients. (A) Platelet count ($\times 10^3/\mu\text{L}$). (B) Leucocyte count ($\times 10^3/\mu\text{L}$). Median, 25th and 75th percentile are shown for each group. Significant differences between MM and CM, and between MM and NCM are shown. P values have been calculated by using Mann-Whitney test (*** $P < 0.001$; ** $P < 0.01$; * $P < 0.05$).

Full-size [DOI: 10.7717/peerj.6048/fig-1](https://doi.org/10.7717/peerj.6048/fig-1)

Table 2 Clinical status of patients for *NCR3-412* polymorphism.

Genotype	Cerebral malaria	Non cerebral malaria	Mild malaria	Total
CC	5	0	3	8
GC	19	9	24	52
GG	58	18	47	123
Total	82 ^a	27	74 ^a	183

Notes.

^aThree CM and 2 MM individuals have not been genotyped.

We applied a logistic regression method when taking into account age and gender, which may influence malaria phenotypes. There was no association with CM ($P = 0.56$), NCM ($P = 0.92$), or severe malaria ($P = 0.61$). The odds of CM, NCM, and severe malaria between *NCR3-412GG* and *NCR3-412GG/GC* were 0.813 (95% confidence interval 0.406–1.628), 1.047 (95% confidence interval 0.396–2.769), and 0.847 (95% confidence interval 0.445–1.614), respectively. Similar results were obtained after adding haemoglobin genotypes and ethnic groups as covariates. In addition, we assessed the power of our analyses on the basis of the sample size used, an α -risk of 5%, and an effect size equal to the one detected by *Delahaye et al. (2007)* for the association of the *NCR3-412C* allele carriage with mild malaria. We obtained a power of 85% and 90% for CM and severe malaria, respectively.

DISCUSSION

A few genes have been associated with both mild malaria and severe malaria. We investigated a *NCR3* genetic variant, which was linked and associated with mild malaria in Burkina Faso (*Delahaye et al., 2007*). We recruited a new study population in Senegal, recorded biological parameters and ethnic groups, genotyped DNA samples for *NCR3-412*, haemoglobin C

and haemoglobin S polymorphism in patients with either mild or severe malaria, and performed an association study based on a case-control design.

Reduced red blood cell count, haematocrit, and haemoglobin concentration were associated with noncerebral malaria, indicating a marked anaemia in these patients compared to mild malaria and cerebral malaria. Thrombocytopenia was associated with both cerebral malaria and severe non-cerebral malaria, as previously reported (*Gerardin et al., 2002; Lampah et al., 2015; Muwonge et al., 2013*). Thrombocytopenia is thought to be due to platelet sequestration and platelet consumption (*Akinosoglou, Solomou & Gogos, 2012; Hochman et al., 2015; Karanikas et al., 2004*). In the same way, platelet adhesion to brain endothelial cells has been associated with cerebral malaria (*Grau et al., 2003*). Furthermore, it has been shown to promote adhesion of infected red blood cells to TNF-stimulated endothelium (*Wassmer et al., 2004*), and to alter gene expression *in vitro* in human brain microvascular endothelial cells (*Barbier et al., 2011*), thus contributing to brain endothelial dysfunction.

Moreover, an increase of leukocyte count was associated with CM, whereas MM patients and NCM patients did not differ in their leukocyte count. In the same way, leukocyte counts were reported to be higher in severe malaria patients compared to those of mild malaria patients (*Berens-Riha et al., 2014; Cserti-Gazdewich et al., 2013; Modiano et al., 2001c; Van Wolfswinkel et al., 2013*), and in malaria patients with high parasitaemia compared to those of individuals with low parasitaemia (*Kotepui et al., 2015; McKenzie et al., 2005*). The increase of leukocyte number is consistent with a strong host inflammatory response associated with severe malaria. It may be due to polymorphonuclear leukocyte or monocyte increase (*Hochman et al., 2015*). T, B, and NK lymphocyte counts have been found to be reduced in severe malaria patients (*Mandala et al., 2015*), whereas the percentage of activated T CD4+, NK, and $\gamma\delta$ T lymphocytes were higher in cerebral malaria patients compared to mild malaria patients (*Mandala et al., 2015*). This is consistent with the role of T cell dependent immunopathology in the pathogenesis of cerebral malaria. This also suggests a role of NK and $\gamma\delta$ T lymphocytes in malaria immunopathology. In the same way, NK and $\gamma\delta$ T lymphocytes produce IFN γ and TNF α in malaria patients (*D'Ombra et al., 2007; Horowitz et al., 2010; Stanisic et al., 2014*), while polymorphisms within the natural killer cell complex modulate mouse cerebral malaria (*Hansen et al., 2014*). In humans, the effect of NK polymorphisms on severe malaria has been poorly investigated, and yielded conflicting results (*Olaniyan et al., 2014; Yindom et al., 2012*).

Since the Peul ethnic group which has a particular genetic background has a lower risk of mild malaria attacks, a lower parasitaemia, and a higher anti-plasmodial antibody response compared to other sympatric ethnic groups (*Arama et al., 2015; Modiano et al., 2001a; Modiano et al., 1996*), we hypothesized that Peul patients may also show a better protection against severe malaria. Although we detected a trend in favour of this hypothesis, the association was not significant at the 5% level. This result urges, nevertheless, new studies with a bigger sample size to increase statistical power.

Since, haemoglobin S and haemoglobin C are well known to protect against severe malaria (*Jallow et al., 2009; Modiano et al., 2001b; Timmann et al., 2012*), we searched for haemoglobin S and haemoglobin C in our study population. Although we detected both




Figure 2 *NCR3* and severity of the disease. *NCR3-412C* allele that increased the risk of mild malaria was shown to decrease both the binding of transcription factors to the promoter and *NCR3* expression. Here we report that there was no difference between *NCR3-412* allele frequency between mild malaria patients and severe malaria patients. Therefore, *NCR3-412* polymorphism unlikely explains why some patients with mild malaria develop severe forms of the disease.

Full-size DOI: [10.7717/peerj.6048/fig-2](https://doi.org/10.7717/peerj.6048/fig-2)

mutations, the frequency of haemoglobin S and haemoglobin C was very low, and we could not provide evidence for a protective effect against severe malaria.

We further assessed the association of *NCR3-412* with severe malaria. Mild malaria patients and severe malaria patients did not differ in their *NCR3-412* allelic frequency. Power studies revealed a power of approximately 85% based on an effect size corresponding to the one found in the mild malaria studies (Delahaye et al., 2007). This indicates that the effect size is significantly lower for severe malaria or that there is no association between the SNP and severe malaria. In the former case, increasing the sample size should allow to detect an association with a small effect size. Alternatively, integrating the polymorphisms of other NK cell receptors should increase the power for detecting a putative association of NK receptors with severe malaria. In the later case, the *NCR3* expression may not influence the occurrence of severe malaria. It should be stressed that very few genes have been associated both with mild and severe malaria.

If we assume that there is no association between *NCR3-412* and severe malaria, we should keep in mind that there is an association between *NCR3-412* and mild malaria (Baaklini et al., 2017; Delahaye et al., 2007). This means that there is no allele frequency difference between mild malaria patients and severe malaria patients, on the one hand, and that there is an allele frequency difference between mild malaria patients and individuals with an asymptomatic infection, on the other hand. Also, *NCR3-412* polymorphism may partly explain why individuals with asymptomatic malaria have mild malaria attacks, but is unlikely to explain why some patients with mild malaria develop severe forms of the disease (Fig. 2).

It is important to point out the limitation of our study. First, the time of our clinical investigation was limited. Also, the absence of CM or NCM was registered on the basis of the declaration of the patients or their parents. Second, it cannot be excluded that clinical status could be due to differential anti-malarial immunity response resulting from different exposure levels. Finally, the sample size of our study population is limited. In spite of this limitation, it should be stressed that we detected an association of leucocyte and platelet counts with clinical status, confirming previous results. This shows that the sample size was sufficient to allow us to detect association with a high effect size. In this way, our power

analysis for genetic association indicated that we could detect an association of *NCR3* with severe malaria when assuming an effect size similar to the one we calculated for mild malaria in Burkina Faso or in the Republic of Congo ([Baaklini et al., 2017](#); [Delahaye et al., 2007](#)). Nevertheless, we cannot exclude that there was an association of *NCR3* with severe malaria with a small effect size.

CONCLUSIONS

To summarize, we conducted an epidemiological study that allowed us to confirm the association of cerebral malaria with thrombocytopenia and leukocyte count. Besides, we did not detect an association of severe malaria with *NCR3* genetic variation. This suggests that genetic variation at this locus does not influence the molecular mechanisms involved in the most severe forms of the disease, although it influences the risk of developing mild malaria attacks.

Abbreviations

CI	confidence interval
CM	cerebral malaria
NCM	non-cerebral malaria
MM	mild malaria
NK	natural killer
OR	Odds ratio

ACKNOWLEDGEMENTS

We thank all the patients who participated in this study.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by the management of the Institute Pasteur of Dakar, the Directorate of International Affairs of the Institute Pasteur in Paris, the Department of Cultural Action of the French Embassy in Senegal, INSERM, and the university of Aix-Marseille. S Baaklini, M Sanka, and S Nisar were supported by a PhD fellowship from the French ministry, PACA regional council, and the Higher Education Commission (HEC) from Pakistan, respectively. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

Institute Pasteur of Dakar.

Directorate of International Affairs of the Institute Pasteur in Paris.

Department of Cultural Action of the French Embassy in Senegal.

INSERM.

University of Aix-Marseille.
French ministry.
PACA regional council.
Higher Education Commission (HEC) from Pakistan.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Alassane Thiam performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Sabrina Baaklini performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Babacar Mbengue conceived and designed the experiments, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Samia Nisar, Sandrine Marquet and Magali Torres performed the experiments, approved the final draft.
- Maryam Diarra analyzed the data, approved the final draft.
- Mouhamadou Mansour Fall contributed reagents/materials/analysis tools, approved the final draft, he was in charge of clinical management of patients, and measurements of clinical and biological parameters.
- Michel Sanka performed the experiments, prepared figures and/or tables, approved the final draft.
- Fatou Thiam contributed reagents/materials/analysis tools, approved the final draft.
- Rokhaya Ndiaye Diallo contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Alioune Dieye conceived and designed the experiments, authored or reviewed drafts of the paper, approved the final draft.
- Pascal Rihet conceived and designed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.

Human Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

The National Ethical Committee and the Ministry of Health of Senegal granted approval to carry out the study.

Data Availability

The following information was supplied regarding data availability:

The raw data are provided in [Supplemental File](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.6048#supplemental-information>.

REFERENCES

- Akinosoglou KS, Solomou EE, Gogos CA. 2012. Malaria: a haematological disease. *Hematology* 17:106–114 DOI [10.1179/102453312X13221316477336](https://doi.org/10.1179/102453312X13221316477336).
- Arama C, Maiga B, Dolo A, Kouriba B, Traore B, Crompton PD, Pierce SK, Troye-Blomberg M, Miller LH, Doumbo OK. 2015. Ethnic differences in susceptibility to malaria: what have we learned from immuno-epidemiological studies in West Africa? *Acta Tropica* 146:152–156 DOI [10.1016/j.actatropica.2015.03.023](https://doi.org/10.1016/j.actatropica.2015.03.023).
- Baaklini S, Afridi S, Nguyen TN, Koukouikila-Koussounda F, Ndounga M, Imbert J, Torres M, Pradel L, Ntoumi F, Rihet P. 2017. Beyond genome-wide scan: association of a cis-regulatory NCR3 variant with mild malaria in a population living in the Republic of Congo. *PLOS ONE* 12:e0187818 DOI [10.1371/journal.pone.0187818](https://doi.org/10.1371/journal.pone.0187818).
- Band G, Le QS, Jostins L, Pirinen M, Kivinen K, Jallow M, Sisay-Joof F, Bojang K, Pinder M, Sirugo G, Conway DJ, Nyirongo V, Kachala D, Molyneux M, Taylor T, Ndila C, Peshu N, Marsh K, Williams TN, Alcock D, Andrews R, Edkins S, Gray E, Hubbart C, Jeffreys A, Rowlands K, Schuldt K, Clark TG, Small KS, Teo YY, Kwiatkowski DP, Rockett KA, Barrett JC, Spencer CC, Malaria Genomic Epidemiological Network. 2013. Imputation-based meta-analysis of severe malaria in three African populations. *PLOS Genetics* 9:e1003509 DOI [10.1371/journal.pgen.1003509](https://doi.org/10.1371/journal.pgen.1003509).
- Barbier M, Faille D, Loriod B, Textoris J, Camus C, Puthier D, Flori L, Wassmer SC, Victorero G, Alessi MC, Fusai T, Nguyen C, Grau GE, Rihet P. 2011. Platelets alter gene expression profile in human brain endothelial cells in an *in vitro* model of cerebral malaria. *PLOS ONE* 6:e19651 DOI [10.1371/journal.pone.0019651](https://doi.org/10.1371/journal.pone.0019651).
- Berens-Riha N, Kroidl I, Schunk M, Alberer M, Beissner M, Pritsch M, Kroidl A, Froschl G, Hanus I, Bretzel G, Von Sonnenburg F, Nothdurft HD, Loscher T, Herbinger KH. 2014. Evidence for significant influence of host immunity on changes in differential blood count during malaria. *Malaria Journal* 13:155–164 DOI [10.1186/1475-2875-13-155](https://doi.org/10.1186/1475-2875-13-155).
- Brisebarre A, Kumulungui B, Sawadogo S, Atkinson A, Garnier S, Fumoux F, Rihet P. 2014. A genome scan for *Plasmodium falciparum* malaria identifies quantitative trait loci on chromosomes 5q31, 6p21.3, 17p12, and 19p13. *Malaria Journal* 13:198–205 DOI [10.1186/1475-2875-13-198](https://doi.org/10.1186/1475-2875-13-198).
- Burt RA, Baldwin TM, Marshall VM, Foote SJ. 1999. Temporal expression of an H2-linked locus in host response to mouse malaria. *Immunogenetics* 50:278–285 DOI [10.1007/s002510050603](https://doi.org/10.1007/s002510050603).

- Cserti-Gazdewich CM, Dhabangi A, Musoke C, Ssewanyana I, Ddungu H, Nakiboneka-Ssenabulya D, Nabukeera-Barungi N, Mpimbaza A, Dzik WH. 2013. Interrelationships of cardinal features and outcomes of symptomatic pediatric *Plasmodium falciparum* MALARIA in 1, 933 children in Kampala, Uganda. *American Journal of Tropical Medicine and Hygiene* 88:747–756 DOI 10.4269/ajtmh.12-0668.
- Delahaye NF, Barbier M, Fumoux F, Rihet P. 2007. Association analyses of NCR3 polymorphisms with *P. falciparum* mild malaria. *Microbes and Infection* 9:160–166 DOI 10.1016/j.micinf.2006.11.002.
- Diakite M, Clark TG, Auburn S, Campino S, Fry AE, Green A, Morris AP, Richardson A, Jallow M, Sisay-Joof F, Pinder M, Kwiatkowski DP, Rockett KA. 2009. A genetic association study in the Gambia using tagging polymorphisms in the major histocompatibility complex class III region implicates a HLA-B associated transcript 2 polymorphism in severe malaria susceptibility. *Human Genetics* 125:105–109 DOI 10.1007/s00439-008-0597-2.
- D’Ombrain MC, Hansen DS, Simpson KM, Schofield L. 2007. gammadelta-T cells expressing NK receptors predominate over NK cells and conventional T cells in the innate IFN-gamma response to *Plasmodium falciparum* malaria. *European Journal of Immunology* 37:1864–1873 DOI 10.1002/eji.200636889.
- Driss A, Hibbert JM, Wilson NO, Iqbal SA, Adamkiewicz TV, Stiles JK. 2011. Genetic polymorphisms linked to susceptibility to malaria. *Malaria Journal* 10:271–281 DOI 10.1186/1475-2875-10-271.
- Gerardin P, Rogier C, Ka AS, Jouvencel P, Brousse V, Imbert P. 2002. Prognostic value of thrombocytopenia in African children with falciparum malaria. *American Journal of Tropical Medicine and Hygiene* 66:686–691 DOI 10.4269/ajtmh.2002.66.686.
- Grau GE, Mackenzie CD, Carr RA, Redard M, Pizzolato G, Allasia C, Cataldo C, Taylor TE, Molyneux ME. 2003. Platelet accumulation in brain microvessels in fatal pediatric cerebral malaria. *Journal of Infectious Diseases* 187:461–466 DOI 10.1086/367960.
- Greenwood BM. 1997. The epidemiology of malaria. *Annals of Tropical Medicine and Parasitology* 91:763–769 DOI 10.1080/00034983.1997.11813201.
- Hansen DS, Ryg-Cornejo V, Ioannidis LJ, Chiu CY, Ly A, Nie CQ, Scalzo AA, Schofield L. 2014. The contribution of natural killer complex loci to the development of experimental cerebral malaria. *PLOS ONE* 9:e93268 DOI 10.1371/journal.pone.0093268.
- Hernandez-Valladares M, Naessens J, Gibson JP, Musoke AJ, Nagda S, Rihet P, Ole-MoiYoi OK, Iraqi FA. 2004. Confirmation and dissection of QTL controlling resistance to malaria in mice. *Mammalian Genome* 15:390–398 DOI 10.1007/s00335-004-3042-4.
- Hernandez-Valladares M, Rihet P, Iraqi FA. 2014. Host susceptibility to malaria in human and mice: compatible approaches to identify potential resistant genes. *Physiological Genomics* 46:1–16 DOI 10.1152/physiolgenomics.00044.2013.
- Hochman SE, Madaline TF, Wassmer SC, Mbale E, Choi N, Seydel KB, Whitten RO, Varughese J, Grau GE, Kamiza S, Molyneux ME, Taylor TE, Lee S, Milner Jr DA, Kim K. 2015. Fatal pediatric cerebral malaria is associated with intravascular

monocytes and platelets that are increased with HIV coinfection. *MBio* 6:e01390–e01315 DOI 10.1128/mBio.01390-15.

- Horowitz A, Newman KC, Evans JH, Korb DS, Davis DM, Riley EM. 2010.** Cross-talk between T cells and NK cells generates rapid effector responses to *Plasmodium falciparum*-infected erythrocytes. *Journal of Immunology* 184:6043–6052 DOI 10.4049/jimmunol.1000106.
- Jallow M, Teo YY, Small KS, Rockett KA, Deloukas P, Clark TG, Kivinen K, Bojang KA, Conway DJ, Pinder M, Sirugo G, Sisay-Joof F, Usen S, Auburn S, Bumpstead SJ, Campino S, Coffey A, Dunham A, Fry AE, Green A, Gwilliam R, Hunt SE, Inouye M, Jeffreys AE, Mendy A, Palotie A, Potter S, Ragoussis J, Rogers J, Rowlands K, Somaskantharajah E, Whittaker P, Widdon C, Donnelly P, Howie B, Marchini J, Morris A, SanJoaquin M, Achidi EA, Agbenyega T, Allen A, Amodu O, Corran P, Djimde A, Dolo A, Doumbo OK, Drakeley C, Dunstan S, Evans J, Farrar J, Fernando D, Hien TT, Horstmann RD, Ibrahim M, Karunaweera N, Kokwaro G, Koram KA, Lemnge M, Makani J, Marsh K, Michon P, Modiano D, Molyneux ME, Mueller I, Parker M, Peshu N, Plowe CV, Puijalon O, Reeder J, Reyburn H, Riley EM, Sakuntabhai A, Singhasivanon P, Sirima S, Tall A, Taylor TE, Thera M, Troye-Blomberg M, Williams TN, Wilson M, Kwiatkowski DP, Wellcome Trust Case Control Consortium, Malaria Genomic Epidemiology Network. 2009.** Genome-wide and fine-resolution association analysis of malaria in West Africa. *Nature Genetics* 41:657–665 DOI 10.1038/ng.388.
- Karanikas G, Zedwitz-Liebenstein K, Eidherr H, Schuetz M, Sauerman R, Dudczak R, Winkler S, Pabinger I, Kletter K. 2004.** Platelet kinetics and scintigraphic imaging in thrombocytopenic malaria patients. *Thrombosis and Haemostasis* 91:553–557 DOI 10.1160/TH03-07-0489.
- Kotepui M, Piwkhram D, PhunPhuech B, Phiwklam N, Chupeerach C, Duangmano S. 2015.** Effects of malaria parasite density on blood cell parameters. *PLOS ONE* 10:e0121057 DOI 10.1371/journal.pone.0121057.
- Koukouikila-Koussounda F, Ntoui F, Ndounga M, Tong HV, Abena AA, Velavan TP. 2013.** Genetic evidence of regulatory gene variants of the STAT6, IL10R and FOXP3 locus as a susceptibility factor in uncomplicated malaria and parasitaemia in Congolese children. *Malaria Journal* 12:9–18 DOI 10.1186/1475-2875-12-9.
- Kwiatkowski DP. 2005.** How malaria has affected the human genome and what human genetics can teach us about malaria. *American Journal of Human Genetics* 77:171–192 DOI 10.1086/432519.
- Lampah DA, Yeo TW, Malloy M, Kenangalem E, Douglas NM, Ronaldo D, Sugiarto P, Simpson JA, Poespoprodjo JR, Anstey NM, Price RN. 2015.** Severe malarial thrombocytopenia: a risk factor for mortality in Papua, Indonesia. *Journal of Infectious Diseases* 211:623–634 DOI 10.1093/infdis/jiu487.
- Malaria Genomic Epidemiology, N, Band G, Rockett KA, Spencer CC, Kwiatkowski DP. 2015.** A novel locus of resistance to severe malaria in a region of ancient balancing selection. *Nature* 526:253–257 DOI 10.1038/nature15390.

- Mandala WL, Msefula CL, Gondwe EN, Gilchrist JJ, Graham SM, Pensulo P, Mwimaniwa G, Banda M, Taylor TE, Molyneux EE, Drayson MT, Ward SA, Molyneux ME, MacLennan CA. 2015. Lymphocyte perturbations in Malawian children with severe and uncomplicated malaria. *Clinical and Vaccine Immunology* 23:95–103 DOI 10.1128/CVI.00564-15.
- Marquet S. 2017. Overview of human genetic susceptibility to malaria: from parasitemia control to severe disease. *Infection, Genetics and Evolution* Epub ahead of print Jun 1 2017 DOI 10.1016/j.meegid.2017.06.001.
- Marsh K, Snow RW. 1999. Malaria transmission and morbidity. *Parassitologia* 41:241–246.
- Mavougou E, Held J, Mewono L, Kremsner PG. 2007. A Duffy binding-like domain is involved in the NKp30-mediated recognition of *Plasmodium falciparum*-parasitized erythrocytes by natural killer cells. *Journal of Infectious Diseases* 195:1521–1531 DOI 10.1086/515579.
- McKenzie FE, Prudhomme WA, Magill AJ, Forney JR, Permpnich B, Lucas C, Gasser Jr RA, Wongsrichanalai C. 2005. White blood cell counts and malaria. *Journal of Infectious Diseases* 192:323–330 DOI 10.1086/431152.
- Milet J, Nuel G, Watier L, Courtin D, Slaoui Y, Senghor P, Migot-Nabias F, Gaye O, Garcia A. 2010. Genome wide linkage study, using a 250K SNP map, of *Plasmodium falciparum* infection and mild malaria attack in a Senegalese population. *PLOS ONE* 5:e11616 DOI 10.1371/journal.pone.0011616.
- Modiano D, Luoni G, Sirima BS, Lanfrancotti A, Petrarca V, Cruciani F, Simpoire J, Ciminelli BM, Foglietta E, Grisanti P, Bianco I, Modiano G, Coluzzi M. 2001a. The lower susceptibility to *Plasmodium falciparum* malaria of Fulani of Burkina Faso (west Africa) is associated with low frequencies of classic malaria-resistance genes. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 95:149–152 DOI 10.1016/S0035-9203(01)90141-5.
- Modiano D, Luoni G, Sirima BS, Simpoire J, Verra F, Konate A, Rastrelli E, Olivieri A, Calissano C, Paganotti GM, D'Urbano L, Sanou I, Sawadogo A, Modiano G, Coluzzi M. 2001b. Haemoglobin C protects against clinical *Plasmodium falciparum* malaria. *Nature* 414:305–308 DOI 10.1038/35104556.
- Modiano D, Petrarca V, Sirima BS, Nebie I, Diallo D, Esposito F, Coluzzi M. 1996. Different response to *Plasmodium falciparum* malaria in west African sympatric ethnic groups. *Proceedings of the National Academy of Sciences of the United States of America* 93:13206–13211 DOI 10.1073/pnas.93.23.13206.
- Modiano D, Sirima BS, Konate A, Sanou I, Sawadogo A. 2001c. Leucocytosis in severe malaria. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 95:175–176 DOI 10.1016/S0035-9203(01)90152-X.
- Muwonge H, Kikomeko S, Sembajjwe LF, Seguya A, Namugwanya C. 2013. How reliable are hematological parameters in predicting uncomplicated *Plasmodium falciparum* malaria in an endemic region? *ISRN Tropical Medicine* 2013:1–9 DOI 10.1155/2013/673798.

- Ohno T, Nishimura M. 2004.** Detection of a new cerebral malaria susceptibility locus, using CBA mice. *Immunogenetics* **56**:675–678 DOI [10.1007/s00251-004-0739-1](https://doi.org/10.1007/s00251-004-0739-1).
- Olaniyan SA, Amodu OK, Yindom LM, Conway DJ, Aka P, Bakare AA, Omotade OO. 2014.** Killer-cell immunoglobulin-like receptors and falciparum malaria in southwest Nigeria. *Human Immunology* **75**:816–821 DOI [10.1016/j.humimm.2014.06.002](https://doi.org/10.1016/j.humimm.2014.06.002).
- Rodriguez S, Gaunt TR, Day IN. 2009.** Hardy-Weinberg equilibrium testing of biological ascertainment for Mendelian randomization studies. *American Journal of Epidemiology* **169**:505–514 DOI [10.1093/aje/kwn359](https://doi.org/10.1093/aje/kwn359).
- Saissy JM, Rouvin B, Koulmann P. 2003.** Severe malaria in intensive care units in 2003. *Médecine tropicale: revue du Corps de santé colonial* **63**:258–266.
- Sakuntabhai A, Ndiaye R, Casademont I, Peerapittayamongkol C, Rogier C, Tortevoye P, Tall A, Paul R, Turbpaiboon C, Phimpraphi W, Trape JF, Spiegel A, Heath S, Mercereau-Puijalon O, Dieye A, Julier C. 2008.** Genetic determination and linkage mapping of *Plasmodium falciparum* malaria related traits in Senegal. *PLOS ONE* **3**:e2000 DOI [10.1371/journal.pone.0002000](https://doi.org/10.1371/journal.pone.0002000).
- Stanisic DI, Cutts J, Eriksson E, Fowkes FJ, Rosanas-Urgell A, Siba P, Laman M, Davis TM, Manning L, Mueller I, Schofield L. 2014.** gammadelta T cells and CD14+ monocytes are predominant cellular sources of cytokines and chemokines associated with severe malaria. *Journal of Infectious Diseases* **210**:295–305 DOI [10.1093/infdis/jiu083](https://doi.org/10.1093/infdis/jiu083).
- Timmann C, Thye T, Vens M, Evans J, May J, Ehmen C, Sievertsen J, Muntau B, Ruge G, Loag W, Ansong D, Antwi S, Asafo-Adjei E, Nguah SB, Kwakye KO, Akoto AO, Sylverken J, Brendel M, Schuldt K, Loley C, Franke A, Meyer CG, Agbenyega T, Ziegler A, Horstmann RD. 2012.** Genome-wide association study indicates two novel resistance loci for severe malaria. *Nature* **489**:443–446 DOI [10.1038/nature11334](https://doi.org/10.1038/nature11334).
- Van Wolfswinkel ME, Vliegthart-Jongbloed K, De Mendonca Melo M, Wever PC, McCall MB, Koelewijn R, Van Hellemond JJ, Van Genderen PJ. 2013.** Predictive value of lymphocytopenia and the neutrophil-lymphocyte count ratio for severe imported malaria. *Malaria Journal* **12**:101–109 DOI [10.1186/1475-2875-12-101](https://doi.org/10.1186/1475-2875-12-101).
- Wassmer SC, Lepolard C, Traore B, Pouvelle B, Gysin J, Grau GE. 2004.** Platelets reorient *Plasmodium falciparum*-infected erythrocyte cytoadhesion to activated endothelial cells. *Journal of Infectious Diseases* **189**:180–189 DOI [10.1086/380761](https://doi.org/10.1086/380761).
- WHO. 2000.** Severe falciparum malaria. World Health Organization, Communicable Diseases Cluster. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **94**(Suppl 1):S1–S90.
- WHO. 2016.** World malaria report 2016. World Health Organization, Geneva.
- Yindom LM, Forbes R, Aka P, Janha O, Jeffries D, Jallow M, Conway DJ, Walther M. 2012.** Killer-cell immunoglobulin-like receptors and malaria caused by *Plasmodium falciparum* in The Gambia. *Tissue Antigens* **79**:104–113 DOI [10.1111/j.1399-0039.2011.01818.x](https://doi.org/10.1111/j.1399-0039.2011.01818.x).