Mitochondrial diversity and phylogeographic analysis of *Pediculus humanus* reveals a new Amazonian clade “F”

Nadia Amanzougaghene, Florence Fenollar, Bernard Davoust, Félix Djossou, Muhammad Ashfaq, Idir Bitam, Didier Raoult, Oleg Mediannikov

► To cite this version:

Nadia Amanzougaghene, Florence Fenollar, Bernard Davoust, Félix Djossou, Muhammad Ashfaq, et al.. Mitochondrial diversity and phylogeographic analysis of *Pediculus humanus* reveals a new Amazonian clade “F”. *Infection, Genetics and Evolution*, 2019, 70, pp.1-8. 10.1016/j.meegid.2019.02.006 . hal-02243787

HAL Id: hal-02243787

<https://hal-amu.archives-ouvertes.fr/hal-02243787>

Submitted on 22 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial| 4.0 International License

**Mitochondrial diversity and phylogeographic analysis of *Pediculus*
humanus reveals a new Amazonian Clade “F”**

Nadia Amanzougaghene¹, Florence Fenollar², Bernard Davoust¹, Félix Djossou³,
Muhammad Ashfaq⁴, Idir Bitam^{2,5}, Didier Raoult^{1*}, Oleg Mediannikov^{1*}

¹Aix Marseille Univ, IRD, APHM, MEPHI, IHU-Méditerranée Infection, Marseille, France

²Aix Marseille Univ, IRD, APHM, SSA, VITROME, IHU-Méditerranée Infection, Marseille,
France

³Service de maladies Infectieuses et Tropicales, Centre Hospitalier de Cayenne, 97306
Cayenne Cedex, Guyane Française, France

⁴Centre for Biodiversity Genomics, Biodiversity Institute of Ontario, University of Guelph,
Guelph, ON, Canada

⁵Ecole Supérieure des Sciences de l'Aliment et des Industries Agro-Alimentaires, Algiers,
Algeria

*Corresponding authors :

Dr. Oleg MEDIANNIKOV

Address: MEPHI, IRD, APHM, IHU-Méditerranée Infection, 19-21 Boulevard Jean Moulin,
13385 Marseille Cedex 05

Tel: +33 (0)4 13 73 24 01, Fax: +33 (0)4 13 73 24 02, E-mail: olegusss1@gmail.com

Prof. Didier RAOULT

Address: MEPHI, IRD, APHM, IHU-Méditerranée Infection, 19-21 Boulevard Jean Moulin,
13385 Marseille Cedex 05

Tel: +33 (0)4 13 73 24 01, Fax: +33 (0)4 13 73 24 02, E-mail: didier.raoult@gmail.com

Word abstract count: 250

Word text count: 3,325

Abstract

Pediculus humanus is an obligate and highly intimate bloodsucking insect parasite of humans that has two ecotypes, head louse and body louse. This study analyzed genetic diversity at three mitochondrial genes (*cytochrome b* [*cytb*], *cytochrome oxidase subunit 1* [*cox1*] and 12S ribosomal RNA [12S]) in 98 head lice collected from an isolated Native American population from the Wayampi community in Trois-Sauts, French Guiana. These results are integrated with all prior data of *P. humanus* (1402 *cytb*, 743 *cox1* and 344 12S) from other parts of the world. The phylogenetic analysis revealed six highly divergent and well-supported monophyletic clades. Five clades corresponded to the previously recognized mitochondrial clades A, D, B, C and E, while the sixth (clade F) was novel, as it exhibited 5.4%, 3.7% and 3.6% divergence at *cytb*, *cox1* and 12S, respectively, from its nearest neighbor clade B. Interestingly, the clade F has only been recovered in a few lice sequences from Mexico and Argentina, while it was the most common lineage in the Amazonian lice, which hints its association with the Native American region. Furthermore, *Pediculus mjobergi*, a New World monkeys' louse, which is thought to be transmitted to monkeys from the first humans that had reached the American continent thousands of years ago, also belonged to this clade, suggesting that this louse may not be a separate species but an evolutionary lineage of *P. humanus*.

The discovery of new Amazonian clade F with the recovery of additional haplotypes within each of the five clades demonstrates that the levels of genetic diversity in *P. humanus* are higher than previously thought.

Keywords: *Pediculus humanus*; phylogeography; genetic diversity; Clade F; Amazonia

1. Introduction

Sucking lice (Phthiraptera: Anoplura) are obligate blood-feeding ectoparasites of placental mammals that have coevolved with their hosts for more than 65 million years (Durden and Musser, 1994; Light et al., 2010). Humans are parasitized by two species of sucking lice, the pubic louse (*Pthirus pubis*) and head/body lice (*Pediculus humanus*) (Reed et al., 2004). The association between the *P. humanus* and its human host can be traced back to at least 6 million years ago (MYa) to a common ancestor of humans and chimpanzees (Reed et al., 2004). *P. humanus* includes two ecotypes, head lice (*P. h. capitis*) and body lice (*P. h. humanus*), that are morphologically and biologically almost similar but ecologically distinct (Reed et al., 2004; Veracx and Raoult, 2012). Head lice are confined to the scalp and feed on human blood every 4–6 hours (Veracx and Raoult, 2012). Body lice live in clothing and feed less frequently but take larger blood meals than head lice (Veracx and Raoult, 2012). Aside from their role as pests (Chosidow, 2000), body lice are the main vectors of serious human pathogens; *Rickettsia prowazekii* (the causative agent of epidemic typhus), *Bartonella quintana* (trench fever), *Borrelia recurrentis* (relapsing fever) and probably *Yersinia pestis* (plague) (Raoult, 2016; Raoult and Roux, 1999). It was once believed that only body lice could transmit disease. However, recently several combined epidemiological and laboratory studies have strongly implicated head lice as a vector of infectious agents, although its vectorial capacity is lower as compared to body lice (Amanzougaghene et al., 2017, 2016a; Angelakis et al., 2011; Diatta et al., 2014; Goldberger and Anderson, 1912; Kim et al., 2017; Murray and Torrey, 1975; Sangaré et al., 2014).

The genetic diversity of human lice has been extensively studied using mitochondrial genes (mainly *cytochrome b* [*cytb*] and *cytochrome oxidase subunit I* [*coxI*] genes) revealing the presence of five highly divergent mitochondrial clades (A, D, B, C and E) (Amanzougaghene et al., 2016b; Ascunce et al., 2013; Ashfaq et al., 2015; Drali et al., 2015;

76 Kittler et al., 2003; Reed et al., 2004). In addition to this inter-clade diversity, human lice also
77 present intra-clade diversity, illustrated by several distinct haplotypes for each clade
78 (Amanzougaghene et al., 2016a; Ascunce et al., 2013; Light et al., 2008). Body lice belong to
79 clades A and D, while head lice encompass the full genetic diversity of clades
80 (Amanzougaghene et al., 2016a; Drali et al., 2016; Light et al., 2008). Clade A is the most
81 common and widely distributed across all continents, whereas the other clades are
82 geographically restricted (Amanzougaghene et al., 2016b; Ascunce et al., 2013; Light et al.,
83 2008; Raoult et al., 2008). Clade D is restricted to Africa and is found in the Democratic
84 Republic of Congo (DR Congo), the Republic of Congo (Congo-Brazzaville), Ethiopia and
85 Zimbabwe (Amanzougaghene et al., 2016a; Drali et al., 2015). Clade B is found in America,
86 Europe, Australia, Algeria, South Africa, Saudi Arabia and has recently been found among
87 the remains of Israeli head lice, which date back about 2,000 years (Al-Shahrani et al., 2017;
88 Amanzougaghene et al., 2016b; Ascunce et al., 2013; Ashfaq et al., 2015; Boutellis et al.,
89 2015; Light et al., 2008; Raoult et al., 2008). Clade C has been found in some African and
90 Asian countries including Ethiopia, Congo-Brazzaville, Nepal, Pakistan and Thailand
91 (Amanzougaghene et al., 2016a; Ashfaq et al., 2015; Kittler et al., 2003; Raoult et al., 2008;
92 Sunantaraporn et al., 2015). Finally, clade E appears to be specific to West African lice,
93 including Senegal and Mali, where it is highly prevalent, and has recently been identified in
94 the head lice from Nigerian refugees arriving in Algeria and from migrant communities living
95 in Bobigny, France (Amanzougaghene et al., 2017; Candy et al., 2018; Louni et al., 2018).

96 Lice are highly host specific and fast-evolving parasites that have evolved in tandem
97 with their primate hosts for thousands of years (Light et al., 2008; Reed et al., 2007).
98 Previous studies have shown that the time of divergence among lice clades (around 2 MYa)
99 was far anterior to the time of modern human divergence, about 200,000 years ago (Light et
100 al., 2008; Tishkoff et al., 2009). Previous reports have also suggested that different lice clades

have evolved on other archaic hominids, likely those known from 2.3 to 0.03 MYa (such as *Homo erectus*, Neandertal and Denisovan), and have only switched to modern humans during the recent period of overlaps (Ashfaq et al., 2015; Reed et al., 2004). Moreover, the presence of highly divergent clades and their geographical isolation can yield important information regarding the evolutionary history of the lice as well as their human hosts (Ascunce et al., 2013; Light et al., 2008). Therefore, a more detailed analysis of genetic diversity in *P. humanus* and current distributions of its major clades will provide a more detailed picture of evolutionary pattern of this parasite and will clarify additional events in human evolution.

In the present study, we obtained and analyzed head lice collected from an isolated Native American population from the Wayampi community in Trois-Sauts, French Guiana. These results are integrated with all prior mitochondrial data from over the world to expand perspectives on the number, distributions and diversification rates of clades of *Pediculus* lice.

2. Materials and methods

2.1 Ethical approval

The Amazonian head lice were collected from infested Amerindians after obtaining their verbal consent or that of their legal representatives in the case of children, because most subjects were illiterate. Local authorities approved and were present at the lice collection. The study was validated by the ethics committee of the Institut Fédératif de Recherche 48, Faculty of Medicine, Marseille, France, under agreement 12-017.

2.2 Louse samples

The Amazonian head louse specimens were recovered in 2013 from Amerindians of the Wayampi community living in Trois-Sauts (2°15'0" N and 52°52'60" W, 122 m), a remote and isolated village in French Guyana. A total of 98 head lice were recovered from 22 individuals. No body lice were found during the examination. Collected lice were then preserved in 70% ethanol before being sent to our laboratory in Marseille (France). All the

samples were photographed with a camera (Olympus DP71, Rungis, France) prior molecular analysis. In addition, a total of 327 louse specimens were also included in this study, corresponding to body and head lice collected from several countries. These samples were obtained from the private frozen collection of world lice belonging to our laboratory. Additional supporting information including, collection locations and numbers of lice tested are described in Table S1.

Moreover, because there were no available 12S sequences of New World monkey louse *Pediculus mjobergi* in the GenBank database, we have also amplified the 12S gene for this louse. In total three *P. mjobergi* lice were amplified in this study targeting 12S gene. These samples were obtained from the same *P. mjobergi* collection, from our laboratory, that was previously used by Drali et al. (Drali et al., 2016). More specifically, the louse specimens were collected from a wild howler monkey, *Alouatta caraya*, (#B2188) from the Iguazú National Park, Misiones Province (Drali et al., 2016).

2. 3 DNA extraction, PCR amplification and sequencing

Genomic DNA was isolated from louse specimens using the DNeasy tissue kit (Qiagen, Courtaboeuf, France) as described previously (Amanzougaghene et al., 2017). Three mitochondrial genes *cytb*, *coxI* and 12S ribosomal RNA were targeted for sequencing. PCR amplifications were conducted in a Peltier PTC-200 thermal cycler (MJ Research Inc., Watertown, MA, USA) with the Hotstart Taq-polymerase (Qiagen) in accordance with the manufacturer's instructions. The success of PCR amplification was then verified by electrophoresis of the PCR product on a 1.5% agarose gels. All primers used for these experiments and PCR conditions are described in Table 1. All PCR products were purified using the PCR filter plate Millipore NucleoFast 96 PCR kit (Macherey-Nagel EURL, Hoerd, France) following the manufacturer's recommendations. The sequence reaction was carried out using the Big Dye Terminator Cycle Sequencing Kit (Perkin Elmer Applied Biosystems,

Foster City, CA) as per the manufacturer's instructions. Sequencing was performed with an ABI Prism 3130xl Genetic Analyzer capillary sequencer (Applied Biosystems). Finally, all the generated sequences were assembled and corrected using ChromasPro 1.7 software (Technelysium Pty Ltd., Tewantin, Australia).

2. 4 Sequence analysis.

All the 98 Amazonian head louse sequences for the three mitochondrial genes (*cytb*, *cox1* and 12S) obtained in this study were combined with all the available mitochondrial sequence data of *P. humanus* (1402 *cytb*, 743 *cox1* and 344 12S) from other parts of the world to generate a global dataset to examine clade diversity in *P. humanus*. In addition, the newly amplified 12S sequence of *P. mjobergi* in this study, as well as, seven *P. mjobergi* sequences (six *cytb* and one *cox1* sequences) reported by Drali et al. (Drali et al., 2016), were also included in the analysis (Table S2).

The DNA sequences obtained from the literature varied in length, so sequences were trimmed to produce a dataset that maximized the number of sequences incorporated. The sequences between nucleotide positions 433–705 of *cytb* (272-bps, according to Genbank Accession KC685778), 748–1031 of *cox1* (283-bps, according to Genbank Accession KC685838) and 109-666 of 12S (557-bps, according to Genbank Accession KC685877) were used for analysis. ClustalW alignments were performed in MEGA6 (Tamura et al., 2013). Haplotypes were identified using DnaSP v5.10 software (Librado and Rozas, 2009). Finally, we created three datasets for which a total of 1500, 841 and 442 sequences were included, respectively, for *cytb*, *cox1* and 12S (Tables S3-S5).

2. 5 Phylogenetic analysis

Neighbor-joining (NJ) analysis was performed in MEGA6 using the K2P model with pairwise-deletion and 500 bootstrap replicates. The Maximum-likelihood (ML) analysis was also performed in MEGA6 using the Kimura 2-parameter model for nucleotide sequences

under 500 bootstrap replicates. The subtree for each clade of lice was collapsed with the “compress/expand subtree” function. *Cytb*, *cox1* and 12S sequences from *P. schaeffi* (AY695999, KC241887, AY696067, KC241883 and KR706169) were employed as outgroups.

2. 6 Genetic diversity and haplotype analysis

For each dataset, population genetic indices including number of sequences (n), number of polymorphic sites (S), average number of pairwise nucleotide differences (k), nucleotide diversity (π), number of haplotypes (H) and haplotype diversity (Hd) were calculated using DNASP v5.10 software (Librado and Rozas, 2009). Kimura-2-parameter (K2P) pairwise distances among the *cytb*, *Cox1* and 12S haplotypes were calculated using MEGA6 with pairwise deletion of gaps and missing data (Tamura et al., 2013). Neutrality tests (Fu & Li’s D and Tajima’s D) were calculated with DNASP v5.10 (Librado and Rozas, 2009). In order to investigate the possible relationships between the haplotypes, networks haplotypes for each of the three genes were constructed with the median joining method of Bandelt available in NETWORK5.0 (www.fluxus-engineering.com/sharenet.htm) using equal weights for all mutations (Bandelt et al., 1999).

3. Results

A total of 98 head lice collected from 22 Amazonian individuals were analyzed, targeting three mitochondrial genes (*cytb*, *cox1* and 12S). We obtained 11 haplotypes of *cytb*, 8 haplotypes of *cox1* and 13 of 12S that were defined by 31, 25 and 41 polymorphic sites, respectively (Fig. S1-S4). The distribution of the head lice haplotypes identified in this study, according to mitochondrial genes, among the 22 infested Amazonian individuals are presented in Table S6. The generated Amazonian sequences (98 sequences for each gene) were then combined with all available sequences for *cytb*, *cox1* and 12S. The number of haplotypes in each dataset within each clade and their distributions were determined for each

gene. The details of the identified haplotypes, their GenBank accession numbers and geographic locations are described in Tables S3-S5.

In addition, six *cytb* and one *cox1* *P. mjobergi* sequences reported by Drali et al. (Drali et al., 2016), as well as, the three *P. mjobergi* 12S sequences amplified in this study were also included in the analysis (Table S2).

For the *cytb* dataset, 1506 sequences were included (1500 sequences of *P. humanus* and 6 sequences of *P. mjobergi*) from which 105 haplotypes, including two haplotypes from *P. mjobergi*, were identified from 45 countries on five continents. For the *cox1* dataset, 842 sequences were included (841 sequences of *P. humanus* and one sequence of *P. mjobergi*) from which 57 haplotypes, including one haplotype from *P. mjobergi*, were identified from 27 countries on five continents. For the 12S dataset, 445 sequences were included (442 sequences of *P. humanus* and 3 sequences of *P. mjobergi*) from which 49 haplotypes, including one haplotype from *P. mjobergi*, were identified from 18 countries on five continents.

Neighbor-joining (NJ) and Maximum-likelihood (ML) analyses, including all haplotypes, were performed for each of the three mtDNA genes, consistently recovered six highly divergent and well-supported monophyletic clades (Fig. 2 and Fig. S5). Five clades corresponded to the mitochondrial clades previously recognized A, D, B, C and E, while the sixth was novel, here named “clade F”.

This novel clade consists mainly of Amazonian head lice (a total of 84 out of 98 [85.7%] Amazonian lice sequences belonged to this clade) as well as one haplotype from the New World monkey louse *P. mjobergi*, while the remaining 14 of the 98 (14.3%) Amazonian lice sequences belonged to clade A.

More precisely, for the 12S gene, the clade F consisted of nine haplotypes; eight haplotypes were for Amazonian lice (referred to here as F19-F26) whose F9 haplotype was

the most common (83.3% of 84 sequences), while the ninth haplotype was from *P. mjobergi*. For the *coxI* gene, 8 haplotypes were identified, six haplotypes were from Amazonian lice (referred to here as F29-F34), one haplotype named F18 consisted of sequences reported by Ascunce et al. (Ascunce et al., 2013) from Argentina (10 *coxI* sequences) and Mexico (2 *coxI* sequences), the eighth haplotype was *P. mjobergi*. Lastly, for the *cytb* gene, the clade F also included nine haplotypes; eight haplotypes were from Amazonian lice (named here F54 and F1-F7) of which haplotype F54 was the most widespread (84.3% of 89 sequences were from Amazonian head lice sequenced in this study, while 15.7% of 89 sequences also from Amazonian lice and were recovered from Genbank database, the remaining haplotype was from *P. mjobergi*. It is important to note that the second *cytb* haplotype of *P. mjobergi* has the *P. humanus*' haplotype A5 within clade A (see Drali et al., 2016).

The median-joining networks for all *cytb*, *coxI* and 12S haplotypes corroborated the neighbor joining and Maximum-likelihood phylogenetic reconstructions, with all the recovered haplogroups forming separate clusters represented by six connected subnetworks corresponding to clades A, D, B, C, E and "F" (Fig. 3-5).

The maximum intra-clades distances at *cytb* were 1.2%, 1.9%, 1.4%, 1.4%, 1.5% and 1.1% for clades A (n= 34 haplotypes), D (n= 17), B (n= 9), C (n= 13), E (n= 23) and "F" (n= 8), respectively. The maximum intra-clades distances at *coxI* were 1.2%, 1.0%, 1.3%, 1.3%, 1.3% and 1.1% for clades A (n= 17), D (n= 7), B (n= 12), C (n= 6), E (n= 7) and "F" (n= 7), respectively. The maximum intra-clades distances at 12S were 0.9%, 1.6%, 0.4%, 1.4%, 0.3% and 0.5% for clades A (n= 15), D (n= 8), B (n= 3), C (n= 10), E (n= 4) and "F" (n= 8), respectively. The nearest neighbors (NN) distances between clades and the nodal supports are presented in Fig. 2A (*cytb*), Fig. 2B (*coxI*) and Fig. 2C (12S). The new clade "F" shows divergence from its NN clade B of 5.4%, 3.7% and 3.6%, respectively, at *cytb*, *coxI* and 12S.

Estimates of genetic diversity indices and the results of neutrality tests for *cytb*, *cox1* and 12S are presented in Table 2. The average number of nucleotide diversity (π), pairwise nucleotide differences (k) and haplotype diversity (Hd) varied among the clades of three genes. The highest haplotype diversity was found within clade D in both *cytb* and 12S genes (Hd= 0.831 and 0.899, respectively, in *cytb* and 12S), while in *cox1* gene, the highest haplotype diversity was found within clade B (Hd = 0.899). Overall, both (k) and (π) were similar in *cytb*, *cox1* and 12S.

4. Discussion

In this study, we analyzed the genetic diversity of head lice collected from Amazonian individuals of the Wayampi community living in Trois-Sauts, a remote and isolated village. In total, three mitochondrial genes were targeted (*cytb*, *cox1* and 12S). By coupling these results with all available mitochondrial data of *P. humanus* from 45 countries, the present study has expanded understanding of its levels, sequence divergence pattern and revealed higher levels of mtDNA diversity in *P. humanus* corroborating results reported by others (Ashfaq et al., 2015). To the best of our knowledge, this is the most geographically widespread study to evaluate the mitochondrial genetic diversity in human lice based on three mtDNA genes. Previous studies on *P. humanus* showed that the maximum distances within clades were 1.4% at *cytb* and 1.9% at *cox1*, while the NN distances at these genes were 4.6% and 2.3%, respectively (Ashfaq et al., 2015). In the present study, the maximum distances within clades were almost similar (*cytb* 1.9%, *cox1* 1.3% and 1.6% 12S), while NN distances were higher (*cytb* 5.6% and *cox1* 6.5%). These results reflect the extended geographic coverage and the large sample sizes, which allowed the recovery of additional clades and haplotypes within each clade.

Three phylogenetic methods (NN, ML and MJ) at three mtDNA genes, have consistently recovered six highly divergent and well-supported monophyletic clades. Five

275 clades corresponded to the previously recognized mitochondrial clades A, D, B, C and E
276 (Amanzougaghene et al., 2016b; Ashfaq et al., 2015), while the sixth (clade F) was novel,
277 consisting mainly of head lice from Amazonian individuals analyzed in this study and 12
278 sequences of head lice from Argentina and Mexico reported by Ascunce et al. (Ascunce et
279 al., 2013). In that study, the authors found these sequences to be highly derived B-haplotypes
280 that are separated from the main group by more than seven mutational steps (Ascunce et al.,
281 2013).

282 Interestingly, the *P. mjobergi* 12S and *cox1* haplotypes, as well as, one of the two *P.*
283 *mjobergi* *cytb* haplotypes previously classified within clade B by Drali et al., (Drali et al.,
284 2016) belong to this new clade F. This result suggests that *P. mjobergi* may not be a distinct
285 species of *P. humanus*, but probably an evolutionary lineage of *P. humanus* species within
286 clades A and F. A similar suggestion was offered by Drali et al. (Drali et al., 2016). Indeed,
287 these authors argued that *P. mjobergi* was originally a human louse and was switched to new
288 world monkeys from the first humans who reached the American continent thousands of
289 years ago (Drali et al., 2016). Further studies are needed to confirm this hypothesis.

290 Clade “F” shows a divergence from its NN clade B of 5.4%, 3.7% and 3.6%,
291 respectively, at *cytb*, *cox1* and 12S. Clade B was first described in contemporary lice in the
292 American continent, where it was highly prevalent and diversified (Ascunce et al., 2013;
293 Light et al., 2008; Reed et al., 2004). This discovery and its identification in the lice of pre-
294 Columbian mummies led researchers to deduce initially an American origin for this clade
295 (Boutellis et al., 2013). However, its recent discovery among head lice remains from Israel,
296 dating back about 2,000 years, has challenged this hypothesis (Amanzougaghene et al.,
297 2016b). In that study, the authors strongly argued in favor of an Asian origin of clade B, that
298 resulted probably from a recent host switch from Neanderthals or Denisovans to modern
299 humans which was followed by its introduction into the New World with the migration of

early peoples (Amanzougaghene et al., 2016b; Ascunce et al., 2013; Light et al., 2008; Reed et al., 2004). Because clade F was found only in head lice from Mexico and Argentina and it was the dominant lineage found in the Amazonian lice, knowing that Amazonia is one of the few places in the world that has not been strongly affected by globalization. This clade F may be the descendants of a pre-Columbian population and was derived from clade B brought by the first humans who had reached the American continent via the Bering straits thousands of years ago, thus representing Native America louse mitochondrial diversity. Interestingly, Ascunce et al. (Ascunce et al., 2013) have also proposed a similar suggestion for their sequences from Mexico and Argentina that we have reclassified here as members of clade F.

Previous studies reported that clade A is the most common and has a global distribution (Amanzougaghene et al., 2017; Ascunce et al., 2013; Ashfaq et al., 2015), results supported by its detection in approximately 46% of the analyzed lice from 49 countries on the five continents. Furthermore, the clade A subnetworks at the three analyzed mitochondrial genes (Fig. 3-5) were star-like in structure, combined with its significant negative Tajima's D value (-2.322, -1.866 and -2.055, respectively at *cytb*, *cox1* and 12S; $P < 0.05$), indicating the signature of population expansion for this clade (Aris-Brosou and Excoffier, 1996), and corroborating the results reported by others (Ascunce et al., 2013; Reed et al., 2004). In addition, Reed et al. (Reed et al., 2004) estimated that the demographic expansion of this clade occurred about 100,000 years ago, coinciding with the out-of-Africa expansion of *Homo sapiens*, thus reflecting a codemographic pattern between lice and humans (Ascunce et al., 2013; Reed et al., 2004). Clade D (referred as clade E in Ashfaq et al. 2015) diverged from clade A between 0.37 and 0.54 MYa (Ashfaq et al., 2015) and is restricted to Central Africa including DR Congo and Congo-Brazzaville, where it was detected mainly among indigenous Pygmy populations (Amanzougaghene et al., 2016a; Drali et al., 2015). This clade

has also been reported in lice in Ethiopia (Amanzougaghene et al., 2016a) and we identified in this study its occurrence, for the first time, in body lice from Zimbabwe.

Our sampling did not encounter any new specimens of clade C, so it remains restricted to Africa and Asia (Amanzougaghene et al., 2016a; Ascunce et al., 2013; Light et al., 2008; Sunantaraporn et al., 2015). Given its early divergence in the *Pediculus* tree around 2 MYa, this clade may have evolved on archaic hominids in Asia or Africa such as *H. erectus* (Light et al., 2008; Reed et al., 2004). Lastly, clade E (referred as clade D in Ashfaq et al, 2015) diverged from the MRCA of clade C between 0.28 and 0.42 MYa (Ashfaq et al., 2015). This clade consists of head lice from West Africa including Senegal and Mali where it was highly prevalent (Amanzougaghene et al., 2017). Its recent detection in the head lice of Nigerian refugees arriving in Algeria and migrant communities living in Bobigny (France) is probably the result of a recent migratory flow from West African countries (Candy et al., 2018; Louni et al., 2018).

5. Conclusion

Our study underlines the importance of the use of mitochondrial genes in the analysis of phylogeographic patterns and genetic diversity of *P. humanus*. Six highly divergent and well-supported monophyletic clades were identified. Five clades corresponded to the previously recognized mitochondrial clades A, D, B, C and E, while the sixth “clade F” was novel. The new clade F was found mainly in Amazonia, where it is also shared with the monkey louse *P. mjobergi* and could therefore represent Native American louse mitochondrial diversity.

The recovery of additional haplotypes within each of the five clades (A, D, C, E and B) along with the discovery of new clade F, demonstrate that levels of genetic diversity in *P.*

humanus is higher than previously thought, reinforcing the importance of continuing to survey and phylogeographically characterize human lice.

Acknowledgments

The authors thank the Amerindian populations of the Wayampi community living in Trois-Sauts for allowing them to collect the louse specimens.

Funding: This study was supported by the Institut Hospitalo-Universitaire (IHU) Méditerranée Infection, the National Research Agency under the program « Investissements d’avenir », reference ANR-10-IAHU-03, the Région Provence-Alpes-Côte d’Azur and the European funding FEDER PRIML.

Figure captions

Figure 1. Map showing the head lice collection site from the Amerindians of the Wayampi community living in Trois-Sauts. A) Geographical localization of louse sampling. B) Amerindian children infested with head lice. C) Human head lice from Trois-Sauts. ♀ female; ♂ male; nymphs and nits.

Figure 2. Neighbor-joining cluster analysis of *Cytb* (A), *Cox1* (B) and 12S (C) haplotypes of *P. humanus*. Bootstrap values (500 replicates) are shown above the branches. The scale bar indicates K2P distances. The node for each clade with multiple haplotypes is collapsed to a vertical triangle, with the horizontal depth indicating the level of intra-clade divergence. Bracketed numbers next to each clade name indicate the number of haplotypes analyzed and the average intra-clade distance. Analyses were conducted in MEGA6

Figure 3. *Cytb* haplotype networks of human body and head lice. Each circle indicates a unique haplotype and variations in circle size are proportional to haplotype frequencies. Pie colors and sizes in circles represent the continents and the number of their sequences for a haplotype. *P. mjobergi* that belong to the haplotype A5 are included in the portion representing human lice from America.

Figure 4. *Cox1* haplotype networks of human body and head lice. Each circle indicates a unique haplotype and variations in circle size are proportional to haplotype frequencies. Pie

colors and sizes in circles represent the continents and the number of their sequences for a haplotype

Figure 5. 12S haplotype networks of human body and head lice. Each circle indicates a unique haplotype and variations in circle size are proportional to haplotype frequencies. Pie colors and sizes in circles represent the continents and the number of their sequences for a haplotype

Table 1. Primer sequences used in this study

Target gene	Primer name	Primer sequences (5'-3')	Product size (bp)	Tm	source
Cytochrome b	cytbF	GAGCGACTGTAATTACTAATC	348	56°C	(Li et al., 2010)
	cytbR	CAACAAAATTATCCGGGTCC			
Cytochrome oxidase subunit 1	cox1F	GGAGTGAGTTCGATTTTAG	828	55°C	This study
	cox1R	GTGCTGAGGAAAGAAAGTC			
12S ribosomal RNA	12SF	CAGCACTAGCGGTCATACAT	596	56°C	This study
	12SR	AATGACGGGCGATATGTAC			

416

417

418

419

420

421

422

Table 2. Analysis of genetic diversity indices and neutrality tests (Fu & Li's D and Tajima's D) on mitochondrial *cytb*, *cox1* and 12S sequences

	n	S	K	π	h	Hd	Fu & Li's D	Tajima's D
<u>Cytb all</u>	<u>1500</u>	96	20.341	0.075	<u>105</u>	0.999	-2,878 (P < 0.05) S*	-0.172 (P > 0.1) NS
Clade A	769	33	3.098	0.011	34	0.750	-3.751 (P < 0.02) S**	-2.322 (P < 0.01) S**
Clade D	69	20	5.044	0.018	17	0.831	-1.425 (P > 0.1) NS	-0.583 (P > 0.1) NS
Clade B	200	13	3.694	0.013	9	0.789	-1.633 (P > 0.1) NS	-1.357 (P > 0.1) NS
Clade F	104	11	2.750	0.010	9	0.700	-1.924 (P < 0.05) S*	-1.757 (P < 0.05) S*
Clade C	205	16	3.744	0.014	13	0.792	-1.471 (P > 0.1) NS	-1.151 (P > 0.1) NS
Clade E	153	26	4.075	0.015	23	0.803	-2.747 (P < 0.05) S*	-1.752 (P > 0.05) NS
<u>COI all</u>	<u>842</u>	67	17.255	0.061	<u>57</u>	0.946	-0.065 (P > 0.1) NS	0.457 (P > 0.1) NS
Clade A	443	21	3.294	0.012	17	0.769	-2.244 (P > 0.05) NS	-1.866 (P < 0.05) S*
Clade D	48	8	2.762	0.010	7	0.739	-0.971 (P > 0.1) NS	-1.318 (P > 0.1) NS
Clade B	157	12	3.697	0.013	12	0.786	-0.564 (P > 0.1) NS	-0.292 (P > 0.1) NS
Clade F	139	8	2.952	0.010	8	0.738	-0.473 (P > 0.1) NS	-0.503 (P > 0.1) NS
Clade C	25	8	3.667	0.012	6	0.782	0.457 (P > 0.1) NS	0.274 (P > 0.1) NS
Clade E	30	8	3.524	0.012	7	0.781	0.321 (P > 0.1) NS	0.415 (P > 0.1) NS
<u>12S all</u>	<u>442</u>	110	26.928	0.048	<u>49</u>	0.964	-0.332 (P > 0.1) NS	-0.130 (P > 0.1) NS
Clade A	145	30	4,886	0.009	15	0.830	-2.502 (P < 0.05) S*	-2.055 (P < 0.05) S*
Clade D	62	23	8.821	0.016	8	0.899	0.337 (P > 0.1) NS	-0.029 (P > 0.1) NS
Clade B	36	3	0.271	0.001	3	0.679	-0.319 (P > 0.1) NS	-1.399 (P > 0.1) NS
Clade F	87	10	2.893	0.005	9	0.746	-1.612 (P > 0.1) NS	-1.589 (P > 0.05) NS
Clade C	69	24	7.622	0.014	10	0.885	-0.096 (P > 0.1) NS	-0.483 (P > 0.1) NS
Clade E	43	3	1.500	0.003	4	0.606	-0.754 (P > 0.1) NS	-0.754 (P > 0.1) NS

425 n: number of sequences; S: number of polymorphic sites; k: average number of pairwise nucleotide
426 differences; π : nucleotide diversity; h: number of haplotypes; Hd: haplotype diversity. Tajima's D: A
427 negative Tajima's D signifies an excess of low frequency polymorphisms relative to expectation. A
428 positive Tajima's D signifies low levels of both low and high frequency polymorphisms. Statistical
429 significance: Not significant, $P > 0.10$.

430

431

432

433

434

435 References

- 436 Al-Shahrani, S.A., Alajmi, R.A., Ayaad, T.H., Al-Shahrani, M.A., Shaurub, E.-S.H., 2017. Genetic
437 diversity of the human head lice, *Pediculus humanus capitis*, among primary school girls in
438 Saudi Arabia, with reference to their prevalence. *Parasitol. Res.* 116, 2637–2643.
439 <https://doi.org/10.1007/s00436-017-5570-3>
- 440 Amanzougaghene, N., Akiana, J., Mongo Ndombe, G., Davoust, B., Nsana, N.S., Parra, H.-J., Fenollar,
441 F., Raoult, D., Mediannikov, O., 2016a. Head Lice of Pygmies Reveal the Presence of
442 Relapsing Fever *Borreliae* in the Republic of Congo. *PLoS Negl. Trop. Dis.* 10, e0005142.
443 <https://doi.org/10.1371/journal.pntd.0005142>
- 444 Amanzougaghene, N., Fenollar, F., Sangaré, A.K., Sissoko, M.S., Doumbo, O.K., Raoult, D.,
445 Mediannikov, O., 2017. Detection of bacterial pathogens including potential new species in
446 human head lice from Mali. *PLOS ONE* 12, e0184621.
447 <https://doi.org/10.1371/journal.pone.0184621>
- 448 Amanzougaghene, N., Mumcuoglu, K.Y., Fenollar, F., Alfi, S., Yesilyurt, G., Raoult, D., Mediannikov,
449 O., 2016b. High Ancient Genetic Diversity of Human Lice, *Pediculus humanus*, from Israel
450 Reveals New Insights into the Origin of Clade B Lice. *PLoS One* 11, e0164659.
451 <https://doi.org/10.1371/journal.pone.0164659>
- 452 Angelakis, E., Diatta, G., Abdissa, A., Trape, J.-F., Mediannikov, O., Richet, H., Raoult, D., 2011.
453 Altitude-dependent *Bartonella quintana* genotype C in head lice, Ethiopia. *Emerg. Infect. Dis.*
454 17, 2357–2359. <https://doi.org/10.3201/eid1712.110453>
- 455 Aris-Brosou, S., Excoffier, L., 1996. The impact of population expansion and mutation rate
456 heterogeneity on DNA sequence polymorphism. *Mol. Biol. Evol.* 13, 494–504.
457 <https://doi.org/10.1093/oxfordjournals.molbev.a025610>
- 458 Ascunce, M.S., Fane, J., Kassu, G., Toloza, A.C., Picollo, M.I., González-Oliver, A., Reed, D.L., 2013.
459 Mitochondrial diversity in human head louse populations across the Americas. *Am. J. Phys.*
460 *Anthropol.* 152, 118–129. <https://doi.org/10.1002/ajpa.22336>
- 461 Ashfaq, M., Prosser, S., Nasir, S., Masood, M., Ratnasingham, S., Hebert, P.D.N., 2015. High diversity
462 and rapid diversification in the head louse, *Pediculus humanus* (Pediculidae: Phthiraptera).
463 *Sci. Rep.* 5. <https://doi.org/10.1038/srep14188>
- 464 Bandelt, H.J., Forster, P., Röhl, A., 1999. Median-joining networks for inferring intraspecific
465 phylogenies. *Mol. Biol. Evol.* 16, 37–48.
466 <https://doi.org/10.1093/oxfordjournals.molbev.a026036>
- 467 Boutellis, A., Bitam, I., Fekir, K., Mana, N., Raoult, D., 2015. Evidence that clade A and clade B head
468 lice live in sympatry and recombine in Algeria. *Med. Vet. Entomol.* 29, 94–98.
469 <https://doi.org/10.1111/mve.12058>
- 470 Boutellis, A., Drali, R., Rivera, M.A., Mumcuoglu, K.Y., Raoult, D., 2013. Evidence of sympatry of clade
471 a and clade B head lice in a pre-Columbian Chilean mummy from Camarones. *PLoS One* 8,
472 e76818. <https://doi.org/10.1371/journal.pone.0076818>
- 473 Candy, K., Amanzougaghene, N., Izri, A., Brun, S., Durand, R., Louni, M., Raoult, D., Fenollar, F.,
474 Mediannikov, O., 2018. Molecular Survey of Head and Body Lice, *Pediculus humanus*, in
475 France. *Vector Borne Zoonotic Dis. Larchmt. N.* <https://doi.org/10.1089/vbz.2017.2206>
- 476 Chosidow, O., 2000. Scabies and pediculosis. *The Lancet* 355, 819–826.
477 [https://doi.org/10.1016/S0140-6736\(99\)09458-1](https://doi.org/10.1016/S0140-6736(99)09458-1)
- 478 Diatta, G., Mediannikov, O., Sokhna, C., Bassene, H., Socolovschi, C., Ratmanov, P., Fenollar, F.,
479 Raoult, D., 2014. Prevalence of *Bartonella quintana* in patients with fever and head lice from
480 rural areas of Sine-Saloum, Senegal. *Am. J. Trop. Med. Hyg.* 91, 291–293.
481 <https://doi.org/10.4269/ajtmh.13-0685>

482 Drali, R., Abi-Rached, L., Boutellis, A., Djossou, F., Barker, S.C., Raoult, D., 2016. Host switching of
483 human lice to new world monkeys in South America. *Infect. Genet. Evol. J. Mol. Epidemiol.*
484 *Evol. Genet. Infect. Dis.* 39, 225–231. <https://doi.org/10.1016/j.meegid.2016.02.008>

485 Drali, R., Shako, J.-C., Davoust, B., Diatta, G., Raoult, D., 2015. A New Clade of African Body and Head
486 Lice Infected by *Bartonella quintana* and *Yersinia pestis*-Democratic Republic of the Congo.
487 *Am. J. Trop. Med. Hyg.* 93, 990–993. <https://doi.org/10.4269/ajtmh.14-0686>

488 Durden, L.A., Musser, G.G., 1994. The sucking lice (Insecta, Anoplura) of the world : a taxonomic
489 checklist with records of mammalian hosts and geographical distributions. *Bulletin of the*
490 *AMNH* ; no. 218. Sucking lice and hosts.

491 Goldberger, J., Anderson, J.F., 1912. The Transmission of Typhus Fever, with Especial Reference to
492 Transmission by the Head Louse (*Pediculus capitis*). *Public Health Rep.* 1896-1970 27, 297–
493 307. <https://doi.org/10.2307/4567527>

494 Kim, J.H., Previte, D.J., Yoon, K.S., Murenzi, E., Koehler, J.E., Pittendrigh, B.R., Lee, S.H., Clark, J.M.,
495 2017. Comparison of the proliferation and excretion of *Bartonella quintana* between body
496 and head lice following oral challenge. *Insect Mol. Biol.* 26, 266–276.
497 <https://doi.org/10.1111/imb.12292>

498 Kittler, R., Kayser, M., Stoneking, M., 2003. Molecular evolution of *Pediculus humanus* and the origin
499 of clothing. *Curr. Biol. CB* 13, 1414–1417.

500 Li, W., Ortiz, G., Fournier, P.-E., Gimenez, G., Reed, D.L., Pittendrigh, B., Raoult, D., 2010. Genotyping
501 of human lice suggests multiple emergencies of body lice from local head louse populations.
502 *PLoS Negl. Trop. Dis.* 4, e641. <https://doi.org/10.1371/journal.pntd.0000641>

503 Librado, P., Rozas, J., 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism
504 data. *Bioinforma. Oxf. Engl.* 25, 1451–1452. <https://doi.org/10.1093/bioinformatics/btp187>

505 Light, J.E., Allen, J.M., Long, L.M., Carter, T.E., Barrow, L., Suren, G., Raoult, D., Reed, D.L., 2008.
506 Geographic distributions and origins of human head lice (*Pediculus humanus capitis*) based
507 on mitochondrial data. *J. Parasitol.* 94, 1275–1281. <https://doi.org/10.1645/GE-1618.1>

508 Light, J.E., Smith, V.S., Allen, J.M., Durden, L.A., Reed, D.L., 2010. Evolutionary history of mammalian
509 sucking lice (Phthiraptera: Anoplura). *BMC Evol. Biol.* 10, 292. [https://doi.org/10.1186/1471-](https://doi.org/10.1186/1471-2148-10-292)
510 2148-10-292

511 Louni, M., Amanzougaghene, N., Mana, N., Fenollar, F., Raoult, D., Bitam, I., Mediannikov, O., 2018.
512 Detection of bacterial pathogens in clade E head lice collected from Niger’s refugees in
513 Algeria. *Parasit. Vectors* 11, 348. <https://doi.org/10.1186/s13071-018-2930-5>

514 Murray, E.S., Torrey, S.B., 1975. Virulence of *Rickettsia prowazeki* for head lice. *Ann. N. Y. Acad. Sci.*
515 266, 25–34.

516 Raoult, D., 2016. A Personal View of How Paleomicrobiology Aids Our Understanding of the Role of
517 Lice in Plague Pandemics. *Microbiol. Spectr.* 4. [https://doi.org/10.1128/microbiolspec.PoH-](https://doi.org/10.1128/microbiolspec.PoH-0001-2014)
518 0001-2014

519 Raoult, D., Reed, D.L., Dittmar, K., Kirchman, J.J., Rolain, J.-M., Guillen, S., Light, J.E., 2008. Molecular
520 identification of lice from pre-Columbian mummies. *J. Infect. Dis.* 197, 535–543.
521 <https://doi.org/10.1086/526520>

522 Raoult, D., Roux, V., 1999. The body louse as a vector of reemerging human diseases. *Clin. Infect. Dis.*
523 *Off. Publ. Infect. Dis. Soc. Am.* 29, 888–911. <https://doi.org/10.1086/520454>

524 Reed, D.L., Light, J.E., Allen, J.M., Kirchman, J.J., 2007. Pair of lice lost or parasites regained: the
525 evolutionary history of anthropoid primate lice. *BMC Biol.* 5, 7.
526 <https://doi.org/10.1186/1741-7007-5-7>

527 Reed, D.L., Smith, V.S., Hammond, S.L., Rogers, A.R., Clayton, D.H., 2004. Genetic Analysis of Lice
528 Supports Direct Contact between Modern and Archaic Humans. *PLOS Biol.* 2, e340.
529 <https://doi.org/10.1371/journal.pbio.0020340>

530 Sangaré, A.K., Boutellis, A., Drali, R., Socolovschi, C., Barker, S.C., Diatta, G., Rogier, C., Olive, M.-M.,
531 Doumbo, O.K., Raoult, D., 2014. Detection of *Bartonella quintana* in African body and head
532 lice. *Am. J. Trop. Med. Hyg.* 91, 294–301. <https://doi.org/10.4269/ajtmh.13-0707>

533 Sunantaraporn, S., Sanprasert, V., Pengsakul, T., Phumee, A., Boonserm, R., Tawatsin, A., Thavara,
 534 U., Siriyasatien, P., 2015. Molecular survey of the head louse *Pediculus humanus capitis* in
 535 Thailand and its potential role for transmitting *Acinetobacter* spp. *Parasit. Vectors* 8, 127.
 536 <https://doi.org/10.1186/s13071-015-0742-4>
 537 Tamura, K., Stecher, G., Peterson, D., Filipski, A., Kumar, S., 2013. MEGA6: Molecular Evolutionary
 538 Genetics Analysis Version 6.0. *Mol. Biol. Evol.* 30, 2725–2729.
 539 <https://doi.org/10.1093/molbev/mst197>
 540 Tishkoff, S.A., Reed, F.A., Friedlaender, F.R., Ehret, C., Ranciaro, A., Froment, A., Hirbo, J.B.,
 541 Awomoyi, A.A., Bodo, J.-M., Doumbo, O., Ibrahim, M., Juma, A.T., Kotze, M.J., Lema, G.,
 542 Moore, J.H., Mortensen, H., Nyambo, T.B., Omar, S.A., Powell, K., Pretorius, G.S., Smith,
 543 M.W., Thera, M.A., Wambebe, C., Weber, J.L., Williams, S.M., 2009. The Genetic Structure
 544 and History of Africans and African Americans. *Science* 324, 1035–1044.
 545 <https://doi.org/10.1126/science.1172257>
 546 Veracx, A., Raoult, D., 2012. Biology and genetics of human head and body lice. *Trends Parasitol.* 28,
 547 563–571. <https://doi.org/10.1016/j.pt.2012.09.003>
 548

Supporting Information

Figure S1. (A) Cytb sequences alignment and (B) DNA sequence chromatograms of clade F haplotypes identified in this study showing the polymorphic sites.

Figure S2. (A) Cox1 sequences alignment and (B) DNA sequence chromatograms of clade F haplotypes identified in this study showing the polymorphic sites.

Figure S2. (A) 12S sequences alignment and (B) DNA sequence chromatograms of clade F haplotypes identified in this study showing the polymorphic sites.

Figure S4. (A) 12S sequences alignment of clade F haplotypes and (B) 12S DNA sequence chromatogram of *P. mjobergi* amplified in this study.

Figure S5. Maximum-likelihood (ML) analysis of *Cytb* (A), *Cox1* (B) and 12S (C) haplotypes of *Pediculus humanus*. Bootstrap values (500 replicates) are shown above the branches. The scale bar shows K2P distances. The node for each clade with multiple haplotypes is collapsed to a vertical triangle, with the horizontal depth indicating the level of intra-clade divergence. Bracketed numbers next to each clade's name indicate the number of haplotypes analyzed and the average intra-clade distance. Analyses were conducted in MEGA6.

Table S1. Additional louse specimens included in this study, obtained from the private frozen collection of world lice belonging to our laboratory.

Table S2. *Pediculus mjobergi* sequences from new world monkey (*Alouatta caraya*) included in this study. The *cox1* and *cytb* *P. mjobergi* sequences analyzed in this study were those reported by Drali et al. (2016) collected from two monkey individuals B2188 and B1395. The 12S sequences were amplified in this study from three *P. mjobergi* specimens from monkey individual B2188.

572 **Table S3. Geographic occurrences and frequencies of *cytb* haplotypes of human head**
573 **and body lice.**

574 **Table S4. Geographic occurrences and frequencies of *cox1* haplotypes of human head**
575 **and body lice.**

576 **Table S5. Geographic occurrences and frequencies of 12S haplotypes of human head**
577 **and body lice.**

578 **Table S6. Distribution of the head lice haplotypes identified in this study, according to**
579 **mitochondrial genes, among the 22 infested Amazonian individuals.**

580

A)

0,02

B)

0,02

C)

0,02

A **C** **E**

● Africa
 ● Asia
 ● Oceania
 ● America
 ● Europe

D **B** **F**

P. mjobergi

- Africa
- Asia
- Oceania
- America
- Europe

D **A** **E** **B** **F** **C**

Head lice from an isolated Native American population were analyzed and integrated with all prior data of *P. humanus* from over the world

⇒ New World monkey louse, *P. mjobergi*, was also included in the analysis