

Complete Genome Sequence of *Escherichia coli* BE104, an MC4100 Derivative Lacking the Methionine Reductive Pathway

Brian Anton, Richard Morgan, Benjamin Ezraty, Bruno Manta, Frederic
Barras, Mehmet Berkmen

► **To cite this version:**

Brian Anton, Richard Morgan, Benjamin Ezraty, Bruno Manta, Frederic Barras, et al.. Complete Genome Sequence of *Escherichia coli* BE104, an MC4100 Derivative Lacking the Methionine Reductive Pathway. *Microbiology Resource Announcements*, American Society for Microbiology, 2019, 8 (28), pp.e00721-19. 10.1128/MRA.00721-19 . hal-02311061

HAL Id: hal-02311061

<https://hal-amu.archives-ouvertes.fr/hal-02311061>

Submitted on 10 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



1 **Complete genome sequence of *E. coli* BE104, an MC4100 derivative lacking the methionine**
2 **reductive pathway**

3

4 Brian P. Anton¹, Rick Morgan¹, Benjamin Ezraty², Bruno Manta¹, Frédéric Barras² and Mehmet
5 Berkmen^{1*}

6

7

8

9

10 ¹New England Biolabs, 240 County Road, Ipswich, MA 01938, USA

11 ²Aix-Marseille Université, CNRS, Laboratoire de Chimie Bactérienne, Marseille, France.

12

13 *Correspondence: Phone: (978) 380-7519; Fax: (978) 921-1350; Email: berkmen@neb.com

14

15

16

17

18

19 **Keywords:** *Escherichia coli*, MC4100, methionine sulfoxide reductase

20 **Abstract:**

21 *Escherichia coli* and its derivatives have played a profound role in the development of
22 molecular biology in the last 50 years. With the advent of the genomic age, many *E. coli* strains
23 are being sequenced, adding to our knowledge of the *E. coli* pan genome and its biology. Here
24 we present the complete, annotated genome sequence of an *E. coli* MC4100 mutant strain. This
25 strain has several methionine sulfoxide reductase deletions, making it ideal for studying enzymes
26 that alter the redox state of methionine.

27

28 **Announcement:**

29 *Escherichia coli* is the predominant prokaryotic model organism. Its use in molecular
30 biology has led to enzyme discovery, the development of myriad research tools, and the
31 understanding of many biological principles. Not surprisingly, over the past half a century, many
32 mutant *E. coli* strains have been genetically engineered or selected.

33 There are two popular ancestral strains of *E. coli*, K-12 and B [1]. In general, *E. coli* K-12
34 is used for DNA manipulation and cloning, while *E. coli* B is mostly used for protein expression.
35 There are many *E. coli* K-12 derivatives currently used in research and production laboratories
36 worldwide, including MC4100, engineered by Malcolm Casadaban [2]. Of the 801 completely
37 sequenced *E. coli* genomes, only 33 are from K-12 derived laboratory strains and of that only 1
38 is from MC4100. MC4100 has been used extensively since its inception [3], resulting in
39 thousands of publications. Its genome has been previously sequenced [4] and its relationship to
40 other *E. coli* K-12 strains has been analyzed [5].

41 The strain BE104, a derivative of MC4100 methionine auxotroph mutant lacking five
42 methionine sulfoxide reductases (Msrs), was previously constructed in order to characterize an

43 enzymatic system (MsrPQ) responsible for repairing proteins containing methionine sulfoxide in
44 the bacterial periplasm [6]. Briefly, BE104 was derived from MC4100 a methionine auxotroph
45 mutant by (1) a series of P1vir crosses to delete all cytoplasmic msrs (*msrA*, *msrB*, *msrC* and
46 *bisC*) by replacement with corresponding alleles from Keio KO strains [7]; (2) selection for
47 suppressor strains that could reduce methionine sulfoxide; and (3) deletion of *msrP*, a
48 consequently discovered periplasmic msr [5].

49 As BE104 is being used in our research and will be further engineered, we sequenced its
50 genome using the Pacific Biosciences (PacBio) RS II sequencing platform as described
51 previously [8]. Genomic DNA was isolated using the Monarch gDNA kit (New England
52 Biolabs). A SMRTbell library was constructed from 5 µg genomic DNA, sheared to ~10 kb
53 using a gTube (Covaris). The library was sequenced on two SMRT cells using P6-C4 chemistry.
54 The first cell yielded 258 Mb sequence from 22,107 (15%) P1 reads with mean polymerase read
55 length of 11,672 and mean read of insert length of 6096 (180-minute data collection time). A
56 second cell was sequenced to increase coverage, yielding 905 Mb sequence from 63,262 (42%)
57 P1 reads with mean polymerase read length of 14,318 and mean read of insert of 4594 (240-
58 minute data collection time). Sequencing reads were processed and assembled with Pacific
59 Biosciences SMRT analysis v2.3.0 software using the HGAP3 protocol (5 Mb expected genome
60 size, filters set to: minimum subread length 1000, minimum polymerase read length 2000,
61 minimum read quality 0.80) and polished using Quiver (4). The 1.1 Gb of sequence assembled
62 into a single closed circular genome of 4,xxx,xxx bp with mean 200-fold coverage. The
63 assembled sequence was annotated using the NCBI Prokaryotic Genomes Annotation Pipeline
64 (PGAP).

65

66 The following expected deletions were all observed: *metB* (*O*-succinyl homoserine
67 lyase) is disrupted by 2 base deletion/frameshift, making this strain auxotrophic for methionine;
68 *msrA* is disrupted by insertion with a spectinomycin cassette; the *msr* loci *msrB/yeaA*,
69 *msrC/yebR*, and *msrP/yedY* are all deleted, as is *bisC* (biotin sulfoxide reductase), an enzyme
70 with weak *msr* activity on free S-methionine sulfoxide [9]. As expected, *yedV::IS2* insertion was
71 not observed as *msrP::kan* deletion was transduced using the kanamycin marker from Keio
72 collection (Strain#), co-transducing wt *yedV* allele. Analysis of the genomic sequence revealed
73 several other changes with respect to the parental strain MC4100 in addition to these expected
74 deletions.

75 This strain should be of general use to the research community studying protein redox
76 states and would be an important addition to the repertoire of sequenced *E. coli* genomes.

77

78 **Nucleotide sequence accession number.** The annotated complete genome sequence is
79 deposited at EMBL, EBI under the following assembly accession numbers:

80

81 **References:**

- 82 [1] F.W. Studier, P. Daegelen, R.E. Lenski, S. Maslov, J.F. Kim, Understanding the differences
83 between genome sequences of *Escherichia coli* B strains REL606 and BL21(DE3) and
84 comparison of the *E. coli* B and K-12 genomes, *J Mol Biol*, 394 (2009) 653-680.
85 [2] Remembering Malcolm J. Casadaban, *J Bacteriol*, 192 (2010) 4261-4263.
86 [3] M.J. Casadaban, Transposition and fusion of the *lac* genes to selected promoters in
87 *Escherichia coli* using bacteriophage lambda and Mu, *J Mol Biol*, 104 (1976) 541-555.
88 [4] T. Ferenci, Z. Zhou, T. Betteridge, Y. Ren, Y. Liu, L. Feng, P.R. Reeves, L. Wang, Genomic
89 sequencing reveals regulatory mutations and recombinational events in the widely used MC4100
90 lineage of *Escherichia coli* K-12, *J Bacteriol*, 191 (2009) 4025-4029.
91 [5] J.E. Peters, T.E. Thate, N.L. Craig, Definition of the *Escherichia coli* MC4100 genome by
92 use of a DNA array, *J Bacteriol*, 185 (2003) 2017-2021.
93 [6] A. Gennaris, B. Ezraty, C. Henry, R. Agrebi, A. Vergnes, E. Oheix, J. Bos, P. Leverrier, L.
94 Espinosa, J. Szewczyk, D. Vertommen, O. Iranzo, J.F. Collet, F. Barras, Repairing oxidized
95 proteins in the bacterial envelope using respiratory chain electrons, *Nature*, 528 (2015) 409-412.

96 [7] T. Baba, T. Ara, M. Hasegawa, Y. Takai, Y. Okumura, M. Baba, K.A. Datsenko, M. Tomita,
97 B.L. Wanner, H. Mori, Construction of *Escherichia coli* K-12 in-frame, single-gene knockout
98 mutants: the Keio collection, Mol Syst Biol, 2 (2006) 2006 0008.
99 [8] C.S. Chin, D.H. Alexander, P. Marks, A.A. Klammer, J. Drake, C. Heiner, A. Clum, A.
100 Copeland, J. Huddleston, E.E. Eichler, S.W. Turner, J. Korlach, Nonhybrid, finished microbial
101 genome assemblies from long-read SMRT sequencing data, Nat Methods, 10 (2013) 563-569.
102 [9] B. Ezraty, J. Bos, F. Barras, L. Aussel, Methionine sulfoxide reduction and assimilation in
103 *Escherichia coli*: new role for the biotin sulfoxide reductase BisC, J Bacteriol, 187 (2005) 231-
104 237.
105