Identification of novel species of marine magnetotactic bacteria affiliated with Nitrospirae phylum

Xin-xin Qian, J Liu, Nicolas Menguy, Jinhua Li, François Alberto, Zhaojie Teng, Tian Xiao, Wenyan Zhang, Long-fei Wu

► To cite this version:

Xin-xin Qian, J Liu, Nicolas Menguy, Jinhua Li, François Alberto, et al.. Identification of novel species of marine magnetotactic bacteria affiliated with Nitrospirae phylum. *Environmental Microbiology Reports*, 2019, 11 (3), pp.330-337. 10.1111/1758-2229.12755 . hal-02333904

HAL Id: hal-02333904

<https://hal-amu.archives-ouvertes.fr/hal-02333904>

Submitted on 10 Feb 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Identification of novel species of marine magnetotactic bacteria affiliated with

***Nitrospirae* phylum**

Xin-Xin Qian ^{a,b,1}, Jia Liu ^{b,d,1,2}, Nicolas Menguy ^{b,c}, Jinhua Li ^{b,e}, François Alberto ^{a,b}, Zhaojie
Teng ^d, Tian Xiao ^{b,d}, Wenyan Zhang ^{b,d,3}, Long-Fei Wu ^{a,b,3}

a Aix Marseille University, CNRS, LCB, Marseille, 13402, France

b International Associated Laboratory of Evolution and Development of Magnetotactic
Multicellular Organisms (LIA-MagMC), CNRS-CAS, Marseille, 13402, France

c Institut de Minéralogie, de Physique des Matériaux et de Cosmochimie, UMR 7590 CNRS-
Sorbonne Université, F-75005 Paris. France

d CAS Key Laboratory of Marine Ecology and Environmental Sciences, Institute of
Oceanology, Chinese Academy of Sciences, Qingdao, 266071, China

e Key Laboratory of Earth and Planetary Physics, Institute of Geology and Geophysics,
Chinese Academy of Sciences, Beijing, 100029, China

1 Contributed equally to this work

2 Current address: Key Laboratory of Earth and Planetary Physics, Institute of Geology and
Geophysics, Chinese Academy of Sciences, Beijing, 100029, China

3 To whom correspondence should be addressed: Long-Fei WU, LCB, CNRS, 31 chemin
Joseph Aiguier, 13402 Marseille cedex 20, France, 33-4-9116 4157, wu@imm.cnrs.fr;
Wenyan ZHANG, CAS Key Laboratory of Marine Ecology and Environmental Sciences,
Institute of Oceanology, Chinese Academy of Sciences, Qingdao, 266071, China, 86-532
8289 8586, zhangwy@qdio.ac.cn

Summary

Magnetotactic bacteria (MTB) are a group of Gram-negative bacteria characterized by synthesizing magnetosomes and swimming along geomagnetic field lines. Phylogenetically, they belong to different taxonomic lineages including *Proteobacteria*, *Nitrospirae*, *Omnitrophica*, *Latescibacteria*, and *Planctomycetes* phyla on the phylogenetic tree. To date, six *Nitrospirae* MTB phylotypes have been identified from freshwater or low salinity environments and described in the literature. Here we report the identification of two *Nitrospirae* MTB phylotypes collected, for the first time, from the marine environment. Both have a spherical morphology with a cell size of ~ 5 μ M and similar motility but are different colors (black-brown and ivory-white) under the optic microscope. They synthesized bullet-shaped iron-oxide magnetosomes that were arranged in multiple bundles of chains. Moreover, the cytoplasm of the black-brown *Nitrospirae* MTB contained sulfur inclusions that conferred on cells a rough, granular appearance. Phylogenetic analysis based on their 16S rRNA gene sequences revealed that they are two novel species and cluster with the previously reported MTB affiliated with the phylum *Nitrospirae*, thus extending the distribution of *Nitrospirae* MTB from freshwater to the marine environment.

Originality-Significance Statement

Magnetotactic bacteria (MTB) consist of a phylogenetic, physiological and morphologic heterogeneous group of gram-negative microbes. They are distributed worldwide at the oxic-anoxic interface in chemically stratified water columns or sediments. To date, despite *Nitrospirae* being one of the major phyla of MTB, they have only been found in freshwater or

low-salinity environments. Here we report the identification of two novel species of *Nitrospirae* MTB, from the marine habitat for the first time. They have a spherical morphology and are relatively big in size approximately 5 x 3 μm . One contains abundant intracellular sulfur inclusions. This study shows the great diversity of *Nitrospirae* phylum and extends the distribution of *Nitrospirae* MTB into marine habitats.

Introduction

Magnetotactic bacteria (MTB) are a group of Gram-negative bacteria and exhibit the unique capacity of aligning and swimming along the geomagnetic field lines, a behavior referred to as magnetotaxis (Uebe and Schüler, 2016). This capacity is based on the intracellular biomineralization of magnetite or greigite magnetic nanocrystal chains that give MTB cells a net magnetic moment dipole. MTB present various morphotypes including cocci, spirilla, rod-shaped, vibrio, and more complex multicellular aggregates, and have been found in aquatic environments from freshwater to marine ecosystems (Blakemore, 1975; Bazylinski and Frankel, 2004; Lefèvre and Bazylinski, 2013). Phylogenetically, magnetotactic bacteria are members of several classes of the *Proteobacteria* phylum including the *Alpha*-, *Gamma*-, *Delta*-, *Zeta*-, candidate *Lambda*-, candidate *Eta*-classes, the *Nitrospirae* phylum, the candidate *Omnitrophica* phylum, the candidate *Latescibacteria* phylum and the *Planctomycetes* phylum (Lin et al., 2017a; Lin et al., 2018).

Compared with the *Proteobacteria* phylum, few MTB have been found to be phylogenetically affiliated with the phylum *Nitrospirae* and none of them have been cultured. This is consistent with the general difficulty in cultivating MTB due to their fastidious

physiological requirements (Bazylinski and Frankel, 2004). The first to be discovered and the most studied was *Candidatus Magnetobacterium bavaricum* (Spring et al., 1993). *Ca. Magnetobacterium bavaricum* is a large, rod-shaped bacterium ranging from 8 to 10 μm in length, and contains 600-1000 bullet-shaped magnetite magnetosomes in 3-5 bundles of multiple chains (Jogler et al., 2010; Jogler et al., 2011). *Candidatus Magnetobacterium casensis* shares similar morphology and 98.2% of the sequence identity of the 16S rRNA gene with *Ca. Magnetobacterium bavaricum*. Analysis of the draft genome sequence of *Ca. Magnetobacterium casensis* reveals an autotrophic lifestyle. Pathways involved in denitrification, sulfur oxidation and sulfate reduction have also been predicted, which indicates its considerable capacity to adapt to variable environments (Lin et al., 2014).

The third identified *Nitrospirae* MTB was a small rod-shaped bacterium collected from sediment of Waller See, in Germany and named MHB-1 (Flies et al., 2005). This organism contains a single bundle of multiple chains of magnetosomes with bullet-shaped magnetite crystals. The fourth reported *Nitrospirae* MTB is a moderately thermophilic magnetotactic bacterium, HSMV-1, which remained viable and motile in samples kept between 25 and 62°C for several months (Lefèvre et al., 2010). HSMV-1 cells are small vibrios and biomineralize a single chain of bullet-shaped magnetosomes. Two *Nitrospirae* MTB phylotypes named LO-1 and MWB-1 were identified from southern Nevada, USA and Beijing, China, respectively (Lefevre et al., 2011; Lin et al., 2012a). The two species have 6% phylogenetic divergence and possess a similar relatively big ($\sim 3 \times 2.5\mu\text{m}$) spherical morphology with bullet-shaped magnetosomes and 4-7 bundles of magnetosome chains parallel to the long axis of cells.

Interestingly, the six described *Nitrospirae* MTB phylotypes have distinct cell

appearance and different magnetosome arrangement including single chain, multiple chains with single bundle or multiple bundles. The obvious common feature is their bullet-shape magnetite magnetosomes. Another common property of the reported *Nitrospirae* MTB is that they dwell in low salinity habitats (< 3 ppt) (Spring et al., 1993; Flies et al., 2005; Lefevre et al., 2011; Lin et al., 2012a; Lin et al., 2012b). Therefore, it has been proposed that salinity might influence the distribution of MTB affiliated with *Nitrospirae* phylum (Lin et al., 2012b). Here, we identified two novel *Nitrospirae* MTBs species and expand the distribution of *Nitrospirae* MTB to marine ecosystems.

Results and discussion

Morphology and swimming behaviour of black-brown cocci and ivory-white cocci

Differential interference contrast (DIC) microscopic examination showed the morphology of black-brown giant cocci (BGC) (Fig. 1 A, black arrows), ivory-white giant cocci (WGC) (Fig. 1 A, white arrows) and small cocci of MTB collected from the sediments in Six-Fours les Plages. BGC have a rough surface appearance with an average length of $5.39 \pm 0.72 \mu\text{m}$ and an average width of $4.04 \pm 0.33 \mu\text{m}$ (n=62). WGC have a smoother cell surface and a similar cell size with an average length of $5.21 \pm 0.66 \mu\text{m}$ and an average width of $3.99 \pm 0.30 \mu\text{m}$ (n=62) (B in Figure 1). During sample preparation for DNA amplification and sequencing, we observed that BGC cells began to disintegrate 5 min following their transfer from seawater to PBS buffer, and small granules were released from the cells (C1 in Figure 1). In contrast, WGC cells shrunk and became smaller in PBS buffer (C2 in Figure 1). Further analysis showed that approximately 50 % of the BGC and WGC cells remained motile and

accumulated at the north edge of the cell suspension droplets where the salinity was decreased from 35 to 16 ppt by dilution with deionized water. The number of cells at the edge of the droplets decreased as the dilution increased, and the BGC and WGC cells almost disappeared completely from the droplets at 13 ppt salinity. This observation suggests a sensitivity of the marine cells to the freshwater conditions. BGC and WGC also exhibited different fluorescent properties. When excited with green light (510-550 nm), BGC emitted red fluorescence while no auto-fluorescence signals were observed in WGC under these conditions (D2 in Figure1). BGC emitted stronger yellow auto-fluorescence than WGC under blue light (450-480 nm) excitation (D3 in Figure1). Similar intensity of green light was emitted by both BGC and WGC when excited by violet light (400-410 nm) (D4 in Figure1). WGC emitted stronger blue fluorescence under ultraviolet light (330-385 nm) compared to BGC (D5 in Figure 1). Illumination at these wavelengths had no impact on the motility of these MTB cells. Lack of culture samples prevents us from performing detailed spectrophotometry analysis to characterize the nature of the fluorescent molecules.

Most of the BGC and WGC exhibited north-seeking magnetotaxis behavior, i.e. swimming along the applied magnetic field lines, whereas 10% of cells were found to be south-seeking. Cells demonstrated a dynamic motility in the hanging drops for approximately 30 min then began to slow down gradually and stopped swimming at the end of one hour. The average swim velocities of BGC and WGC cells were $126.3 \pm 45.0 \mu\text{m}\cdot\text{s}^{-1}$ (n=52) and $148.9 \pm 28.0 \mu\text{m}\cdot\text{s}^{-1}$ (n=54), respectively (E in Figure 1). The corresponding maximal swim speeds of BGC ($225.4 \mu\text{m}\cdot\text{s}^{-1}$) and WGC ($188.6 \mu\text{m}\cdot\text{s}^{-1}$) were slower than the closest freshwater relative MTB MWB-1 ($260 \mu\text{m}\cdot\text{s}^{-1}$).

Phylogenetic analysis

In the environmental samples analyzed, BGC cells were the dominant MTB while WGC were approximately 4-fold less abundant as estimated by direct counting of the two morphotypes in droplets (Experimental Procedures and Supporting Information Figure S1). We collected 8 WGC cells using the magnetic micro-sorting approach, and performed Whole Genome Amplification to get sufficient template DNA for further 16S rRNA gene amplification and cloning. We obtained 24 sequences of 16S rRNA genes that shared >99.3% identity between each other and thus formed one operational taxonomic unit (OTU-1, GenBank number MK203828). In parallel, we obtained 40 full-length 16S rRNA gene sequences of MTB prepared by a routine two-step magnetic collection (see Experimental Procedures). Among them 31 sequences share more than 97.1% identity and form another operational taxonomic unit (OTU-2, GenBank number MK073023). The nine other 16S rRNA sequences consisted of 6 OTUs that were all affiliated with *Alphaproteobacteria*. The high abundance of OTU-2 sequences detected is correlated with the dominant occurrence of BGC in this sample (Fig. 1). A Blast search in NCBI indicated that the two OTUs sharing 92% identity form a cluster that is affiliated with the phylum *Nitrospirae* (Fig. 2). They possess less than 97% identity with all 16S rRNA genes in the GenBank. The most closely related phylotypes are uncultured *Nitrospirae* MTBs clone MWB-1 (93% with OTU-1, 92% with OTU-2) and *Candidatus* *Magnetoovum mohavensis* LO-1 (94% with OTU-1, 92% with OTU-2). Therefore, the two OTUs represent a novel species of *Nitrospirae*.

Being one of the deep-branching MTB group, *Nitrospirae* MTB plays an important role when analyzing the origin and evolution of magnetosome biomineralization and magnetotaxis.

Lin et al. suggested that the magnetosome biomineralization and magnetotaxis was present before the separation of the *Nitrospirae* and *Proteobacteria* phyla, which was between ~3.38 and 3.21 Ga (Lin et al., 2017b). However, it remains unclear whether the acquisition of magnetosome island (MAI), that encodes essential proteins for magnetosome biosynthesis in the genome, occurred before or after the divergence between freshwater and marine *Nitrospirae* MTB. It is equally possible that the spherical cells of *Nitrospirae* phylum got MAI by horizontal gene transfer (HGT) then adapted into two different habitats or that they diverged into different habitats then acquired MAI separately. Future genomic comparison studies should provide answers to elucidate the evolutionary mechanism.

Corroboration of authenticity of giant cocci by fluorescence *in situ* hybridization

The BGC authenticity of OTU-2 was inferred from the fact that BGC is the most abundant morphotype in the analyzed samples. To corroborate this inference we designed a specific probe based on the conserved OTU-2 sequence and performed fluorescence *in situ* hybridization (FISH) analysis. As shown in Figure 3 A2, the universal bacterial probe hybridized with all the MTB, while the specific probe hybridized only with black-brown giant cocci. The WGC cells only hybridized with the bacteria universal probe and were used as a negative control. Therefore, the OTU-2 really corresponds to the abundant BGC morphotype.

We also confirmed the authenticity of the OTU-1 to the scarce WGC. A probe specific to the conserved sequences of OTU-1 was designed. We found that the specific probe hybridized only with the WGC morphotype while the universal probe hybridized with all bacterial cells including BGC that served as the negative control (Fig. 3, B1). After FISH identification, the

same cover-slips were observed under scanning electron microscopy (SEM). As shown in B3 of Fig. 3, more than 8 bundles of multiple magnetosome chains aligned parallel to the long axis and close to the outer surface in the WGC cell. Correlative FISH-SEM analysis not only allowed identification of the OTU but also revealed characteristics of the magnetosome chains.

Magnetosome and element composition analyses

After corroborating their authenticities, we analyzed the magnetosome chains and element composition in BGC and WGC. Under DIC microscopy BGC appeared to contain brown granules inside the cells while WGC had a smooth appearance (Fig. 4A). Observations using High-angle annular dark-field scanning transmission electron microscopy (HAADF-STEM) revealed that the BGC cells were filled with electron dense granules rich in sulfur (Fig. 4B, top cell with white inclusions, and 4C in red color) in addition to the magnetosomes chains (Fig. 4C green color). Further energy-dispersive X-ray spectroscopy (XEDS) analysis confirmed that the granules were rich in sulfur, the most abundant element in BGC cells but barely detectable in WGC (Fig. 4, D2 versus D1). The magnetosomes in WGC were arranged in bundles of multiple chains as revealed by FISH-SEM analysis. We attempted to identify the magnetosome crystals inside of BGC and WGC cells by electron diffraction or high-resolution electron microscopy without success due to the thickness of the cells. Although we identified isolated bullet-shaped crystals as magnetites on grids, but it is uncertain if they were from BGC or WGC because environmental samples containing other MTBs were used. We performed STEM-XEDS elemental map (Fig. 4C) and XEDS spectrum

analyses (Fig. 4B, E1, E2, F1 and F2), and unambiguously showed that the crystals in BGC and WGC cells are iron-oxide minerals, which is consistent with the finding of only magnetite minerals in *Nitrospirae* MTB magnetosomes in the literature.

One essential difference between BGC and WGC is that BGC contain numerous sulfur-rich inclusions. Intracellular sulfur globules were also detected in *Nitrospirae* magnetotactic bacteria *Ca. Magnetobacterium bavaricum*, LO-1, MWB-1 and MYR-1 (Jogler et al., 2010; Li et al., 2010; Lefevre et al., 2011; Lin et al., 2012a; Li et al., 2016), and those in the *Ca. Magnetobacterium bavaricum* are present in the form of cyclo-octa sulfur (S₈) (Eder et al., 2014). Intracellular deposition of sulfur may serve as an electron donor or an electron acceptor to provide energy for bacterial growth (Cox et al., 2002). We observed that BGC became the dominant population during storage under laboratory conditions and were able to survive for more than ten months while maintaining vivacious magnetotactic motility. Therefore, it is plausible that the metabolism of intracellular sulfur-rich inclusions might contribute to the long-term survival of BGC under laboratory conditions.

In conclusion, we reported the discovery of two novel marine species of *Nitrospirae* MTB and their morphology and structural differences. Future genomic analysis would be helpful to understand the evolution and adaptation mechanisms of *Nitrospirae* magnetotactic bacteria to freshwater and marine environments.

Experimental Procedures

Sediment collection and MTB enrichment

Sediment samples were collected from Six-Four les Plages (43°07'N, 5°79'E) in the

Mediterranean Sea, Southern France. The site was characterized by a salinity of ~35 ppt, pH of 7.5 and a temperature of 19°C at the time of sampling. Collected sediments were quickly transferred to 1000 ml plastic bottles, covered with approximately 600 ml of *in situ* seawater, transported to the laboratory, and incubated at room temperature without disturbance.

Black-brown giant cocci cells (BGC) in the sediment were magnetically enriched using the race-track method (Wolfe et al., 1987; Zhu et al., 2010). The 16S rRNA genes were directly amplified from the collections. White giant cocci cells (WGC) were further collected by magnetic micro-sorting using a Transfer Man ONM-2D micromanipulator and an IM-9B CellTram Oil manual hydraulic pressure-control system installed on an IX51 microscope equipped with a 40 × LD objective (Zhang et al., 2014). WGC cells were directly transferred to 3µl of PBS of the REPLI-g Single Cell Kit (QIAGEN). The PBS-suspended sample was frozen and thawed repeatedly, and the whole genome amplification (WGA) reaction was carried out using the REPLI-g Single Cell Kit (Chen et al., 2015). The 16S rRNA gene sequences were obtained as previously described by Chen et al. (Chen et al., 2015).

Phylogenetic analysis

The 16S rDNA sequences were aligned using CLUSTAL W multiple alignment software. Sequence identities were calculated with the BIOEDIT software. The phylogenetic tree was constructed via the neighbor-joining method using MEGA VERSION (Kumar et al., 2016), and boot-strap values were calculated from a 1000 replicates.

Optical microscopy

The morphologies and motilities of the enriched MTB cells were examined under DIC microscopy (Olympus BX51 equipped with a DP71 camera; Olympus) using the hanging-drop method in an applied magnetic field. The ratio between BGC and WGC was estimated by counting the two morphotypes of cells at the edge of the droplets as shown in Supporting Information Figure S1. To analyze the sensitivity to the osmolality the collected cell suspensions were diluted with deionized water before observing their magnetotactic behavior in the droplets. The trajectories of the cells were obtained using the dark-field mode with an increased exposure time. To assess auto-fluorescence, fluorescence mirror units were used to expose the cells to green light (510-550 nm), blue light (450-480 nm), violet light (400-410 nm) and UV light (330-385 nm) with the corresponding filter sets of Olympus.

FISH analyses

BGC-specific probe (5'-ATACTTAATCTTCCGCAGACC-3') and WGC-specific probes (5'-GGTTCCACACCGAAGCATT-3') were designed according to the conserved 16S rRNA gene sequences of OTU-2 (GenBank number MK073023) and OTU-1 (GenBank number MK203828), respectively and evaluated for their specificity by using the online probe evaluation tools Ribosomal Database Project (RDP) <https://rdp.cme.msu.edu/probematch/search.jsp> (Cole et al., 2009; Cole et al., 2014). The BGC- and WGC-specific probes were synthesized and fluorescently labeled with the hydrophilic sulfoindocyanine dye Cy3 at the 5' end. The universal bacterial probe EUB338 (5'-GCTGCCTCCCGTAGGAGT-3') was fluorescently labeled with fluorescein phosphoramidite 200 FAM at the 5' end and used as a positive control for FISH (Amann et al.,

1990).

Correlative FISH-SEM analyses

The FISH and correlative FISH-SEM analyses were carried out as described by Li et al. (Li et al., 2017). The same FISH coverslips of WGC were treated for SEM observations using a Zeiss Ultra-55 field-emission gun SEM (Carl Zeiss) operating at 5 kV (Li et al., 2017).

HAADF-STEM elemental composition analysis

Magnetosome biomineralization was investigated using a JEM-2100F microscope (JEOLLtd) operating at 200 kV equipped with a field emission gun, an ultra-high resolution (UHR) pole piece and a Gatan energy filter (GIF2001) system (Gatan, Inc.). HAADF-STEM was used for Z-contrast imaging. Chemical compositional analysis was performed by EDXS elemental mapping in the HAADF-STEM mode. By using this method, nanometer-scale spatial distribution of most elements can be determined, and the relative abundance of each element at a certain location can be semi-quantitatively calculated (Li et al., 2017).

Acknowledgments

We are grateful to H. Zhang and F.X. Wang for their assistance in the coordinated FISH-SEM experiments. X.X. Qian is grateful for her fellowship provided by the Chinese Scholarship Council. This work was supported by grants 41776131 and U1706208 from NSFC, grants 2016YFC0302502_02 and 2018YFC0309904 from National Key R&D Program of China, a grant XDB06010203 from the Strategic Priority Research Program, by a

grant from the Excellence Initiative of Aix-Marseille University - A*Midex, a French “Investissements d’Avenir” programme, and grant from CNRS for LIA-MagMC. The FISH-SEM experiments were performed at the IGGCAS, Beijing, China and were supported by a NSFC grant 41522402.

Conflict of interest

The authors declare that they have no conflict of interest

References

- Amann, R.L., Krumholz, L., and Stahl, D.A. (1990) Fluorescent-oligonucleotide probing of whole cells for determinative, phylogenetic, and environmental studies in microbiology. *J bacteriol* **172**: 762-770.
- Bazylinski, D.A., and Frankel, R.B. (2004) Magnetosome formation in prokaryotes. *Nat Rev Microbiol* **2**: 217.
- Blakemore, R. (1975) Magnetotactic bacteria. *Science* **190**: 377.
- Chen, Y.R., Zhang, R., Du, H.J., Pan, H.M., Zhang, W.Y., Zhou, K. et al. (2015) A novel species of ellipsoidal multicellular magnetotactic prokaryotes from Lake Yuehu in China. *Environ Microbiol* **17**: 637-647.
- Cole, J.R., Wang, Q., Fish, J.A., Chai, B., McGarrell, D.M., Sun, Y. et al. (2014) Ribosomal Database Project: data and tools for high throughput rRNA analysis. *Nucleic Acids Res* **42**: D633-D642.
- Cole, J.R., Wang, Q., Cardenas, E., Fish, J., Chai, B., Farris, R.J. et al. (2009) The Ribosomal

295 Database Project: improved alignments and new tools for rRNA analysis. *Nucleic Acids Res*
296 **37**: D141-D145.

297 Cox, B.L., Popa, R., Bazylnski, D.A., Lanoil, B., Douglas, S., Belz, A. et al. (2002)
298 Organization and Elemental Analysis of P-, S-, and Fe-rich Inclusions in a Population of
299 Freshwater Magnetococci. *Geomicrobiol J* **19**: 387-406.

300 Eder, S.H.K., Gigler, A.M., Hanzlik, M., and Winklhofer, M. (2014) Sub-micrometer-scale
301 mapping of magnetite crystals and sulfur globules in magnetotactic bacteria using confocal
302 Raman micro-spectrometry. *PloS one* **9**: e107356-e107356.

303 Flies, C.B., Peplies, J., and Schöler, D. (2005) Combined Approach for Characterization of
304 Uncultivated Magnetotactic Bacteria from Various Aquatic Environments. *Appl Environ*
305 *Microbiol* **71**: 2723-2731.

306 Jogler, C., Wanner, G., Kolinko, S., Niebler, M., Amann, R., Petersen, N. et al. (2011)
307 Conservation of proteobacterial magnetosome genes and structures in an uncultivated member
308 of the deep-branching Nitrospira phylum. *Proc Natl Acad Sci U S A* **108**: 1134-1139.

309 Jogler, C., Niebler, M., Lin, W., Kube, M., Wanner, G., Kolinko, S. et al. (2010)
310 Cultivation-independent characterization of 'Candidatus Magnetobacterium bavaricum' via
311 ultrastructural, geochemical, ecological and metagenomic methods. *Environ Microbiol* **12**:
312 2466-2478.

313 Kumar, S., Stecher, G., and Tamura, K. (2016) MEGA7: Molecular Evolutionary Genetics
314 Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* **33**: 1870-1874.

315 Lefèvre, C.T., Abreu, F., Schmidt, M.L., Lins, U., Frankel, R.B., Hedlund, B.P., and
316 Bazylnski, D.A. (2010) Moderately Thermophilic Magnetotactic Bacteria from Hot Springs

317 in Nevada. *Appl Environ Microbiol* **76**: 3740-3743.

318 Lefevre, C.T., and Bazylinski, D.A. (2013) Ecology, diversity, and evolution of magnetotactic
319 bacteria. *Microbiol Mol Biol Rev* **77**: 497-526.

320 Lefevre, C.T., Frankel, R.B., Abreu, F., Lins, U., and Bazylinski, D.A. (2011)
321 Culture-independent characterization of a novel, uncultivated magnetotactic member of the
322 Nitrospirae phylum. *Environ Microbiol* **13**: 538-549.

323 Li, J., Menguy, N., Gatel, C., Boureau, V., Snoeck, E., Leroy, E., and Patriarche, G. (2016)
324 Crystal growth of bullet-shaped magnetite in magnetotactic bacteria of the Nitrospirae phylum.
325 *J Royal Soc Interface* **12**: 103.

326 Li, J., Zhang, H., Menguy, N., Benzerara, K., Wang, F., Lin, X. et al. (2017) Single-Cell
327 Resolution of Uncultured Magnetotactic Bacteria via Fluorescence-Coupled Electron
328 Microscopy. *Appl Environ Microbiol* **83**: e00409-00417.

329 Li, J., Pan, Y., Liu, Q., Yu-Zhang, K., Menguy, N., Che, R. et al. (2010) Biomineralization,
330 crystallography and magnetic properties of bullet-shaped magnetite magnetosomes in giant
331 rod magnetotactic bacteria. *Earth Planet Sci Lett* **293**: 368-376.

332 Lin, W., Li, J., and Pan, Y. (2012a) Newly isolated but uncultivated magnetotactic bacterium
333 of the phylum Nitrospirae from Beijing, China. *Appl Environ Microbiol* **78**: 668-675.

334 Lin, W., Pan, Y., and Bazylinski, D.A. (2017a) Diversity and ecology of and biomineralization
335 by magnetotactic bacteria. *Environ Microbiol Rep* **9**: 345-356.

336 Lin, W., Wang, Y., Li, B., and Pan, Y. (2012b) A biogeographic distribution of magnetotactic
337 bacteria influenced by salinity. *ISME J* **6**: 475-479.

338 Lin, W., Zhang, W., Zhao, X., Roberts, A.P., Paterson, G.A., Bazylinski, D.A., and Pan, Y.

(2018) Genomic expansion of magnetotactic bacteria reveals an early common origin of magnetotaxis with lineage-specific evolution. *ISME J* **12**: 1508-1519.

Lin, W., Deng, A., Wang, Z., Li, Y., Wen, T., Wu, L.-F. et al. (2014) Genomic insights into the uncultured genus ‘Candidatus Magnetobacterium’ in the phylum Nitrospirae. *ISME J* **8**: 2463.

Lin, W., Paterson, G.A., Zhu, Q., Wang, Y., Kopylova, E., Li, Y. et al. (2017b) Origin of microbial biomineralization and magnetotaxis during the Archean. *Proc Natl Acad Sci* **114**: 2171-2176.

Spring, S., Amann, R.I., Ludwig, W., Schleifer, K., van Gernerden, H., and Petersen, N. (1993) Dominating Role of an Unusual Magnetotactic Bacterium in the Microaerobic Zone of a Freshwater Sediment. *Appl Environ Microbiol* **59**: 2397-2403.

Uebe, R., and Schüler, D. (2016) Magnetosome biogenesis in magnetotactic bacteria. *Nat Rev Microbiol* **14**: 621.

Wolfe, R.S., Thauer, R.K., and Pfennig, N. (1987) A ‘capillary racetrack’ method for isolation of magnetotactic bacteria. *FEMS Microbiol Lett* **45**: 31-35.

Zhang, R., Chen, Y.R., Du, H.J., Zhang, W.Y., Pan, H.M., Xiao, T., and Wu, L.F. (2014) Characterization and phylogenetic identification of a species of spherical multicellular magnetotactic prokaryotes that produces both magnetite and greigite crystals. *Res Microbiol* **165**: 481-489.

Zhu, K., Pan, H., Li, J., Yu-Zhang, K., Zhang, S.-D., Zhang, W. et al. (2010) Isolation and characterization of a marine magnetotactic spirillum axenic culture QH-2 from an intertidal zone of the China Sea. *Res Microbiol* **161**: 276-283.

Figure legends

Figure 1. Morphology and velocity of black-brown giant cocci (BGC) and ivory-white giant cocci (WGC).

(A) Differential interference contrast (DIC) micrograph shows magnetotaxis and accumulation of BGC (indicated by black arrows) and WGC (white arrows) at the north edge in a droplet. The big black arrow B indicates the direction of the magnetic field. (B) DIC micrograph shows the rough surface of BGC and smooth surface of WGC. (C) DIC micrograph shows both BGC

and WGC under low salinity conditions. (C1) Cell inclusions released from BGC. (C2) Shrunken shapes of WGC. (D) DIC micrograph (D1) and fluorescence micrograph (D1-D5) shows auto-fluorescence of BGC and WGC excited by light at different wavelengths. (D1) BGC and WGC under white light. (D2) BGC emitted red fluorescence while WGC had no fluorescence emission when exposed to green light. (D3) BGC emitted stronger yellow fluorescence than WGC under blue light. (D4) BGC and WGC emitted similar green fluorescence under violet light. (D5) WGC emitted stronger blue fluorescence than BGC under ultraviolet light. (E) Swim tracks and velocities of WGC (n=54) and BGC (n=52). The scale bars: 5µm in A, C, D. 10µm in B and E.

Figure 2. Phylogenetic tree of 16S rRNA gene sequences from *Nitrospirae* MTB.

All currently characterized *Nitrospirae* MTB and some typical *Nitrospirae* non-MTBs were used to construct the phylogenetic tree. The *Nitrospirae* MTB from seawater clustered to closely related freshwater *Nitrospirae* MTB. Bootstrap values (higher than 50) at nodes are

given as percentages of 1000 replicates. GenBank accession numbers of the sequences are given in parentheses. Scale bar represents 2% sequence divergence.

Figure 3. FISH identification of magnetotactic giant cocci.

Optical micrograph (A1) and fluorescence micrograph (A2) show BGC cells that were *in situ* hybridized (FISH) with the 5'-FAM-labeled universal bacterial probe EUB338 and the 5'-Cy3-labeled BGC-specific probe. Green color indicates bacteria hybridized only with the universal bacterial probe and yellow color shows the cell hybridized with both the universal bacterial probe and the BGC-specific probe. (B) Correlative FISH - SEM microscopic analysis of the WGC. (B1) In fluorescence micrograph, green color indicates bacteria hybridized only with universal bacterial probe EUB338 and yellow color shows the cell hybridized with both the universal bacterial probe and the 5'-Cy3-labeled WGC-specific probe. (B2) Correlated

396 SEM image of image B1. (B3) Correlated SEM image of the same red frame field in image B2
397 revealed the multiple magnetosome chains and bullet-shape magnetosome morphology of
398 WGS. The scale bars: 5 μ m in A1, A2, B1, B2. 2 μ m in B3.

399 **Figure 4. Element composition of BGC and WGC**

400

401 A) DIC image showed overall morphology difference between BGC (top cell) and WGC

402 (bottom cell). (B) STEM-HAADF image of BGC and WGC revealed multiple bundles of

magnetosome chains in both and granules filled in BGC, dashed rectangle is related to the area shown in E1. (C) STEM-XEDS elemental map of BGC and WBC revealing the presence of sulfur (red) and iron (green) overlaid with granules in BGC and iron-oxide crystals of magnetosomes in both BGC and WGC. (D1) XEDS spectrum corresponding to WGC showing that carbon is the most abundant element. (D2) XEDS spectrum corresponding to BGC showing that sulfur is the most abundant element. (E1) STEM-HAADF image related to the selected area in (A) and corresponding elemental map (E2). (F1) and (F2) represent XEDS spectra of the magnetosome chains in the selected areas #1 and #2 (in E2) in WBC and BGC cells, respectively. These spectra indicate the iron-oxide nature of magnetosome crystals. The scale bars: 5 μm in A, 1 μm in B and C, 0.5 μm in E1.