

Design of New Probes for Oxidized Amino Acids Localization

Mathieu Esgulian, Luc Camoin, Mathieu Cassien, Yves Toiron, Sylvia Pietri,
Sophie Thétiot-Laurent

► **To cite this version:**

Mathieu Esgulian, Luc Camoin, Mathieu Cassien, Yves Toiron, Sylvia Pietri, et al.. Design of New Probes for Oxidized Amino Acids Localization. Proceedings, MDPI, 2019, 22 (1), pp.39. 10.3390/proceedings2019022039 . hal-02393325

HAL Id: hal-02393325

<https://hal-amu.archives-ouvertes.fr/hal-02393325>

Submitted on 5 Dec 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Abstract

Design of New Probes for Oxidized Amino Acids Localization [†]

Mathieu Esgulian ^{1,*}, Luc Camoin ², Mathieu Cassien ¹, Yves Toiron ², Sylvia Pietri ¹ and Sophie Thétiot-Laurent ^{1,*}

¹ Aix Marseille Université, CNRS UMR 7273, Institut de Chimie Radicalaire, Equipe Sonde Moléculaires en Biologie et Stress Oxydant, 13013 Marseille, France

² CRCM, Inserm, 13009 Marseille, France

* Correspondence: mathieu.esgulian@univ-amu.fr (M.E.); sophie.thetiot-laurent@univ-amu.fr (S.T.-L.)

[†] Presented at the 2nd Molecules Medicinal Chemistry Symposium (MMCS): Facing Novel Challenges in Drug Discovery, Barcelona, Spain, 15–17 May 2019.

Published: 8 August 2019

Keywords: protein carbonyls; probes; hydrazine; click chemistry

Protein carbonyls (PC) are oxidative damage observed in many diseases. Proteins are possibly the most immediate vehicle for inflicting oxidative damage on cells because they are often catalysts [1]. A fluorometric and UV-absorption method have been developed to quantify PC in blood and tissues samples by labeling with two hydrazines: 7-hydrazino-4-nitrobenzo-2,1,3-oxadiazole (NBDH) and dinitrophenylhydrazine (DNPH), respectively [2,3]. These methods are based on the selective hydrazone formation between the carbonyls group of oxidized protein to yield a strong fluorescent/UV-absorption adduct that is then quantified.

Here, we will describe our study on NBDH's and DNPH's derivatives to generate new PC-probes bearing an alkyne moiety. In a first step, the hydrazine moiety reacted specifically with protein carbonyls and in a second step, a click reaction was performed between the alkyne moiety and a cleavable resin [4] to yield a PC's enrichment. These probes have been explored on oxidized bovine serum albumin (OxBSA) for PC-labeling, and the possible sites of oxidation of isolated labelled PC will be studied by LC-MS and proteomics experiments.

References

1. Dalle-Donne, I.; Rossi, R.; Giustarini, D.; Milzani, A.; Colombo, R. Protein carbonyl groups as biomarkers of oxidative stress. *Clin. Chim. Acta* **2003**, *329*, 23–38.
2. Stocker, P.; Ricquebourg, E.; Vidal, N.; Villard, C.; Lafitte, D.; Sellami, L.; Pietri, S. Fluorimetric screening assay for protein carbonyl evaluation in biological samples. *Anal. Biochem.* **2015**, *482*, 55–61.
3. Vidal, N.; Cavaille, J.P.; Graziani, F.; Robin, M.; Ouari, O.; Stocker, P. High throughput assay for evaluation of reactive carbonyl scavenging capacity. *Redox Biology* **2014**, *2*, 590–598.
4. Sibbersen, C.; Lykke, L.; Gregersen, N.; Jørgensen, K.A.; Johannsen, M. A cleavable azide resin for direct click chemistry mediated enrichment of alkyne-labeled proteins. *Chem. Commun.* **2014**, *50*, 12098–12100.



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).