



**HAL**  
open science

## Phylogenetically based establishment of a dengue virus panel, representing all available genotypes, as a tool in dengue drug discovery

Franck Touret, Cecile Baronti, Olivia Goethals, Marnix van Loock, Xavier de Lamballerie, Gilles Quérat

### ► To cite this version:

Franck Touret, Cecile Baronti, Olivia Goethals, Marnix van Loock, Xavier de Lamballerie, et al.. Phylogenetically based establishment of a dengue virus panel, representing all available genotypes, as a tool in dengue drug discovery. *Antiviral Research*, Elsevier Masson, 2019, 168, pp.109-113. 10.1016/j.antiviral.2019.05.005 . hal-02617069

HAL Id: hal-02617069

<https://hal-amu.archives-ouvertes.fr/hal-02617069>

Submitted on 25 Oct 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial | 4.0 International License

# Phylogenetically based establishment of a dengue virus panel, representing all available genotypes, as a tool in dengue drug discovery

1

2

3 Franck Touret<sup>1\*</sup>, Cécile Baronti<sup>1</sup>, Olivia Goethals<sup>2</sup>, Marnix Van Loock<sup>2</sup>, Xavier de

4 Lamballerie<sup>1</sup> and Gilles Querat<sup>1</sup>

5 <sup>1</sup>: Unité des Virus Émergents (UVE : Aix-Marseille Univ – IRD 190 – Inserm 1207 – IHU  
6 Méditerranée Infection), Marseille, France

7 <sup>2</sup> : Janssen Infectious Diseases Discovery, Janssen Pharmaceutica NV, Turnhoutseweg 30,  
8 2340, Beerse, Belgium.

9 \*Corresponding author: [franck.touret@hotmail.fr](mailto:franck.touret@hotmail.fr)

10

## 11 **Abstract**

12 Dengue fever is the most widespread of the human arbovirus diseases, with approximately  
13 one third of the world's population at risk of infection. Dengue viruses are members of the  
14 genus *Flavivirus* (family *Flaviviridae*) and, antigenically, they separate as four closely related  
15 serotypes (1-4) that share 60 to 75 % amino acid homology. This genetic diversity  
16 complicates the process of antiviral drug discovery. Thus, currently no approved dengue-  
17 specific therapeutic treatments are available. With the aim of providing an efficient tool for  
18 dengue virus drug discovery, a collection of nineteen dengue viruses, representing the  
19 genotypic diversity within the four serotypes, was developed. After phylogenetic analysis of  
20 the full-length genomes, we selected relevant strains from the EVAg collection at Aix-  
21 Marseille University and completed the virus collection, using a reverse genetic system based  
22 on the infectious sub-genomic amplicons technique. Finally, we evaluated this dengue virus  
23 collection against three published dengue inhibitory compounds. NITD008, which targets the

24 highly conserved active site of the viral NS5 polymerase enzyme, exhibited similar antiviral  
25 potencies against each of the different dengue genotypes in the panel. Compounds targeting  
26 less conserved protein subdomains, such as the capsid inhibitor ST-148, or SDM25N, a  $\delta$   
27 opioid receptor antagonist which indirectly targets NS4B, exhibited larger differences in  
28 potency against the various genotypes of dengue viruses. These results illustrate the  
29 importance of a phylogenetically based dengue virus reference panel for dengue antiviral  
30 research. The collection developed in this study, which includes such representative dengue  
31 viruses, has been made available to the scientific community through the European Virus  
32 Archive to evaluate novel DENV antiviral candidates.

### 33 **Keywords**

34 Dengue virus; phylogenetic analysis; reverse genetic; drug discovery; virus panel; dengue inhibitors

### 35 **Short communication**

36 Dengue virus (DENV) is a major threat to human health, with approximately one third of the  
37 world's population at risk of being infected. DENV is the causative agent of dengue fever, as  
38 well as the more severe dengue haemorrhagic fever (DHF)(Messina et al., 2014) and dengue  
39 shock syndrome (DSS). It belongs to the genus *Flavivirus* (*Flaviviridae* family), which  
40 comprises other clinically important human pathogens, such as yellow fever virus, West Nile  
41 virus and the recently emerging Zika virus(Vasilakis and Weaver, 2017). DENV is an  
42 arthropod borne virus transmitted through the bite of infected mosquitoes from the genus  
43 *Aedes* (*Stegomyia*). Epidemiological transmission of DENV is confined to urban and peri-  
44 urban cycles for which *Aedes aegypti* and *Ae albopictus* mosquitoes, respectively, are the  
45 primary transmission vectors(Chen and Vasilakis, 2011). Dengue is a positive-sense single  
46 stranded RNA virus with a 10.7 kb genome encoding a single polyprotein which is post-  
47 translationally processed into three structural proteins, viz., capsid (C), pre-membrane (prM),

48 envelope (E) and seven non-structural proteins (NS1, NS2A, NS2B, NS3, NS4A, NS4B and  
49 NS5)(Gebhard et al., 2011). Four antigenically closely related serotypes of DENV (1-4)  
50 which share 60 to 75 % amino acid homology, have been identified(Guzman and Harris,  
51 2015). Within this serotype demarcation, the DENV are also grouped into genotypes, with  
52 varying terminology between authors(Carrillo-Valenzo et al., 2010; Weaver and Vasilakis,  
53 2009) (hereunder we refer to the grouping proposed by Weaver and Vasilakis(Weaver and  
54 Vasilakis, 2009)).

55 Hence, many of the DENV diagnostic tools do not readily distinguish between DENV  
56 serotypes. Moreover, co-circulation of different serotypes during DENV epidemics(Vilela et  
57 al., 2016) increases the complexity of virus identification. Added to these factors, antibody  
58 dependent enhancement of the disease i.e., when patients contract a heterotypic secondary  
59 DENV infection(Katzelnick et al., 2017) is a potential additional complication for effective  
60 treatment of patients. Consequently, scientists are faced with the challenge of developing  
61 Directly Active Antivirals (DAA) that can inhibit the entire spectrum of genetically diverse  
62 serotypes and/or genotypes of DENV. However, despite the tireless efforts to provide an  
63 antiviral therapy(Canard, 2012; Coutard et al., 2008; van Cleef et al., 2013; Yin et al., 2009),  
64 there are still no approved drugs on the market to treat dengue infections. At present, the  
65 treatments available are merely supportive(Kaptein and Neyts, 2016).

66 A major barrier to evaluating the activity spectrum of potential DENV-inhibitory molecules  
67 arises from the non-availability of a well-defined panel of viruses that specifically represents  
68 the genetic variability of all characterised DENV isolates. With the aim of providing a tool for  
69 DENV research, with which to assess the antiviral activity of potential inhibitory molecules,  
70 we have developed a collection of DENV with sequences that include representative  
71 genotypes from within the four DENV serotypes (figure 1). Wherever possible, we selected  
72 clinical strains with a limited number of passages in cell culture. Strains were selected from

73 either the European Virus Archive (EVA) collection(Romette et al., 2018), the French  
74 National Reference Centre for arboviruses (CNR), or the World Reference Center for  
75 Emerging Viruses and Arboviruses (WRCEVA). Viruses that could not be obtained but for  
76 which full length genome sequences were available, were re-created using the versatile  
77 infectious sub-genomic amplicons (ISA) reverse genetics technology(Aubry et al., 2015,  
78 2014).

79 In order to select representative genotypes, we collected dengue full-length genome sequences  
80 from the NCBI database and complemented this database with those of our, still unpublished,  
81 “in house” and CNR strains. We performed phylogenetic reconstructions with the maximum  
82 likelihood method to assign all available genome sequences to a genotype in a serotype  
83 (supplementary material Fig 1, 2, 3 and 4). Within each genotype, we focused on strains that  
84 were not subjected to extensive cell passage and were either available as biological isolates in  
85 virus collections or as full-length sequences in GenBank. Six dengue genotypes were  
86 available only as complete genome sequences in the NCBI database without any biological  
87 strain counterparts in referenced collections (DENV-1 genotype III, DENV-2 genotype  
88 sylvatic and Asian II, DENV-3 genotype V and DENV-4 genotype III and sylvatic). Two  
89 genotypes were not available at all because of incomplete genome sequence (DENV-1  
90 genotype II and DENV-3 genotype IV). To obtain the biological viruses from the completely  
91 sequenced strains, we designed reverse genetics systems based on the ISA technique(Atieh et  
92 al., 2016; Aubry et al., 2015, 2014) and generated synthetic overlapping DNA fragments that  
93 covered each of the entire genome, bordered by a CMV promoter on the 5’ end and a  
94 Ribozyme and poly-adenylation signal on the 3’ end. The overlapping fragments were co-  
95 transfected into a mix of human and hamster embryonic kidney cell lines (HEK 293 and  
96 BHK-21 purchased from the American Cell Culture Collection). This enabled us to recover  
97 the missing biological strains to complete the collection. The initial viral stocks were

98 amplified in Vero E6 cells and fully sequenced. All the DENV strains used, have been made  
99 available through the EVAg collection (<https://www.european-virus-archive.com/>).

100 Various specific dengue inhibitors that target several viral proteins involved in different  
101 replication steps, have been discovered. ST-148, an inhibitor targeting the capsid structural  
102 protein, has been reported to inhibit all DENV serotypes in cell culture, although with varying  
103 efficiency. This inhibitor also appears promising in the AG-129 mouse model when infected  
104 with a strain of DENV-2(Byrd et al., 2013). NITD008, an adenosine analogue inhibitor that  
105 targets the RNA-dependent RNA polymerase activity, was shown to be inhibitory against all  
106 dengue serotypes as well as other flaviviruses, including West Nile virus, yellow fever virus  
107 and tick-borne Powassan virus(Yin et al., 2009). SDM25N, a  $\delta$  opioid receptor antagonist, has  
108 been reported to target the NS4B protein, probably indirectly through a cellular factor. Thus  
109 far, it has only been shown to be active against a DENV-2 strain(van Cleef et al., 2013).

110 Based on the different mechanisms of action of the 3 compounds, their respective target and  
111 its associated sequence variability across the different genotypes, we hypothesize that the  
112 antiviral activity of the compounds might differ between all of the genotypes of DENV.  
113 Therefore, the antiviral activity of these three compounds was assessed using a single  
114 common protocol based on a viral RNA yield reduction assay(Delang et al., 2016). The assay  
115 did not depend on the cytopathogenic potential of the strain, thus allowing for the inclusion of  
116 any dengue strain in the panel tested. Because all these strains differed in their replication  
117 kinetics, prior to the assay, all DENV MOI and times of readout of the assay were calibrated  
118 **so that** the replication growth were still in the log growth curve at time of the collection of the  
119 supernatant. Although the maximum reduction of virus yield may depend of the specific strain  
120 and assay conditions, the half inhibitory doses (IC50s) are not expected to be affected in these  
121 settings and will depend only on the inhibitor efficiency. The compounds were assayed from  
122 10 to 0.004 $\mu$ M, with 3-fold step-dilution in triplicate. The amount of viral RNA in the

123 supernatant medium, sampled at pre-determined time in the growth cycle, was quantified by  
124 qRT-PCR to determine the 50% maximal effective concentration (EC<sub>50</sub>) (Table 1).

125 The DENV strains of the collection showed similar sensitivity towards the nucleoside  
126 analogue inhibitor NITD008 with EC<sub>50</sub>'s ranging from 0.2µM to 2.8µM, which is in  
127 accordance with previously published results (Xie et al., 2015).

128 The capsid inhibitor ST-148 inhibited all DENV-2 genotypes with EC<sub>50</sub>'s ranging from 0.25  
129 to 1.1µM. However, only one genotype of DENV-1 (DENV-1 GIII at 0.5µM), and one of  
130 DENV-4 (DENV-4 GIII at 0.3µM), were inhibited by this compound. Finally, no activity was  
131 observed against our DENV-3 genotypes, with all EC<sub>50</sub>'s > 10µM. Although Byrd and co-  
132 workers(Byrd et al., 2013) found that the DENV-2 serotype was the most sensitive serotype to  
133 this capsid inhibitor and showed up to two log of variability in the inhibition against other  
134 serotypes, they did not fully evaluate the variation in susceptibility to other serotypes  
135 sufficiently comprehensively to draw conclusions. In their study, they associated ST-148  
136 resistance to a Leucine at position 34 instead of a Serine in DENV-2. However, looking at  
137 capsid amino-acid alignment of all our DENV panel and their study's strains, and regardless  
138 of their sensibility to ST-148, all strains exhibited a Serine at position 34 except DENV-2  
139 from Trinidad (1751 TC 544), which presented a Proline at this position, as the Modoc virus  
140 that they reported to be sensitive to ST-148 (Byrd et al., 2013). Thus, if resistance for ST-148  
141 can arise from S34L mutations in some DENV-2 strains it is clear that it cannot be  
142 unequivocally associated to a Leucine in position 34 in other serotypes and genotypes. This  
143 suggests that other residues or domains in the capsid protein may be involved in the  
144 interaction.

145 SDM25N showed moderate efficacy, with EC<sub>50</sub>'s ranging from 1.7 – 7.7 µM against a large  
146 proportion of the DENV-2 genotype strains, and half of the DENV-1 genotypes. However, no

147 activity was observed against any of the DENV-3 and 4 genotypes, as EC<sub>50</sub> were all above 10  
148 µM. This result suggests that the binding affinity of NS4B to the hypothetical cellular factor  
149 targeted by SDM25N varies greatly among various DENV genotypes and/or that this cellular  
150 factor might be dispensable for efficient replication of some DENV genotypes.

151 Overall, the results demonstrate that compounds targeting highly conserved sites, exemplified  
152 by nucleoside analogue inhibitor NITD008 (targeting the active site of the polymerase), had a  
153 broader pan-serotypic activity, with similar EC<sub>50</sub>'s regardless of the DENV genotype. In  
154 contrast, compounds targeting less conserved proteins or protein subdomains, either directly  
155 (*e.g.* the capsid) or indirectly through an interaction with a host factor of the cell (*e.g.*  
156 SDM25N), exhibited larger differences in activity towards the various genotypes of DENV.

157 Importantly, these data illustrate the fact that a sound *in cellulo* evaluation of anti-dengue  
158 candidate molecules requires the use of a complete reference virus panel that enables  
159 estimates of the antiviral activity against each of the identified DENV genotypes to be  
160 obtained. Modern reverse genetics techniques have enabled us to develop such a  
161 representative collection, and it has been made available to the scientific community through  
162 the European Virus Archive collection (EVA). We believe that the availability of this new  
163 tool will enable the independent assessment of pan-serotypic activity of anti-dengue  
164 candidates in the future, fulfilling a critical requirement for a successful dengue antiviral  
165 small molecule.

## 166 **Funding**

167 This research was carried out under sponsorship of Janssen-Cilag S. A., a pharmaceutical  
168 company of Johnson & Johnson (Research agreement ICD#1041950) and EVAg European  
169 Union – Horizon 2020 program under grant agreement no. 653316; [http://www.european-](http://www.european-virusarchive.com)  
170 [virusarchive.com](http://www.european-virusarchive.com)).



171 **Contributions**

172 OG, MVL, GQ and XDL generated the idea of the panel. FT, XDL and GQ conceived the  
173 experiments. XDL proposed the study design. FT, CB, and GQ performed the experiments.  
174 FT and GQ analysed the results. FT and GQ wrote the paper. FT, CB, GQ, OG, MVL and  
175 XDL reviewed and edited the paper.

176 **Acknowledgments**

177 We would thank Dr Robert Tesh from UTMB (Texas) for providing the dengue 4 Genotype I  
178 strain G11337 from India. We thank Dr Raphaelle Klitting for her help and expertise  
179 regarding the dengue phylogeny. We would thank Pr E.A Gould and Dr Stéphane Priet for  
180 their careful reading of the manuscript. As well as Magali Gilles, and Fiona Baudino from the  
181 UMR UVE (Marseille, France) for excellent technical help. We also thank Geraldine  
182 Piorkowski and Karine Barthelemy from UMR UVE (Marseille, France) for the sequencing.

183 **Bibliography**

- 184 Atieh, T., Baronti, C., de Lamballerie, X., Nougairède, A., 2016. Simple reverse genetics  
185 systems for Asian and African Zika viruses. *Sci. Rep.* 6, 39384.  
186 <https://doi.org/10.1038/srep39384>
- 187 Aubry, F., Nougairède, A., de Fabritus, L., Piorkowski, G., Gould, E.A., de Lamballerie, X.,  
188 2015. “ISA-Lation” of Single-Stranded Positive-Sense RNA Viruses from Non-  
189 Infectious Clinical/Animal Samples. *PloS One* 10, e0138703.  
190 <https://doi.org/10.1371/journal.pone.0138703>
- 191 Aubry, F., Nougairède, A., de Fabritus, L., Querat, G., Gould, E.A., de Lamballerie, X., 2014.  
192 Single-stranded positive-sense RNA viruses generated in days using infectious  
193 subgenomic amplicons. *J. Gen. Virol.* 95, 2462–2467.  
194 <https://doi.org/10.1099/vir.0.068023-0>
- 195 Byrd, C.M., Dai, D., Grosenbach, D.W., Berhanu, A., Jones, K.F., Cardwell, K.B., Schneider,  
196 C., Wineinger, K.A., Page, J.M., Harver, C., Stavale, E., Tyavanagimatt, S., Stone,  
197 M.A., Bartenschlager, R., Scaturro, P., Hrubby, D.E., Jordan, R., 2013. A Novel  
198 Inhibitor of Dengue Virus Replication That Targets the Capsid Protein. *Antimicrob.*  
199 *Agents Chemother.* 57, 15–25. <https://doi.org/10.1128/AAC.01429-12>
- 200 Canard, B., 2012. Antiviral research and development against dengue virus. Available  
201 Accessed 9.
- 202 Carrillo-Valenzo, E., Danis-Lozano, R., Velasco-Hernández, J.X., Sánchez-Burgos, G.,  
203 Alpuche, C., López, I., Rosales, C., Baronti, C., de Lamballerie, X., Holmes, E.C.,

204 Ramos-Castañeda, J., 2010. Evolution of dengue virus in Mexico is characterized by  
205 frequent lineage replacement. *Arch. Virol.* 155, 1401–1412.  
206 <https://doi.org/10.1007/s00705-010-0721-1>

207 Chen, R., Vasilakis, N., 2011. Dengue — Quo tu et quo vadis? *Viruses* 3, 1562–1608.  
208 <https://doi.org/10.3390/v3091562>

209 Coutard, B., Gorbalenya, A.E., Snijder, E.J., Leontovich, A.M., Poupon, A., De Lamballerie,  
210 X., Charrel, R., Gould, E.A., Gunther, S., Norder, H., Klempa, B., Bourhy, H.,  
211 Rohayem, J., L’hermite, E., Nordlund, P., Stuart, D.I., Owens, R.J., Grimes, J.M.,  
212 Tucker, P.A., Bolognesi, M., Mattevi, A., Coll, M., Jones, T.A., Åqvist, J., Unge, T.,  
213 Hilgenfeld, R., Bricogne, G., Neyts, J., La Colla, P., Puerstinger, G., Gonzalez, J.P.,  
214 Leroy, E., Cambillau, C., Romette, J.L., Canard, B., 2008. The VIZIER project:  
215 Preparedness against pathogenic RNA viruses. *Antiviral Res., Special Issue:*  
216 *Treatment of highly pathogenic RNA viral infections* 78, 37–46.  
217 <https://doi.org/10.1016/j.antiviral.2007.10.013>

218 Delang, L., Li, C., Tas, A., Quérat, G., Albulescu, I.C., De Burghgraeve, T., Guerrero, N.A.S.,  
219 Gigante, A., Piorkowski, G., Decroly, E., Jochmans, D., Canard, B., Snijder, E.J.,  
220 Pérez-Pérez, M.J., van Hemert, M.J., Coutard, B., Leyssen, P., Neyts, J., 2016. The  
221 viral capping enzyme nsP1: a novel target for the inhibition of chikungunya virus  
222 infection. *Sci. Rep.* 6, 31819. <https://doi.org/10.1038/srep31819>

223 Gebhard, L.G., Filomatori, C.V., Gamarnik, A.V., 2011. Functional RNA Elements in the  
224 Dengue Virus Genome. *Viruses* 3, 1739–1756. <https://doi.org/10.3390/v3091739>

225 Guzman, M.G., Harris, E., 2015. Dengue. *The Lancet* 385, 453–465.  
226 [https://doi.org/10.1016/S0140-6736\(14\)60572-9](https://doi.org/10.1016/S0140-6736(14)60572-9)

227 Kaptein, S.J., Neyts, J., 2016. Towards antiviral therapies for treating dengue virus infections.  
228 *Curr. Opin. Pharmacol.* 30, 1–7. <https://doi.org/10.1016/j.coph.2016.06.002>

229 Katzelnick, L.C., Gresh, L., Halloran, M.E., Mercado, J.C., Kuan, G., Gordon, A., Balmaseda,  
230 A., Harris, E., 2017. Antibody-dependent enhancement of severe dengue disease in  
231 humans. *Science* 358, 929–932. <https://doi.org/10.1126/science.aan6836>

232 Messina, J.P., Brady, O.J., Scott, T.W., Zou, C., Pigott, D.M., Duda, K.A., Bhatt, S.,  
233 Katzelnick, L., Howes, R.E., Battle, K.E., Simmons, C.P., Hay, S.I., 2014. Global  
234 spread of dengue virus types: mapping the 70 year history. *Trends Microbiol.* 22, 138–  
235 146. <https://doi.org/10.1016/j.tim.2013.12.011>

236 Romette, J.L., Prat, C.M., Gould, E.A., de Lamballerie, X., Charrel, R., Coutard, B., Fooks,  
237 A.R., Bardsley, M., Carroll, M., Drosten, C., Drexler, J.F., Günther, S., Klempa, B.,  
238 Pinschewer, D., Klimkait, T., Avsic-Zupanc, T., Capobianchi, M.R., Dicaro, A.,  
239 Ippolito, G., Nitsche, A., Koopmans, M., Reusken, C., Gorbalenya, A., Raoul, H.,  
240 Bourhy, H., Mettenleiter, T., Reiche, S., Batten, C., Sabeta, C., Paweska, J.T.,  
241 Eropkin, M., Zverev, V., Hu, Z., Mac Cullough, S., Mirazimi, A., Pradel, F., Lieutaud,  
242 P., 2018. The European Virus Archive goes global: A growing resource for research.  
243 *Antiviral Res.* <https://doi.org/10.1016/j.antiviral.2018.07.017>

244 van Cleef, K.W.R., Overheul, G.J., Thomassen, M.C., Kaptein, S.J.F., Davidson, A.D.,  
245 Jacobs, M., Neyts, J., van Kuppeveld, F.J.M., van Rij, R.P., 2013. Identification of a  
246 new dengue virus inhibitor that targets the viral NS4B protein and restricts genomic  
247 RNA replication. *Antiviral Res.* 99, 165–171.  
248 <https://doi.org/10.1016/j.antiviral.2013.05.011>

249 Vasilakis, N., Weaver, S.C., 2017. Flavivirus transmission focusing on Zika. *Curr. Opin.*  
250 *Virol., Emerging viruses: intraspecies transmission • Viral immunology* 22, 30–35.  
251 <https://doi.org/10.1016/j.coviro.2016.11.007>



252 Vilela, A.P.P., Miranda, D.P.J., Andrade, E.H.P., Abrahão, J.S., Araújo, V.E.M., Zibaoui,  
253 H.M., Oliveira, J.G., Rosa, J.C.C., Figueiredo, L.B., Kroon, E.G., Ferreira, P.C.P.,

254 2016. Spatial–Temporal Co-Circulation of Dengue Virus 1, 2, 3, and 4 Associated  
255 with Coinfection Cases in a Hyperendemic Area of Brazil: A 4-Week Survey. *Am. J.*  
256 *Trop. Med. Hyg.* 94, 1080–1084. <https://doi.org/10.4269/ajtmh.15-0892>  
257 Weaver, S.C., Vasilakis, N., 2009. Molecular evolution of dengue viruses: Contributions of  
258 phylogenetics to understanding the history and epidemiology of the preeminent  
259 arboviral disease. *Infect. Genet. Evol.* 9, 523–540.  
260 <https://doi.org/10.1016/j.meegid.2009.02.003>  
261 Xie, X., Zou, J., Wang, Q.-Y., Shi, P.-Y., 2015. Targeting dengue virus NS4B protein for  
262 drug discovery. *Antiviral Res.* 118, 39–45.  
263 <https://doi.org/10.1016/j.antiviral.2015.03.007>  
264 Yin, Z., Chen, Y.-L., Schul, W., Wang, Q.-Y., Gu, F., Duraiswamy, J., Kondreddi, R.R.,  
265 Niyomrattanakit, P., Lakshminarayana, S.B., Goh, A., Xu, H.Y., Liu, W., Liu, B.,  
266 Lim, J.Y.H., Ng, C.Y., Qing, M., Lim, C.C., Yip, A., Wang, G., Chan, W.L., Tan,  
267 H.P., Lin, K., Zhang, B., Zou, G., Bernard, K.A., Garrett, C., Beltz, K., Dong, M.,  
268 Weaver, M., He, H., Pichota, A., Dartois, V., Keller, T.H., Shi, P.-Y., 2009. An  
269 adenosine nucleoside inhibitor of dengue virus. *Proc. Natl. Acad. Sci. U. S. A.* 106,  
270 20435–20439. <https://doi.org/10.1073/pnas.0907010106>  
271

272 **Table 1 Dengue virus collection-susceptibility to three antiviral compounds assessed by yield**  
 273 **reduction assay.** Anti-capsid ST-148, nucleoside analogue NITD008 and  $\delta$  opioid receptor antagonist  
 274 SDM25N were independently tested twice, with 3 replicates per experiment, against the dengue  
 275 collection from 10 $\mu$ M to 0.005 $\mu$ M. AA: Asian American, A: American, C: cosmopolitan.

	Virus	Genotype	ST-148	NITD008	SDM25N
			EC50 ( $\mu$ M)	EC50 ( $\mu$ M)	EC50 ( $\mu$ M)
<b>Dengue 1</b>	D1/H/IMTSSA/98/606 <b>Djibouti</b>	<b>I</b>	>10	0,9 $\pm$ 0,1	>10
	JKT 1186 TYP 949 <b>Indonesia</b>	<b>IV</b>	>10	0,3 $\pm$ 0,03	5,5 $\pm$ 3,67
	CNR_25329 <b>France</b>	<b>V</b>	>10	2,7 $\pm$ 4	7,4 $\pm$ 0,04
	P72-1244 <b>Malaysia</b>	<b>III</b>	3 $\pm$ 0,5	0,9 $\pm$ 0,2	>10
<b>Dengue 2</b>	H/IMTSSA-MART/98-703 <b>France</b>	<b>AA</b>	0,8 $\pm$ 0,5	0,9 $\pm$ 0,3	2,9 $\pm$ 0,95
	_1751 TC 544 <b>Trinidad</b>	<b>A</b>	1 $\pm$ 0,7	0,3 $\pm$ 0,06	2,9 $\pm$ 0,01
	CNR_25679 <b>France</b>	<b>C</b>	1,1 $\pm$ 0,3	0,2 $\pm$ 0,07	1,9 $\pm$ 0,03
	CNR 25326 <b>Thailand</b>	<b>Asian I</b>	0,1 $\pm$ 0,03	0,9 $\pm$ 0,2	7,7 $\pm$ 0,04
	DENV-2/PG/BID-V2618/2008 <b>Papua New Guinea</b>	<b>Asian II</b>	0,2 $\pm$ 0,16	0,3 $\pm$ 0,5	4,1 $\pm$ 0,02
DKD811 <b>Malaysia</b>	<b>Sylvatic</b>	0,4 $\pm$ 0,18	0,4 $\pm$ 0,1	>10	
<b>Dengue 3</b>	DENV3/BR/D3LIMHO/2006 <b>Brazil</b>	<b>V</b>	>10	1 $\pm$ 0,09	>10
	4025 <b>Bolivia</b>	<b>III</b>	>10	1 $\pm$ 0,05	>10
	CNR 17046 <b>Malaysia</b>	<b>I</b>	>10	2,8 $\pm$ 0,3	>10
	CNR 15418 <b>Thailand</b>	<b>II</b>	>10	1,2 $\pm$ 0,3	>10
<b>Dengue 4</b>	G11337 <b>India</b>	<b>I</b>	>10	1,2 $\pm$ 0,03	>10
	Dakar HD 34460 <b>Senegal</b>	<b>IIb</b>	>10	0,9 $\pm$ 0,3	>10
	CNR_16861 <b>Malaysia</b>	<b>IIa</b>	>10	0,4 $\pm$ 0,01	>10
	ThD4_0476_97 <b>Thailand</b>	<b>III</b>	0,3 $\pm$ 0,08	0,2 $\pm$ 0,08	>10
	P75-514 <b>Malaysia</b>	<b>Sylvatic</b>	>10	1 $\pm$ 0,05	>10

276

**A****B**

**Figure 1 : Global serotype-representative DENV collection** A: Pipeline of the workflow employed for the virus collection. B: Maximum likelihood phylogenetic tree (GTR+G+I model with 500 bootstraps), based on the complete nucleic acid sequences of the virus collection. Strain information: DENV-1: Genotype I: (Djibouti, 1998, AF298808); Genotype III (Malaysia, 1972, EF457905.1); Genotype IV (Indonesia, 1977, EUO74031); Genotype V (France, 2014, MF004384); DENV-2: Genotype Asian-America (France Martinique, 1998, AF208496); Genotype American (Trinidad, 1953, EU073981.1); Genotype Cosmopolitan (France, 2014, MF004385); Genotype Asian 1 (Thailand, 2014, MH888331); Genotype Asian 2 (Papua New Guinea, 2008, FJ906959.1); Genotype Sylvatic (Malaysia, 2008, FJ467493.1); DENV-3: Genotype I (Malaysia, 2012, MF004386); Genotype II (Thailand, 2012, MH888332); Genotype III (Bolivia, 2011, MH888333); Genotype V (Brazil, 2006, JN697379.1); DENV-4: Genotype IIb (Senegal, 1981, MF004387); Genotype IIa (Malaysia, 2013, MH888334); Genotype III (Thailand, 1997, AY618988.1); Genotype Sylvatic (Malaysia, 1975, JF262779.1); Genotype I (INDIA, 1961, JF262783.1). Complete information relevant to the strains of the collection are more fully detailed in the supplemental material.