

Soil properties, grassland management, and landscape diversity drive the assembly of earthworm communities in temperate grasslands

Kevin Hoeffner, Mathieu Santonja, Cécile Monard, Lou Barbe, Mathilde Le Moing, Daniel Cluzeau

▶ **To cite this version:**

Kevin Hoeffner, Mathieu Santonja, Cécile Monard, Lou Barbe, Mathilde Le Moing, et al.. Soil properties, grassland management, and landscape diversity drive the assembly of earthworm communities in temperate grasslands. Pedosphere, 2021, 31 (3), pp.375-383. 10.1016/S1002-0160(20)60020-0 . hal-03105585

HAL Id: hal-03105585
https://amu.hal.science/hal-03105585

Submitted on 22 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Soil properties, grassland management, and landscape diversity drive the assembly of earthworm communities in temperate grasslands

Kevin HOEFFNER[ab] Mathieu SANTONJA[ac] Cécile MONARD[a] Lou BARBE[a] Mathilde LEMOING[a] Daniel CLUZEAU[a]

[a] University of Rennes, CNRS, ECOBIO UMR 6553, Rennes F-35000 (France)

[b] INRAE, AGROCAMPUS OUEST UMR SAS, Rennes F-35000 (France)

[c] Aix Marseille Université, Avignon Université, CNRS, IRD, IMBE, Marseille F-13397 (France)

Revised manuscript

Abstract

Earthworms are widespread soil organisms that contribute to a wide range of ecosystem services. As such, it is important to improve our knowledge, still scanty, of the factors that drive the assembly of earthworm communities. The aim of the present study was to conjointly evaluate the effects on the assembly of earthworm communities of i) soil properties (texture, organic matter content, and pH), ii) grassland management (grassland age, livestock unit, and type of fertilization), iii) landscape diversity (richness, diversity of surrounding habitats, and grassland plant diversity), and iv) presence of hedgerows. The study was conducted in temperate grasslands of Brittany, France. Earthworms were sampled in 24 grasslands and, in three of these grasslands, they were sampled near a hedgerow or near a ditch (control without a hedgerow). Soil properties explained the larger portion of the variation in the earthworm community parameters compared to grassland management or landscape diversity. The increase in soil organic matter content and pH were the most favorable factors for earthworm abundance and biomass, in particular for endogeic species. Regarding grassland management, the increase in the livestock unit was the most damaging factor for earthworm communities, in particular for the anecic earthworm biomass and endogeic species richness. Surprisingly, landscape diversity negatively affected the total earthworm abundance and epigeic earthworm biomass, but it was related to an increase in the epi-anecic species. At a finer scale, we also demonstrated that the presence of hedgerows surrounding grasslands enhanced earthworm species richness, especially within the epigeic and anecic ecological categories. This study highlights that the earthworm ecological categories respond specifically to environmental filters; further studies need to be conducted to elucidate the factors that drive the assembly of earthworm communities at this ecological category level. We recommend that policymakers should act on landscape management to favor earthworm diversity in order to improve the ecosystem services they drive.

INTRODUCTION

Earthworms are widespread soil organisms constituting the most important terrestrial biomass in temperate climate zones (Hole, 1981; Bar-On et al., 2018). They are usually classified into three main ecological categories depending on their physiology, morphology and behaviour: epigeic, anecic and endogeic species (Bouché, 1972, 1977). Briefly, epigeic earthworms live in and consume surface organic matter, anecic earthworms burrow vertical galleries to feed on a mixture of surface and soil organic matter, and endogeic earthworms burrow horizontal galleries to feed on soil organic matter (Bouché and Kretzschmar, 1974; Bouché, 1977; Jégou et al., 1998). Additionally, within the anecic earthworms, epi-anecic species feed preferentially on fresh surface organic matter (i.e. leaf litter) and are thereby distinguished from strict-anecic species that feed preferentially on humified organic matter already incorporated into the soil (Jégou et al., 1998; Larsen et al., 2016; Hoeffner et al., 2019). Depending on their ecological categories and associated feeding and burrowing behaviour, earthworms contribute to important ecosystem services provided by the soil such as nutrient cycling, water and climate regulation and primary production (Blouin et al., 2013; Bertrand et al., 2015). For example, van Groenigen et al. (2014) reported in a meta-analysis that an increase in crop production was observed in presence of earthworms, this increase ranging from 18% in presence of epigeic species up to 32% in presence of anecic species.

Earthworm communities are governed by different environmental filters, including biogeographical history, soil properties, land use and management as well as species interactions within the community (e.g. competition or facilitation; Lavelle, 1983; Curry, 2004; Decaëns et al., 2008). Previous studies focusing on the impact of soil properties on earthworm communities highlighted the key role played by soil pH, soil organic matter content and soil texture (Joschko et al., 2006; Lee, 1985; Decaëns et al., 2008). Other studies focused on the impact of land use on these earthworm communities (Boag et al., 1997; Decaëns et al., 2003, 2008; Cluzeau et al., 2012). For example, Ponge et al. (2013) reported that grasslands exhibited higher anecic earthworm abundance than croplands. In addition, Zaller and Arnone (1999) observed a positive correlation between the density and the biomass of earthworm communities and the plant species richness of grasslands, and in particular for endogeic species. Concerning land management, previous studies reported that ploughing (Chan, 2001; Briones and Schmidt, 2017), pesticide application (Pelosi et al., 2014) and low permanent cover (Vršic, 2011) negatively impact earthworm communities with a response intensity depending on the ecological category considered.

Other studies have been undertaken at a greater scale to evaluate the impact of landscape diversity on earthworm communities within croplands (Vanbergen et al., 2007; Lüscher et al., 2014; Frazão et al., 2017). For example, Flohre et al. (2011) observed that the earthworm species richness in croplands

Revised manuscript

decreased with the percentage of surrounding agricultural fields. Regulska and Kolaczkowska (2015) also reported that a cropland surrounded by a diverse landscape supported a higher earthworm diversity, density and biomass than the same type of cropland surrounded by a simpler landscape. However, the majority of the previous studies did not report effect of landscape diversity on earthworm communities of croplands and vineyards (Kovács-Hostyánszki et al., 2013; Frazão et al., 2017; Buchholz et al., 2017). Moreover, field margins of croplands were reported to exhibit higher abundance and diversity of earthworms than in the croplands itself but, surprisingly, these field margins were not reported to favor earthworm populations of these croplands (Smith et al., 2008; Roarty and Schmidt, 2013; Crittenden et al., 2015). Whether and how earthworms disperse within agricultural landscapes hence remains an unresolved issue.

A strong research effort has been done in the past decades to study the earthworm communities of croplands. Grasslands are the largest terrestrial ecosystem in the globe and produce many key ecosystem services, such as carbon storage, soil erosion mitigation or support for pollinators (Costanza et al., 1997; Conant and Paustian, 2002; Werling et al., 2014). The main objective of the present study was to conjointly evaluate the effects of soil properties, grassland management and landscape diversity on the assembly of grassland earthworm communities. Specifically, we hypothesized that the intensity of grassland management would negatively affect earthworm community parameters while the landscape diversity surrounding the grasslands would increase earthworm community parameters. The second objective was to evaluate the effect of hedgerows on these earthworm communities. By increasing the number of available niches, we hypothesized that the presence of a hedgerow in the grassland edge would increase earthworm community parameters (Tews et al., 2004). We conducted the study in an agricultural landscape of Brittany, France. Earthworms were sampled in 24 grasslands and, within three of them, they were oversampled near a hedgerow and near a ditch (control without hedgerow). Several parameters of the earthworm communities were evaluated including (i) the total abundance, total biomass, species richness and species diversity and (ii) the abundance, biomass and richness within each earthworm ecological category.

MATERIALS AND METHODS

Study site

The study site covers 10 km^2 and is a part of the Long Term Ecological Research (LTER) ‘‘Zone Atelier Armorique’’, located in Brittany, France (48°50’ N, -1°58’ W). The climate of the area is oceanic with a mean annual temperature of 11.7 °C, a mean annual rainfall of 815.0 mm and a mean annual relative humidity of 80.9 % (mean values over the period 2010-2016, data from Météo France). The main soil types encountered are Cambisols (IUSS Working Group, 2015) with high bedrock heterogeneity (granite, soft schist and aeolian loam). Moreover, the study area presents a substantial micro-topography, mainly due to

Revised manuscript

a high variability of landscape structures (e.g. hedges and ditches as field margins) with a hedge density ranging from 50 to 100 $m.ha^{-1}$ (Baudry et al., 2000; Thomas et al., 2016). Land use comprises mainly annual crops (corn, wheat, barley) and temporary or permanent grasslands, forest and unmanaged areas.

We used ground-truth aerial photos, which were taken every year since 1990, to construct a detailed land-use history for all grasslands, allowing us to precisely determine the age of each grassland. Based on this land-use history and after verification with grassland owners, we selected 24 grasslands ranging from 1 to 25 years since the last crop. Among them, three grasslands with an age gradient of 1-, 2- and 7-year-old were selected and oversampled from a hedgerow and a ditch at their surroundings to take into account a specific effect of hedgerow on soil properties (Marshall and Moonen, 2002; Walter et al., 2003).

Earthworm sampling and laboratory analyses

Earthworms were sampled in 2016 within the 24 grasslands at a 30 m distance from any grassland edge, and then in the 3 selected grasslands near a ditch and near a hedgerow. For the 3 selected grasslands, we standardized the sampling with 3 sampling points in order to consider 3 replicates with hedgerow (at 1, 5 and 10 m from the hedgerow) and 3 replicates without hedgerow (at 1, 5 and 10 m from the ditch).

Earthworms sampling followed the normalized protocol ISO 23 611-1, that was modified and validated during the RMQS BioDiv program (Cluzeau et al., 2012) combining chemical and physical extractions. Briefly, each earthworm sampling was characterised by a mean of three sub-sampling spaced of 10 m in line. Earthworm sub-sampling consisted of three waterings of 10 L with a gradient concentration of formaldehyde (0.25, 0.25 and 0.4%) on one square meter. After each watering, earthworms were collected for 15 min. Afterwards, a block of soil (25 × 25 × 20 cm, length × width × depth) was excavated within each sub-sampling area and earthworms were hand-sorted. The number of hand-sorted earthworms (HS) was multiplied by 16 to obtain an estimation per square meter. This number was then added to the number of earthworms counted with the formaldehyde extraction (F) to obtain the total number of earthworms per square meter (FHS): $FHS = F + (16 \times HS)$. Earthworms were fixed and preserved in formaldehyde solution (4%).

In the laboratory, each earthworm individual was counted, weighed, assigned to a stage of development (juvenile, sub-adult and adult), identified at the sub-species level and assigned to its ecological category: epigeic, anecic or endogeic (Bouché, 1972, 1977). Additionally, we distinguished within anecic earthworms, the epi-anecic (genus *Lumbricus*) from the strict-anecic earthworms (genus *Aporrectodea*) (Ferrière, 1980; Jégou et al., 1998). For juvenile individuals, identification was first limited to the genus and thereafter they were attributed a species name according to the proportions of sub-adults and adults present

Revised manuscript

of the same genus on each square meter. Earthworm diversity was analysed through three levels: total species richness, Shannon diversity index and species evenness index.

Environmental filters

We selected three environmental filters to explain earthworm community parameters: soil properties, grassland management and landscape diversity.

Soil properties were characterized by the soil texture, organic matter content and pH (water). Ten soil samples were randomly collected at 3 m around the earthworm sub-samplings using a cylindrical soil corer (5 cm diameter × 20 cm depth) in each grassland. These 10 soil samples were pooled and homogenized in order to consider one composite soil sample per grassland and sent to the analytical laboratory of LABOCEA (Combourg, France). Briefly, clay content ranged from 9.5% to 19.7%, sand content from 13.3% to 68.9%, organic matter content from 1.8% to 5.2% and soil pH from 5.5 to 6.7 (Supplementary Table S1).

Grassland management was assessed from interviews with farmers (Supplementary Table S1) and from ground-truth aerial photos. The grassland age ranged from 1 to 25 years since the last row-cropping using quite similar species sown (*Lolium perenne* and *Trifolium repens* or *pratensis*). In addition, livestock unit per hectare varied from 0 to 4.3. Fertilisation rate was declarative so we used only the distinction between organic and mineral input.

Landscape structure within 100 m radius around the sampled fields was classified into 9 habitats based on aerial photos (forest, grassland, crop, hedge, water, building, garden, asphalt area, road). The radius of 100 m was chosen to reflect the overall low mobility of earthworms (Bardgett et al., 2005; Eijsackers, 2010, 2011). Landscape diversity was characterized by two indexes: total richness of habitats within the radius and Shannon Diversity Index of habitats (hereafter called SHDI). Mapping and analysis were done using the softwares QGis 2.8.1 and FRAGSTATS 4.296. In addition, we characterized the plant community of the 24 grasslands in spring 2015 using 10 quadrats (1 × 1 m) evenly distributed in each grassland, characterizing for each plant species its covering percentage. Among the 24 grasslands selected, landscape richness varied from 1 to 7 habitats (maximum number of habitats has never been observed), SHDI from 0.1 to 1.6 and plant Shannon index (hereafter called Plant diversity) within grasslands from 1.2 to 3.2 (Supplementary Table S1).

Statistical analysis

We used multiple linear regression models to test the effects of soil properties (decomposed in clay, sand, organic matter contents and pH), grassland management (decomposed in grassland age, livestock unit

Revised manuscript

and fertilisation), and landscape diversity (decomposed in landscape richness, SHDI and plant diversity) on all earthworm community parameters (i.e. total abundance and biomass, total diversity indexes, ecological categories abundance and biomass). We constructed a full model comprising all environmental filters, and then we selected the significant environmental filters using a backward stepwise selection procedure that selects the best model using the AIC criterion (Crawley, 2012; stepAIC function of the "MASS" package). We also evaluated the variance inflation factor (VIF) of each variable selected by the previous procedure to test for multicollinearity among environmental filters. We removed all environmental filters that showed a VIF > 5, even if significant from the model. Data met the conditions of normality and homoscedasticity.

Second, within each of the three selected grasslands (i.e. 1-, 2- and 7-year-old), we compared earthworm communities with and without hedgerow (ditch) using the three sampling points per plot as replicates. We used separated *t*-tests within the three selected grasslands to assess the differences in earthworm abundance, earthworm biomass, and species richness according to the presence or absence of a hedgerow.

Statistical analyses were performed with the R software 3.2.3 (R. Core Team, 2017). Significance was evaluated in all cases at $P < 0.05$.

RESULTS

Impact of soil properties, grassland management and landscape diversity on earthworm communities

Over the 24 grasslands sampled, the average earthworm abundance and biomass were 517.0 ± 57 individual.m^{-2} and 219.4 ± 20 g.m^{-2}, respectively. The mean earthworm species richness was 10.8 ± 0.3. Eighteen species belonging to the three ecological categories were identified (Supplementary Table S2). *Allolobophora chlorotica* and *Aporrectodea caliginosa* were the most abundant species whereas *Eisenia tetraedra, Dendrobaena rubida and Octalasium lacteum* were present in one grassland only (Supplementary Table S2).

Higher soil organic matter content increased the total earthworm abundance ($F = 5.3$, $P = 0.033$, Table 1), the endogeic species abundance ($F = 5.7$, $P = 0.028$, Supplementary Table S3) and the endogeic species richness ($F = 5.4$, $P = 0.031$, Supplementary Table S4), while the endogeic species abundance was negatively correlated to the sand content ($F = 6.9$, $P = 0.017$, Supplementary Table S3). In addition the total earthworm abundance and biomass increased when soil pH was more alkaline ($F = 5.0$ and 6.8, $P < 0.05$, Fig. 1, Table 1) but no category-specific impact was observed with respect to pH variation.

The increase in livestock unit decreased total earthworm biomass ($F = 5.7$, $P = 0.028$, Table 1), and in particular the biomass of anecic species ($F = 9.6$, $P = 0.005$, Fig. 2a, Supplementary Table S5). However,

Revised manuscript

this negative effect was only confirmed for the biomass of epi-anecic species (F = 4.4, P = 0.049, Fig. 2b, Supplementary Table S5). The increase in livestock unit also decreased the earthworm species richness, the Shannon diversity index and the species evenness (F = 2.8 to 9.6, P < 0.05, Fig. 2c, Table 1), and in particular the endogeic species richness (F = 9.5, P = 0.006, Supplementary Table S4). Mineral fertilisation enhanced the epigeic species abundance and biomass compared to organic fertilisation (F = 6.6 and 8.6, P < 0.02, Supplementary Tables S3 and S5).

Landscape richness decreased the biomass of epigeic species (F = 4.9, P = 0.041, Supplementary Table S4) but enhanced the epi-anecic species richness (F = 6.6, P = 0.019, Supplementary Table S4). The increase of SHDI decreased the total earthworm abundance (F = 4.6, P = 0.047, Table 1). In addition, the increase in plant diversity was positively correlated to Shannon diversity index and species evenness (F = 5.0 and 4.8, P < 0.04, Table 1).

Interestingly, the abundance of strict-anecic species, their biomass and richness were not affected by any of the environmental filters measured (Supplementary Tables S3, S4 and S5).

Impact of hedgerow presence on earthworm communities

Over the 3 grasslands oversampled, earthworm abundance was higher in the 2-year-old grassland (834 ± 76 individuals.m^{-2}) compared to the 1-year-old (306 ± 32 individuals.m^{-2}) and 7-year-old grasslands (385 ± 32 individuals.m^{-2}). Earthworm species richness was higher in the 2- and 7-year-old grasslands (11.0 ± 0.4 and 10.2 ± 0.3, respectively) compared to the 1-year-old grassland (7.9 ± 0.4). Earthworm species composition was also strongly different between these three grasslands. For example, the presence of *Eisenia tetraedra* occurred only in the 2-year-old grassland and the presence of *Aporrectodea caliginosa meridionalis* occurred only in the 7-year-old grassland.

Earthworm species richness was 21.0% and 23.2 % higher with the presence of a hedgerow, compared to the presence of a ditch, in the grasslands of 1- and 2-year-old (t = 5.8 and 13.9, P < 0.03, Fig. 3a and b). It was however not affected in the 7-year-old grassland (t = 0.0, P = 0.85, Fig. 3c). The abundance of earthworms was not affected by the presence of hedgerows in the three selected grasslands (t= 0.0 to 0.03, P > 0.865).

Overall, except *Allolobophora icterica* and *Aporrectodea nocturna* that were more abundant with the presence of a hedgerow, the strict-anecic and endogeic species were evenly distributed between the plots with and without a hedgerow. The distribution of epi-anecic earthworm species was heterogeneous, but *Lumbricus rubellus rubellus* and *Lumbricus terrestris* were more often observed in presence of a hedgerow. The distribution of epigeic earthworm was species dependent: *Dendrobaena mammalis* occurrence was higher in presence of a hedgerow and *Eisenia tetraedra* was observed in presence of a hedgerow in the 2-

Revised manuscript

246 year-old grassland only. *Lumbricus castaneus* and *Lumbricus rubellus castanoïdes* occurrences were overall
247 similar between the plots, independent from the presence of a hedgerow.
248
249 DISCUSSION
250 In the present study, we clearly demonstrated that soil properties, grassland management and
251 landscape diversity conjointly affected the selected parameters of the earthworm communities. Our findings
252 hence contrast with those of Frazão et al. (2017) who reported that earthworm communities of the croplands
253 were impacted by agricultural practices only but neither by soil properties nor landscape diversity.
254 Contrary to previous studies that observed an effect of soil properties at the regional scale (Decaëns
255 et al., 2003; Vanbergen et al., 2007; Decaëns et al., 2008), here, by taking the earthworm ecological category
256 into account, we evidenced that soil properties impact on a finer scale (i.e. 10 km^2), the abundance, biomass
257 and richness of earthworm ecological categories. This result might be due to the strong spatial heterogeneity
258 of the soil properties in the studied region (Jamagne, 2011). In agreement with previous studies, we observed
259 that higher soil sand content decreased the total abundance of earthworms (Hendrix et al., 1992; Lapied et
260 al., 2009), which could be due to the low capacity of sandy soils to hold water, leading to an unfavorable
261 habitat for earthworms (Lee, 1985). In addition, the increase in soil pH was positively correlated to both
262 earthworm species richness (Joschko et al., 2006) and total abundance (Ma et al., 1990; McCallum et al.,
263 2016). Nonetheless, several reviews observed that earthworm preference to soil pH was species-dependent
264 due to their synecology (Bouché, 1972; Edwards and Lofty, 1977; Lee, 1985) but the underlying
265 mechanisms for pH preference are not fully understood yet. In line with their feeding behaviour that consists
266 in consuming mainly humified organic matter, endogeic earthworm communities were more abundant and
267 diversified in grasslands presenting high contents of soil organic matter (Bouché, 1977; Piearce, 1978;
268 Ferrière, 1980).
269 Regarding grassland management, increasing livestock unit was the most damaging factor for
270 earthworm communities as it decreased the total biomass, species richness, the Shannon diversity index and
271 the species evenness. This strong negative effect could be associated to the trampling at high stocking levels
272 that damages soil structure and thus adversely affect earthworm communities and burrows (Cluzeau et al.,
273 1992; Pietola et al., 2005; Chan and Barchia, 2007). Interestingly, earthworms' response to livestock unit
274 was almost entirely confined to the largest epi-anecic and endogeic species and only the earthworm biomass
275 was affected, contrary to their abundance, suggesting a decrease in the mean body size rather than in
276 individuals' number. Surprisingly, mineral fertilisation enhanced the abundance and biomass of epigeic
277 species, but this finding is nonetheless in line with some previous studies that reported an increase in
278 earthworm abundance in relation to N mineral fertilisation (Muldowney et al., 2003; King and Hutchinson,

Revised manuscript

2007; Curry et al., 2008). Mineral fertilisation would probably allow a better primary production leading to higher leaf litter inputs that constitute a source of refuge and food for earthworms. Further studies are needed, in grassland, to elaborate the different impacts of manure versus mineral fertilisation on earthworms. Overall, we observed that within grasslands, grazing pressure led to smaller and less-diversified earthworm communities.

We observed a negative effect of increasing landscape diversity (richness and Shannon Index) on the total abundance of earthworms and, to our knowledge, for the first time, the biomass of epigeic earthworms in grasslands. A negative correlation between the total abundance of earthworms and landscape diversity was also observed by Flohre et al., (2011) in croplands, and the authors hypothesized that landscape diversity increases the number of earthworm predators. Indeed, several studies highlighted that landscape diversity enhance the abundances of invertebrates, mammals and birds (Marshall and Moonen, 2002; Maudsley et al., 2002; Vickery et al., 2009) that are potential predators for earthworms (Granval and Aliaga, 1988; O'Brien et al., 2016). We can also hypothesize that the capacity of epigeic species to disperse is hindered by physical barriers (i.e. hedge or ditch) and different soil properties (shelter and litter availability) in neighboring habitats that nonetheless constitute landscape diversity. In contrast, the species richness of epi-anecic earthworm was enhanced by the landscape diversity. As epi-anecic earthworm species have a great mobility varying from 1.5 to 14 m. year^{-1} (Hoogerkamp et al., 1983; Eijsackers, 2011; Nuutinen et al., 2014) and the ability to burrow into the soil to protect themselves, higher landscape diversity around grasslands could enhance their areas of emigration. Endogeic earthworm species were not impacted by landscape diversity and were highly abundant in each grassland as previously reported (Lavelle, 1983; Decaëns et al., 2008). Overall, it is possible that low agricultural practices in grasslands, compared to croplands or vineyard, could increase the effect of the surrounding landscape diversity on earthworm communities (Roarty and Schmidt, 2013; Buchholz et al., 2017; Frazão et al., 2017).

In addition to the effect of landscape diversity, we highlighted the importance of hedgerows surrounding grasslands. Hedgerows especially acted in young grasslands (i.e. 1- and 2-year-old grassland), which is probably due to the increase earthworm species aggregation with the age of the grasslands (Richard et al., 2012). It is well known that hedgerows locally modify soil properties (i.e. soil moisture, temperature or organic matter content; Marshall and Moonen, 2002), and especially the amount and type of litter deposited at the soil surface (Walter et al., 2003). This litter input is a key factor for the development of earthworm communities (Lee, 1985; Edwards, 2004), and in particular for epigeic and epi-anecic species that have a diet mainly composed of fresh leaf litter (Bouché and Kretzschmar, 1974; Piearce, 1978; Ferrière, 1980). In field, earthworm communities living in grasslands surrounded by a hedgerow were richer in earthworm species compared to earthworm communities in grasslands surrounded by a ditch, especially for

Revised manuscript

epigeic and epi-anecic earthworm species. Thus, hedgerows presence could promote earthworm diversity in grasslands. Increasing epi-anecic earthworm diversity in grasslands landscape could have consequences on ecosystem services provided by these species. Hoeffner et al. (2018) observed that burrows' fungal communities were regulated by epi-anecic species identity, which could increase the diversity of the drilospheric microbiota and improve soil functioning. Besides, as it is difficult to monitor the earthworm diversity response to global change drivers, earthworm databases often concern surveys carried out at regional or national scales (Rutgers et al., 2009; Cluzeau et al., 2012; Cameron et al., 2016). A first predictive model on the abundance and diversity of earthworms was created by Rutgers et al. (2016) taking into account soil occupation and properties. Future predictive models could therefore take into account the landscape as an additional factor regulating these earthworm communities.

CONCLUSION

Our study clearly illustrated that earthworm communities in grasslands were affected by the three environmental filters considered: soil properties, grassland management and landscape diversity. Soil properties was the main environmental filter controlling earthworm communities. However, we also highlighted important effects of grassland management, for instance a strong decrease in abundance of earthworms with increasing livestock unit. We observed various effects of landscape diversity, such as a surprising overall decrease of earthworm abundance or a higher epi-anecic richness in diverse landscapes. Therefore, our findings demonstrated conjoint effects of various environmental filters as drivers of earthworm communities. Taken together, our results suggest a strong context dependency in the assembly rules of earthworm communities, despite the fact that these communities are well known to be ubiquitous and resilient.

ACKNOWLEDGEMENTS

We thank the LABOCEA (Combourg, France) for mineral soil analyses and Météo France for meteorological data. We also thank Sarah Guillocheau, Daniel Cylly, Albin Fertil, André Bastin, Valentin Tripet, Olivier Jambon, Stéphanie Llopis, Cendrine Mony and Vincent Jung for their help during field sampling. We sincerely thank Claudia Wiegand for helpful comments and improvement of the English. Kevin Hoeffner and Lou Barbe were supported by a French PhD grant from the French Ministère de la Recherche.

REFERENCES

Revised manuscript

Bardgett, R., Hopkins, D., Usher, M., 2005. Biological diversity and function in soils. Cambridge University Press, Cambridge.

Bar-On, Y.M., Phillips, R., Milo, R., 2018. The biomass distribution on Earth. Proceedings of the National Academy of Sciences 201711842.

Baudry, J., Burel, F., Thenail, C., Le Cœur, D., 2000. A holistic landscape ecological study of the interactions between farming activities and ecological patterns in Brittany, France. Landscape and Urban Planning 50, 119–128.

Bertrand, M., Barot, S., Blouin, M., Whalen, J., Oliveira, T. de, Roger-Estrade, J., 2015. Earthworm services for cropping systems. A review. Agronomy for Sustainable Development 35, 553–567.

Blouin, M., Hodson, M.E., Delgado, E.A., Baker, G., Brussaard, L., Butt, K.R., Dai, J., Dendooven, L., Peres, G., Tondoh, J.E., Cluzeau, D., Brun, J.-J., 2013. A review of earthworm impact on soil function and ecosystem services: Earthworm impact on ecosystem services. European Journal of Soil Science 64, 161–182.

Boag, B., Palmer, L.F., Neilson, R., Legg, R., Chambers, S.J., 1997. Distribution, prevalence and intensity of earthworm populations in arable land and grassland in Scotland. Annals of Applied Biology 130, 153–165.

Bouché, M.B., 1977. Strategies lombriciennes. Ecological Bulletins, Soil Organisms as Components of Ecosystems 25, 122–132.

Bouché, M.B., 1972. Lombriciens de France: écologie et systématique, INRA-Annales de Zoologie Ecologie Animale. ed. INRA, France.

Bouché, M.B., Kretzschmar, A., 1974. Fonctions des lombriciens II. Recherches méthodologiques pour l'analyse qualitative de la matière organique végétale ingéré (étude du peuplement de la station RCP-165/PBI). Revue d'Ecologie et de Biologie Du Sol 11, 127–139.

Briones, M.J.I., Schmidt, O., 2017. Conventional tillage decreases the abundance and biomass of earthworms and alters their community structure in a global meta-analysis. Global Change Biology 23, 4396–4419.

Buchholz, J., Querner, P., Paredes, D., Bauer, T., Strauss, P., Guernion, M., Scimia, J., Cluzeau, D., Burel, F., Kratschmer, S., Winter, S., Potthoff, M., Zaller, J.G., 2017. Soil biota in vineyards are more influenced by plants and soil quality than by tillage intensity or the surrounding landscape. Scientific Reports 7, 17445.

Butt, K.R., Frederickson, J., Morris, R.M., 1994. Effect of earthworm density on the growth and reproduction. Pedobiologia 38, 254–261.

Revised manuscript

376 Cameron, E.K., Vila, M., Cabeza, M., 2016. Global meta-analysis of the impacts of terrestrial invertebrate
377 invaders on species, communities and ecosystems. Global Ecology and Biogeography 25, 596–606.
378 Chan, K.Y., 2001. An overview of some tillage impacts on earthworm population abundance and diversity
379 - Implications for functioning in soils. Soil and Tillage Research 57, 179–191.
380 Chan, K.Y., Barchia, I., 2007. Soil compaction controls the abundance, biomass and distribution of
381 earthworms in a single dairy farm in south-eastern Australia. Soil and Tillage Research 94, 75–82.
382 Cluzeau, D., Binet, F., Vertes, F., Simon, J.C., Riviere, J.M., Trehen, P., 1992. Effects of intensive cattle
383 trampling on soil-plant-earthworms system in two grassland types. Soil Biology and Biochemistry
384 24, 1661–1665.
385 Cluzeau, D., Guernion, M., Chaussod, R., Martin-Laurent, F., Villenave, C., Cortet, J., Ruiz-Camacho, N.,
386 Pernin, C., Mateille, T., Philippot, L., Bellido, A., Rougé, L., Arrouays, D., Bispo, A., Pérès, G.,
387 2012. Integration of biodiversity in soil quality monitoring: Baselines for microbial and soil fauna
388 parameters for different land-use types. European Journal of Soil Biology, Bioindication in Soil
389 Ecosystems 49, 63–72.
390 Conant, R.T., Paustian, K., 2002. Potential soil carbon sequestration in overgrazed grassland ecosystems.
391 Global Biogeochemical Cycles 16, 90-1-90–9.
392 Costanza, R., d'Arge, R., Groot, R. de, Farber, S., Grasso, M., Hannon, B., Limburg, K., Naeem, S., O'Neill,
393 R.V., Paruelo, J., Raskin, R.G., Sutton, P., Belt, M. van den, 1997. The value of the world's
394 ecosystem services and natural capital. Nature 387, 253–260.
395 Crawley, M.J., 2012. The R Book, 2nd Edition. ed. Wiley & Sons, Chichester.
396 Crittenden, S.J., Huerta, E., de Goede, R.G.M., Pulleman, M.M., 2015. Earthworm assemblages as affected
397 by field margin strips and tillage intensity: An on-farm approach. European Journal of Soil Biology
398 66, 49–56.
399 Curry, J.P., 2004. Factors affecting the abundance of earthworms in soils, in: Edwards, C.A. (Ed.),
400 Earthworm Ecology. CRC Press, Boca Raton, pp. 91–113.
401 Curry, J.P., Doherty, P., Purvis, G., Schmidt, O., 2008. Relationships between earthworm populations and
402 management intensity in cattle-grazed pastures in Ireland. Applied Soil Ecology 39, 58–64.
403 Decaëns, T., Bureau, F., Margerie, P., 2003. Earthworm communities in a wet agricultural landscape of the
404 Seine Valley (Upper Normandy, France). Pedobiologia, The 7th International Symposium on
405 Earthworm Ecology 47, 479–489.
406 Decaëns, T., Margerie, P., Aubert, M., Hedde, M., Bureau, F., 2008. Assembly rules within earthworm
407 communities in North-Western France - A regional analysis. Applied Soil Ecology 39, 321–335.

Revised manuscript

Edwards, C.A., 2004. The importance of earthworms as key representatives of the soil fauna, in: Edwards, C.A. (Ed.), Earthworm Ecology. CRC Press, Boca Raton, pp. 3–11.

Edwards, C.A., Lofty, J.R., 1977. Biology of Earthworms, 2nd ed. Wiley & Sons, New York.

Eijsackers, H., 2011. Earthworms as colonizers of natural and cultivated soil environments. Applied Soil Ecology 50, 1–13.

Eijsackers, H., 2010. Earthworms as colonisers: Primary colonisation of contaminated land, and sediment and soil waste deposits. Science of The Total Environment 408, 1759–1769.

Ferrière, G., 1980. Fonctions des Lombriciens. VII. Une méthode d'analyse de la matière organique végétale ingérée. Pedobiologia 20, 263–273.

Flohre, A., Rudnick, M., Traser, G., Tscharntke, T., Eggers, T., 2011. Does soil biota benefit from organic farming in complex vs. simple landscapes? Agriculture, Ecosystems & Environment 141, 210–214.

Frazão, J., de Goede, R.G.M., Brussaard, L., Faber, J.H., Groot, J.C.J., Pulleman, M.M., 2017. Earthworm communities in arable fields and restored field margins, as related to management practices and surrounding landscape diversity. Agriculture, Ecosystems & Environment 248, 1–8.

Granval, P., Aliaga, R., 1988. Analyse critique des connaissances sur les prédateurs de lombriciens. Gibier Faune Sauvage 5, 71–94.

Groenigen, J.W. van, Lubbers, I.M., Vos, H.M.J., Brown, G.G., Deyn, G.B.D., Groenigen, K.J. van, 2014. Earthworms increase plant production: a meta-analysis. Scientific Reports 4, 6365.

Hartenstein, R., 1984. Rate of production and loss of earthworm biomass in relation to species and size. Soil Biology and Biochemistry 16, 643–649.

Hartenstein, R., Amico, L., 1983. Production and carrying capacity for the earthworm Lumbricus terrestris in culture. Soil Biology and Biochemistry 15, 51–54.

Hendrix, P.F., Mueller, B.R., Bruce, R.R., Langdale, G.W., Parmelee, R.W., 1992. Abundance and distribution of earthworms in relation to landscape factors on the Georgia Piedmont, U.S.A. Soil Biology and Biochemistry 24, 1357–1361.

Hoeffner, K., Monard, C., Santonja, M., Cluzeau, D., 2018. Feeding behaviour of epi-anecic earthworm species and their impacts on soil microbial communities. Soil Biology and Biochemistry 125, 1–9.

Hoeffner, K., Santonja, M., Cluzeau, D., Monard, C., 2019. Epi-anecic rather than strict-anecic earthworms enhance soil enzymatic activities. Soil Biology and Biochemistry 132, 93–100.

Hole, F.D., 1981. Effects of animals on soil. Geoderma 25, 75–112.

Hoogerkamp, M., Rogaar, H., Eijsackers, H.J.P., 1983. Effect of earthworms on grassland on recently reclaimed polder soils in the Netherlands, in: Satchell, J.E. (Ed.), Earthworm Ecology - from Darwin to Vermiculture. Chapman and Hall, London, pp. 85–105.

Revised manuscript

IUSS Working Group, W.R.B., 2015. World Reference Base for Soil Resources 2014, update 2015. International soil classification system for naming soils and creating legends for soil maps. (No. 106), World Soil Resources Reports. FAO, Rome.

Jamagne, M., 2011. Grands paysages pédologiques de France, 1st ed. Editions QUAE GIE, Versailles.

Jégou, D., Cluzeau, D., Balesdent, J., Trehen, P., 1998. Effects of four ecological categories of earthworms on carbon transfer in soil. Applied Soil Ecology 9, 249–255.

Joschko, M., Fox, C.A., Lentzsch, P., Kiesel, J., Hierold, W., Krück, S., Timmer, J., 2006. Spatial analysis of earthworm biodiversity at the regional scale. Agriculture, Ecosystems & Environment 112, 367–380.

King, K.L., Hutchinson, K.J., 2007. Pasture and grazing land: assessment of sustainability using invertebrate bioindicators. Australian Journal of Experimental Agriculture 47, 392–403.

Kovács-Hostyánszki, A., Elek, Z., Balázs, K., Centeri, C., Falusi, E., Jeanneret, P., Penksza, K., Podmaniczky, L., Szalkovszki, O., Báldi, A., 2013. Earthworms, spiders and bees as indicators of habitat quality and management in a low-input farming region—A whole farm approach. Ecological Indicators, Biodiversity Monitoring 33, 111–120.

Lapied, E., Nahmani, J., Rousseau, G.X., 2009. Influence of texture and amendments on soil properties and earthworm communities. Applied Soil Ecology 43, 241–249.

Larsen, T., Pollierer, M.M., Holmstrup, M., D'Annibale, A., Maraldo, K., Andersen, N., Eriksen, J., 2016. Substantial nutritional contribution of bacterial amino acids to earthworms and enchytraeids: A case study from organic grasslands. Soil Biology and Biochemistry 99, 21–27.

Lavelle, P., 1983. The structure of earthworm communities, in: Satchell, J.E. (Ed.), Earthworm Ecology - from Darwin to Vermiculture. Chapman and Hall, London, pp. 449–466.

Lee, K.E., 1985. Earthworms – Their ecology and relationships with soils and land use. Academic Press, Sydney.

Lüscher, G., Jeanneret, P., Schneider, M.K., Turnbull, L.A., Arndorfer, M., Balázs, K., Báldi, A., Bailey, D., Bernhardt, K.G., Choisis, J.-P., Elek, Z., Frank, T., Friedel, J.K., Kainz, M., Kovács-Hostyánszki, A., Oschatz, M.-L., Paoletti, M.G., Papaja-Hülsbergen, S., Sarthou, J.-P., Siebrecht, N., Wolfrum, S., Herzog, F., 2014. Responses of plants, earthworms, spiders and bees to geographic location, agricultural management and surrounding landscape in European arable fields. Agriculture, Ecosystems & Environment 186, 124–134.

Ma, W.-C., Brussaard, L., de Ridder, J.A., 1990. Long-term effects of nitrogenous fertilizers on grassland earthworms (Oligochaeta: Lumbricidae): Their relation to soil acidification. Agriculture, Ecosystems & Environment 30, 71–80.

Revised manuscript

474 Marshall, E.J.P., Moonen, A.C., 2002. Field margins in northern Europe: their functions and interactions
475 with agriculture. Agriculture, Ecosystems & Environment, The Ecology of Field Margins in
476 European Farming Systems 89, 5–21.
477 Maudsley, M., Seeley, B., Lewis, O., 2002. Spatial distribution patterns of predatory arthropods within an
478 English hedgerow in early winter in relation to habitat variables. Agriculture, Ecosystems &
479 Environment, The Ecology of Field Margins in European Farming Systems 89, 77–89.
480 McCallum, H.M., Wilson, J.D., Beaumont, D., Sheldon, R., O'Brien, M.G., Park, K.J., 2016. A role for
481 liming as a conservation intervention? Earthworm abundance is associated with higher soil pH and
482 foraging activity of a threatened shorebird in upland grasslands. Agriculture Ecosystems &
483 Environment 223, 182–189.
484 Michon, J., 1954. Contribution expérimentale à l'étude de la biologie des Lumbricidae. Les variations
485 pondérales au cours des différentes modalités du développement postembryonnaire. Poitiers
486 University, Poitiers.
487 Muldowney, J., Curry, J.P., O'Keeffe, J., Schmidt, O., 2003. Relationships between earthworm populations,
488 grassland management and badger densities in County Kilkenny, Ireland. Pedobiologia, The 7th
489 International Symposium on Earthworm Ecology 47, 913–919.
490 Neuhauser, E.F., Hartenstein, R., Kaplan, D.L., 1980. Growth of the earthworm Eisenia foetida in relation
491 to population density and food rationing. Oikos 35, 93–98.
492 Nuutinen, V., Butt, K.R., Jauhiainen, L., Shipitalo, M.J., Sirén, T., 2014. Dew-worms in white nights: High-
493 latitude light constrains earthworm (Lumbricus terrestris) behaviour at the soil surface. Soil Biology
494 and Biochemistry 72, 66–74.
495 O'Brien, J., Elliott, S., Hayden, T.J., 2016. Use of hedgerows as a key element of badger (Meles meles)
496 behaviour in Ireland. Mammalian Biology 81, 104–110.
497 Pelosi, C., Barot, S., Capowiez, Y., Hedde, M., Vandenbulcke, F., 2014. Pesticides and earthworms. A
498 review. Agronomy for Sustainable Development 34, 199–228.
499 Piearce, T.G., 1978. Gut contents of some lumbricid earthworms. Pedobiologia 18, 153–157.
500 Pietola, L., Horn, R., Yli-Halla, M., 2005. Effects of trampling by cattle on the hydraulic and mechanical
501 properties of soil. Soil and Tillage Research 82, 99–108.
502 Ponge, J.F., Pérès, G., Guernion, M., Ruiz-Camacho, N., Cortet, J., Pernin, C., Villenave, C., Chaussod, R.,
503 Martin-Laurent, F., Bispo, A., Cluzeau, D., 2013. The impact of agricultural practices on soil biota:
504 A regional study. Soil Biology and Biochemistry 67, 271–284.
505 R. Core Team, 2017. R: A language and environment for statistical computing. R foundation for statistical
506 computing, Vienna, Austria.

Revised manuscript

Richard, B., Legras, M., Margerie, P., Mathieu, J., Barot, S., Caro, G., Desjardins, T., Dubs, F., Dupont, L., Decaëns, T., 2012. Spatial organization of earthworm assemblages in pastures of northwestern France. European Journal of Soil Biology 53, 62–69.

Roarty, S., Schmidt, O., 2013. Permanent and new arable field margins support large earthworm communities but do not increase in-field populations. Agriculture, Ecosystems & Environment 170, 45–55.

Rutgers, M., Orgiazzi, A., Gardi, C., Römbke, J., Jänsch, S., Keith, A.M., Neilson, R., Boag, B., Schmidt, O., Murchie, A.K., Blackshaw, R.P., Pérès, G., Cluzeau, D., Guernion, M., Briones, M.J.I., Rodeiro, J., Piñeiro, R., Cosín, D.J.D., Sousa, J.P., Suhadolc, M., Kos, I., Krogh, P.-H., Faber, J.H., Mulder, C., Bogte, J.J., Wijnen, H.J. va., Schouten, A.J., Zwart, D. de, 2016. Mapping earthworm communities in Europe. Applied Soil Ecology 97, 98–111.

Rutgers, M., Schouten, A.J., Bloem, J., Eekeren, N.V., Goede, R.G.M.D., Akkerhuis, G.A.J.M.J., Wal, A.V. der, Mulder, C., Brussaard, L., Breure, A.M., 2009. Biological measurements in a nationwide soil monitoring network. European Journal of Soil Science 60, 820–832.

Smith, J., Potts, S.G., Woodcock, B.A., Eggleton, P., 2008. Can arable field margins be managed to enhance their biodiversity, conservation and functional value for soil macrofauna? Journal of Applied Ecology 45, 269–278.

Tews, J., Brose, U., Grimm, V., Tielbörger, K., Wichmann, M.C., Schwager, M., Jeltsch, F., 2004. Animal species diversity driven by habitat heterogeneity/diversity: the importance of keystone structures. Journal of Biogeography 31, 79–92.

Thomas, Z., Abbott, B.W., Troccaz, O., Baudry, J., Pinay, G., 2016. Proximate and ultimate controls on carbon and nutrient dynamics of small agricultural catchments. Biogeosciences 13, 1863–1875.

Vanbergen, A.J., Watt, A.D., Mitchell, R., Truscott, A.-M., Palmer, S.C.F., Ivits, E., Eggleton, P., Jones, T.H., Sousa, J.P., 2007. Scale-specific correlations between habitat heterogeneity and soil fauna diversity along a landscape structure gradient. Oecologia 153, 713–725.

Vickery, J.A., Feber, R.E., Fuller, R.J., 2009. Arable field margins managed for biodiversity conservation: A review of food resource provision for farmland birds. Agriculture, Ecosystems & Environment 133, 1–13.

Vršic, S., 2011. Soil erosion and earthworm population responses to soil management systems in steep-slope vineyards. Plant Soil Environ 57, 258–263.

Walter, C., Merot, P., Layer, B., Dutin, G., 2003. The effect of hedgerows on soil organic carbon storage in hillslopes. Soil Use and Management 19, 201–207.

Revised manuscript

Werling, B.P., Dickson, T.L., Isaacs, R., Gaines, H., Gratton, C., Gross, K.L., Liere, H., Malmstrom, C.M., Meehan, T.D., Ruan, L., Robertson, B.A., Robertson, G.P., Schmidt, T.M., Schrotenboer, A.C., Teal, T.K., Wilson, J.K., Landis, D.A., 2014. Perennial grasslands enhance biodiversity and multiple ecosystem services in bioenergy landscapes. Proceedings of the National Academy of Sciences 111, 1652–1657.

Zaller, J.G., Arnone, J.A., 1999. Earthworm responses to plant species' loss and elevated CO2 in calcareous grassland. Plant and Soil 208, 1–8.

Revised manuscript

548 TABLES

549

550 **Table 1** ANOVA results of multiple linear models testing for the effects of soil properties, grassland management and landscape diversity
551 on total earthworm abundance, total biomass, species richness and Shannon diversity index and evenness index (when VIF > 5). *F*-
552 values and associated *P*-values are indicated. Significant *P*-values are indicated in bold (P < 0.05). df = degrees of freedom, %SS =
553 percentage of sum of square.

	Total abundance				Total biomass				Total richness				Shannon				Equitability			
	df	%SS	*F*	*P*	df	%SS	*F*	*P*	df	%SS	*F*	*P*	df	%SS	*F*	*P*	df	%SS	*F*	*P*
Soil properties																				
Clay content	1	5.9	2.2	0.157																
Sand content	1	5.2	1.9	0.182	1	0.4	0.1	0.711	1	0.1	0.0	0.841								
Organic matter content	1	14.4	5.3	**0.033**	1	6.0	2.1	0.164												
pH	1	13.4	5.0	**0.039**	1	19.4	6.8	**0.018**												
Grassland management																				
Grassland age													1	7.5	9.9	0.107	1	7.3	2.4	0.141
Livestock unit					1	16.3	5.7	**0.028**	1	31.3	9.6	**0.005**	1	26.2	2.8	**0.005**	1	15.1	4.8	**0.040**
Fertilisation																				
Landscape diversity																				
Landscape Richness																				
SHDI	1	12.3	4.6	**0.047**	1	6.4	2.2	0.153												
Plant diversity													1	13.2	5.0	**0.037**	1	15.1	4.8	**0.040**
Residuals	18	48.7			18	51.5			21	68.6			20	53.1			20	62.5		

554

Revised manuscript

555 FIGURE LEGENDS

556

557 **Fig. 1.** Relationship between total earthworm biomass and soil pH. R^2 and associated *P*-value of
558 the linear regression are indicated.

559

560 **Fig. 2.** Relationships between livestock unit and (a) anecic earthworm abundance, (b) epi-anecic
561 earthworm abundance and (c) Shannon index. R^2 and associated *P*-values of linear regressions are
562 indicated.


563

564 **Fig. 3.** Earthworm species richness in plots with a hedgerow or with a ditch (i.e. control plot without
565 hedgerow) for grassland of (a) 1-year-old, (b) 2-year-old and (c) 7-year-old. Values are means ±
566 SD; n = 3. Different letters denote significant differences between the two plots with a > b (post
567 hoc Tukey test results).

568

Revised manuscript


569 **Fig. 1.**



570
571
572

Revised manuscript


573 **Fig. 2.**
574



575
576

Revised manuscript

577 **Fig. 3.**



578
579

Revised manuscript