



**HAL**  
open science

## Dynamics and genetic diversity of *Haemophilus influenzae* carriage among French pilgrims during the 2018 Hajj: A prospective cohort survey

Van-Thuan Hoang, Thi-Loi Dao, Tran Duc Anh Ly, Khadidja Belhouchat, Kamel Larbi Chaht, Saber Yezli, Badriah Alotaibi, Didier Raoult, Philippe Parola, Vincent Pommier De Santi, et al.

► **To cite this version:**

Van-Thuan Hoang, Thi-Loi Dao, Tran Duc Anh Ly, Khadidja Belhouchat, Kamel Larbi Chaht, et al.. Dynamics and genetic diversity of *Haemophilus influenzae* carriage among French pilgrims during the 2018 Hajj: A prospective cohort survey. *TRAVEL MEDICINE AND INFECTIOUS DISEASE*, 2020, 38, 10.1016/j.tmaid.2020.101883 . hal-03149263

**HAL Id: hal-03149263**

**<https://hal-amu.archives-ouvertes.fr/hal-03149263>**

Submitted on 26 Sep 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution-NonCommercial 4.0 International License

**Dynamics and genetic diversity of *Haemophilus influenzae* carriage among French pilgrims during the 2018 Hajj: A prospective cohort survey**

Van-Thuan Hoang<sup>1,2,3</sup>, Thi-Loi Dao<sup>1,2,3</sup>, Tran Duc Anh Ly<sup>1,2</sup>, Khadidja Belhouchat<sup>1,2</sup>, Kamel Larbi Chaht<sup>1,2</sup>, Saber Yezli<sup>4</sup>, Badriah Alotaibi<sup>4</sup>, Didier Raoult<sup>2,5</sup>, Philippe Parola<sup>1,2</sup>, Vincent Pommier de Santi<sup>1,2,6</sup>, Pierre-Edouard Fournier<sup>1,2</sup>, Philippe Gautret<sup>1,2\*</sup>

<sup>1</sup>Aix Marseille Univ, IRD, AP-HM, SSA, VITROME, Marseille, France

<sup>2</sup>IHU-Méditerranée Infection, Marseille, France

<sup>3</sup>Thai Binh University of Medicine and Pharmacy, Thai Binh, Viet Nam

<sup>4</sup>The Global Centre for Mass Gatherings Medicine, Ministry of Health, Riyadh, Saudi Arabia

<sup>5</sup>Aix Marseille Univ, MEPHI, Marseille, France

<sup>6</sup>French Military Center for Epidemiology and Public Health Marseille, France

Running title: Dynamics and genetic diversity of *H. influenzae*

\*Corresponding author:

Philippe Gautret

VITROME, Institut Hospitalo-Universitaire Méditerranée Infection, 19-21 Boulevard Jean Moulin 13385 Marseille Cedex 05, France. Phone: + 33 (0) 4 13 73 24 01. Fax: + 33 (0) 4 13 73 24 02. E-mail address: [philippe.gautret@club-internet.fr](mailto:philippe.gautret@club-internet.fr)

## Abstract

**Background:** To investigate the genetic diversity of *Haemophilus influenzae* positive samples among French pilgrims attending the 2018 Hajj pilgrimage.

**Method:** After screening by qPCR, multilocus sequence typing was performed for all *H. influenzae*-positive samples. The following housekeeping genes were amplified and assigned: *adk*, *atpG*, *frdB*, *fucK*, *mdh*, *pgi* and *recA*.

**Results:** 121 pilgrims were included. *H. influenzae* was positive in 35.5% pre-Hajj samples, 12.4% at day five post-arrival, 15.7% at day 12 post-arrival, and 43.0% post-Hajj. Of the 129 positive swabs for *H. influenzae*, only one sample at D12 was negative for all seven genes amplified by standard PCR. The *adk*, *atpG*, *frdB*, *mdh*, *pgi*, *recA* and *fucK* genes were positive in 123, 107, 122, 70, 127, 118 and 69 samples, respectively. One sequence of *atpG* and two of *recA* genes were not possible to assign. None of the sequences of *fucK* gene was successfully obtained. Consequently, a complete sequence type characterisation was not possible. Of the 128 obtained strains, 111 had distinct patterns of alleles.

**Conclusion:** *H. influenzae* genotypes acquired were completely different from those present at pre-Hajj. We observed a great biodiversity and a lack of clonality of *H. influenzae* among French pilgrims during the 2018 Hajj. Further studies aiming at studying the genome of Hajj-acquired *H. influenzae* isolates are needed to define the clinical burden of *H. influenzae* infection during Hajj and to evaluate the potential interest of vaccination in Hajj pilgrims.

**Key words:** Hajj, pilgrims; *Haemophilus influenzae*; MLST; genetic diversity

## Introduction

Every year, the Kingdom of Saudi Arabia (KSA) hosts more than two million people from about 180 countries during the Hajj pilgrimage. This number continues to increase, and the estimated number of annual attendees is expected to reach 4.5 million by 2030 [1]. Crowding and the extreme heat in Mecca, are known risk factors for the transmission of respiratory infections at the Hajj [2]. Infectious diseases accounted for 53% of diagnoses in outpatients consulting at the Indian Medical Mission in Mecca during the 2014–2016 Hajj, with respiratory tract infections (RTIs) and gastroenteritis being the most common [3]. The effectiveness of individual non-pharmaceutical preventive measures against infectious diseases is uncertain [3]. Despite several preventive measures [4], including hand hygiene, handkerchief use, face mask use, and vaccinations against influenza and pneumococcal infections in at-risk individuals being recommended, RTIs remain frequent among Hajj pilgrims. As an example, up to 90% of French pilgrims suffered an RTI during the 2012–2014 Hajj [5, 6]. Besides the widespread acquisition of respiratory viruses, bacteria are also frequently isolated among ill and asymptomatic pilgrims with *Haemophilus influenzae* being the most common [7-10]. Most cohort surveys among pilgrims used real-time PCR to detect the carriage of *H. influenzae*. Based on a culture method, Nik Zuraina showed that *H. influenzae* was the most predominant bacterium isolated (60%) from Malaysian pilgrims with RTIs during the Hajj [9]. In a 2018 study conducted among French Hajj pilgrims the carriage of *Staphylococcus aureus*, *Moraxella catarrhalis*, *Streptococcus pneumoniae* and *Klebsiella pneumoniae* increased following participation in the Hajj [11]. In contrast, *H. influenzae* carriage first decreased on D5 and D12 post-arrival by a factor of 2.5 and then increased in post-Hajj samples to reach carriage rates higher than that of pre-Hajj samples

[11]. The objective of this study was to investigate the genetic diversity of these *H. influenzae* positive samples among French pilgrims before, during and after the 2018 Hajj.

## **Methods**

### **Participants and study design**

This study included pilgrims travelling from Marseille, France, to Mecca, KSA during the 2018 Hajj. Participant were recruited at a single specialized travel agency and followed-up during the travel by two bilingual (French and Arabic) medical doctors. All pilgrims departed to KSA at the same date, were housed in the same accommodation and performed the rituals together during their stay. To evaluate the dynamic of *H. influenzae* during the Hajj, all pilgrims underwent four successive systematic nasopharyngeal swabs at different times: pre-travel, five to six days post arrival (D5 sample), 12 to 13 days post arrival (D12 sample) and just prior to leaving KSA (post-Hajj sample). The Hajj rituals took place from 19–24 August, corresponding to days 14 to 19 post-arrival (Figure 1).

### **Respiratory specimen and screening for *H. influenzae* by real-time qPCR**

The sampling was done by the doctors accompanying the group, in a standardized way (3 cm in the nostril, 5 turns; post wall of the pharynx, 5 streaks). Obtained swabs were transferred to Sigma-Virocult® medium and kept at room temperature (20°C) during travel and then stored at -80°C until processing.

The DNA were extracted from the samples using the EZ1 Advanced XL (Qiagen, Hilden, German) with the Virus Mini Kit v2.0 (Qiagen) according to the manufacturer's

recommendations. All quantitative real-time PCR were performed using a C1000 Touch™ Thermal Cycle (Bio-Rad, Hercules, CA, USA).

Real-time PCR amplifications were carried out using LightCycler® 480 Probes Master kit (Roche diagnostics, France) according to the manufacturer's recommendations. The *SHD* gene of *H. influenzae* was amplified with internal DNA extraction controls TISS, as previously described [12]. Negative controls (PCR mix) and positive controls (DNA from *H. influenzae* strain) were included in each run. Positive results of DNA amplification were defined as those with a cycle threshold (CT) value  $\leq 35$ .

### **Multilocus sequence typing (MLST) of *H. influenzae***

MLST was performed for all positively screened samples. The housekeeping genes *adk*, *atpG*, *frdB*, *fucK*, *mdh*, *pgi* and *recA* were amplified by standard PCR, as previously described [13], then sequenced. The sequences were compared to those existing in GenBank. Positive samples which presented at least one of seven genes using standard PCR were considered as being *H. influenzae* MLST positive samples. Purified PCR products were sequenced using specific primers and the BigDye Terminator® version 1.1 cycle sequencing ready reaction mix (Applied Biosystems, Foster City, CA, USA). Sequencing was performed using an Applied Biosystems 3130 platform (ABI PRISM, PE Applied Biosystems, USA). The sequences obtained were edited and assembled using Chromas Pro 1.77 (Technelysium Pty Ltd, Australia), aligned with *H. influenzae* strains from GenBank. Allele numbers and sequence types (ST) were assigned using the *H. influenzae* MLST website (<http://haemophilus.mlst.net>).

## **Results**

### **Characteristics of study participants and clinical symptoms**

The characteristics of study participants and clinical symptoms have been detailed elsewhere [11]. To summarise, 121 pilgrims were included, with a sex ratio of 1:1.3 and a median age of 61 years (interquartile = 56-66 years, range = 26–83 years). A total of 113/121 (93.4%) pilgrims presented at least one respiratory symptom during their stay in the KSA. Antibiotic use for RTIs was reported by 58.7% pilgrims. The mean time between arrival in the KSA and the onset of respiratory symptoms was  $8.7 \pm 4.6$  days (range = 1 – 21 days).

### **Identification of *H. influenzae* among French pilgrims during the Hajj**

#### ***H. influenzae* screening by real-time PCR**

Of the 484 swabs tested, 129 (26.7%) were positive [11]. Of the 121 pilgrims included, 75 (62.0%) at least one of their samples was positive for *H. influenzae* and all four samples of the remaining 46 were negative. The prevalence of *H. influenzae* carriage was 43/121 (35.5%) in samples obtained before leaving France, 15/121 (12.4%) at D5, 19/121 (15.7%) at D12 and 52/121 (43.0%) before leaving Saudi Arabia [11].

#### **Diversity of encapsulated and non-encapsulated *H. influenzae* using multi-locus sequence typing**

Of the 129 swabs that were positive for *H. influenzae*, only one sample at D12 was negative for all seven genes amplified by standard PCR. Figure 2 shows the distribution of the 128 MLST positive samples among the 75 pilgrims who presented at least one positive sample. A total of 40/121 (33.0%) pilgrims presented only one of the four samples that was positive for at least one of the seven genes (14 at pre-Hajj, 1 at D5, 3 at D12 and 12 at post-Hajj samples). In addition, 21/121 (17.4%), 10/121 (8.3%) and 4/121 (3.3%) pilgrims had two, three and four MLST positive samples, respectively.

The *adk*, *atpG*, *frdB*, *mdh*, *pgi*, *recA* and *fucK* genes were positive in 123, 107, 122, 70, 127, 118 and 69 samples, respectively. All sequences of *adk*, *frdB*, *mdh* and *pgi* genes were obtained from GenBank or using the *H. influenzae* MLST website (GenBank accession numbers MN555328, MN607727–MN607848, MN617868–MN617990, MN617991–MN618060 and MN627489–MN627615). A total of 106/107 and 116/118 sequences of *atpG* and *recA* genes were assigned with success respectively (GenBank accession numbers MN607849 – MN607953 and MN627616 – MN627731). One sequence of *atpG* and two of *recA* genes were not possible to assign, using Chromas Pro 1.77. However, none of the sequences of the *fucK* gene was successfully obtained (Figure 3). Consequently, complete sequence type characterisation was not possible. Nevertheless, our results enabled us to differentiate between strains. The supplementary data shows the diversity of the six genes obtained from pilgrim strains. Of the 128 obtained strains, 111 had distinct patterns of alleles. In post-Hajj samples, five distinct strains were each found in two pilgrims and one was found in three pilgrims. Samples with similar strains were also found in one pre-Hajj and one post-Hajj samples in two distinct individuals and another strain was found in one D12 and one post-Hajj samples in two other pilgrims.

## **Discussion**

In this study, we evaluated the dynamics and genetic diversity of *H. influenzae* carriage among French pilgrims during the 2018 Hajj season. Our main results are as follows: 1 - *H. influenzae* carriage was frequent among pilgrims before leaving France and increased following participation in the Hajj, following a transient decrease; 2 - *H. influenzae* genotypes acquired in the KSA were completely different from those present before leaving France and 3 - we



observed a great biodiversity and a lack of clonality of *H. influenzae* among pilgrims during the Hajj.

*H. influenzae* is a commensal organism of the human respiratory tract and is a common cause of upper and lower RTIs in adults [14]. Occasionally, it also causes life-threatening invasive diseases such as meningitis in adults [15]. This organism has been described as commensal in 75% of the healthy adult population [15] and typing of the *H. influenzae* isolates is essential in order to confirm its pathogenicity in pilgrims. Within the species, *H. influenzae* type b (Hib) is known to cause severe forms of infections [16]. Non-capsulated (non-typeable) *H. influenzae* is commonly carried in the pharynx and is one of the major causes of acute otitis media in children. It can cause diseases of the upper and lower respiratory tracts, including sinusitis and pneumonia [13]. However, previous studies conducted in some countries using the Hib conjugate vaccines reported that non-typeable and non-b serotypes caused most invasive diseases in adults over the age of 65 [17-19]. The World Health Organization recommends the epidemiological surveillance of *H. influenzae* to evaluate the current burden of its associated diseases and, consequently, to ascertain the practicality of Hib vaccine in the affected regions.

Pneumonia is a leading cause of hospitalisation during the annual Islamic pilgrimage. Memish conducted a study among pilgrims with severe community-acquired pneumonia who were admitted to 15 healthcare facilities in the cities of Makkah and Medina in Saudi Arabia [20]. Bacterial pathogens were detected in 84.6% of cases. *H. influenzae* and *S. pneumoniae* were the predominant bacteria, detected in 57.7% and 53.8% of patients, respectively. In a cohort study conducted among international pilgrims, Memish showed that the carriage of *H. influenzae* detected by qPCR was low. Only 2.3% of pilgrims were positive for this bacterium pre-Hajj and 11.7% post-Hajj [21]. Likewise, in a longitudinal survey conducted in 254,823 Iranian pilgrims

recruited at 1,352 Hajj caravans over a five-year period (2004-2009), *H. influenzae* was identified in only 9.1% of the 357 tested samples [22]. Among French pilgrims from Marseille, *H. influenzae* pre-Hajj prevalence varied according to the Hajj season: 50% in 2014, 0.9% in 2015, 2.8% in 2016 and 52.8% in 2017, while post-Hajj prevalence was consistently high with 67.8%, 45.5%, 41.0% and 53.5%, respectively [8, Gautret, unpublished data]. This suggests that *H. influenzae* carriage in Marseille may be subject to yearly variations. In contrast, a frequent acquisition of this bacterium at the Hajj is observed each year. In line with this result, Wilkinson conducted a prospective analysis in the United Kingdom, surveying 127 patients with chronic obstructive pulmonary disease (COPD) aged between 40 and 85 over a period of two years (from June 2011 to June 2012) for the occurrence of acute exacerbation and respiratory infections in COPD. Their results showed, for the first time, that changes in the yearly COPD exacerbation rate may be associated with variations in *H. influenzae* colonisation [23].

In our study, *H. influenzae* alleles varied according to the sampling time during Hajj. In pilgrims who were positive for *H. influenzae* in all four samples, *H. influenzae* genotypes were different in most samples, suggesting that pilgrims successively acquired different new *H. influenzae* genotypes during the pilgrimage. The reason for the initial clearance of *H. influenzae* carriage following arrival to the KSA may have resulted from the high rate of antibiotic consumption by pilgrims [24]. The large diversity of genotypes acquired during the stay in the KSA suggests that the source of infection was polyclonal and was likely to be external to the group of French pilgrims, with very little transmission between them.

In our study, none of the sequences of the *fucK* gene was successfully assigned. Other researchers have reported that certain *H. influenzae* isolates may lack the *fucK* gene, one of the seven genes used in the *H. influenzae* MLST scheme [13, 25-27]. The absence of this gene has

historically confounded attempts at *H. influenzae* speciation, such as the probable incorrect assignment of *fucK*-negative *H. influenzae* strains as *H. haemolyticus* [28]. These ‘fuzzy’ isolates should therefore be considered to be fucose-negative *H. influenzae*. It is unfortunate that the developers of the *H. influenzae* MLST scheme inadvertently chose a variably present gene for genotyping purposes. Price confirmed the existence of fucose-negative *H. influenzae* strains [29]. Redesign of the existing *H. influenzae* MLST scheme to incorporate a conserved seventh locus in *H. influenzae* and *H. haemolyticus* in place of the *fucK* gene may lead to greater uptake of this highly useful genotyping scheme around the world.

Our study has several limitations. First, we performed sequencing directly from samples, not on cultured isolates and we could not succeed in amplifying all seven genes in the MLST system. In addition, we could not eliminate the possibility that an individual sample could be positive for several genotypes of *H. influenzae*. The study was conducted among French pilgrims only and cannot be generalised to all pilgrims. Using qPCR to detect respiratory pathogens does not distinguish between dead and living micro-organisms. Nevertheless, our study is the first study on the dynamic and genetic diversity of *H. influenzae* in the Hajj context. Continuous monitoring of the molecular evolution in *H. influenzae* among Hajj pilgrims is necessary to explore how *H. influenzae* biofilm formation involves in the pathological process of infection and may help for clinical treatment. Moreover, constant vigilance and precaution such as a vaccine development is necessary to respond to new epidemiological trends of *H. influenzae* strains acquired by Hajj pilgrims. *H. influenzae* type b vaccination has been described by previous study that induces serum antibody production and reduces the nasopharyngeal carriage prevalence of this bacterium. Reduction in carriage also reduces transmission of *H. influenzae* between individuals [30]. Further studies aiming at studying the genome of Hajj-acquired *H. influenzae* isolates are

needed to better elucidate their ecological changes, to define the clinical burden of *H. influenzae* infection during Hajj and to evaluate the potential interest of vaccination in Hajj pilgrims. Currently, vaccination against invasive *H. influenzae* disease is not recommended for Hajj pilgrims [4].

### **Funding**

This study was supported by the Institut Hospitalo-Universitaire (IHU) Méditerranée Infection, the French National Research Agency under the “Investissements d’avenir” programme, reference ANR-10-IAHU-03, the Région Provence Alpes Côte d’Azur and European funding FEDER PRIMI.

### **Conflict of Interest**

The authors declare that they have no conflict of interest

### **Author Contributions Statement**

VTH, VPS, PEF and PG contributed to experimental design, interpretation and writing. PEF supported the technique. VTH conducted the technique. TDAL and TLD provided technical assistance. KB and KLC administered questionnaires, followed patient and collected samples. SY, BA, DR and PP contributed to critically reviewing the manuscript. PG coordinated the work.

## References

1. Ebrahim SH, Doumbia S, Elachola H, Memish ZA. "Healthy Hajj 2019" -- what you need to know, before you go. *Travel Med Infect Dis* 2019;30:1-3.
2. Memish ZA, Steffen R, White P, Dar O, Azhar EI, Sharma A, et al. Mass gatherings medicine: public health issues arising from mass gathering religious and sporting events. *Lancet* 2019;393:2073-84.
3. Benkouiten S, Brouqui P, Gautret P. Non-pharmaceutical interventions for the prevention of respiratory tract infections during Hajj pilgrimage. *Travel Med Infect Dis* 2014;12:429-42.
4. The Ministry of Health in the Kingdom of Saudi Arabia. 2019/1440H-Hajj and Umrah Health Regulations. <https://www.moh.gov.sa/en/Hajj/HealthGuidelines/HealthGuidelinesDuringHajj/Pages/HealthRequirements.aspx>; 2019
5. Gautret P, Benkouiten S, Griffiths K, Sridhar S. The inevitable Hajj cough: surveillance data in French pilgrims, 2012–2014. *Travel Med Infect Dis* 2015;13:485–9.
6. Hashim S, Ayub ZN, Mohamed Z, Hasan H, Harun A, Ismail N, et al. The prevalence and preventive measures of the respiratory illness among Malaysian pilgrims in 2013 Hajj season. *J Travel Med* 2016;23:tav019.
7. Hoang VT, Gautret P. Infectious Diseases and Mass Gatherings. *Curr Infect Dis Rep* 2018;20:44.
8. Hoang VT, Meftah M, Anh Ly TD, Drali T, Yezli S, Alotaibi B, et al. Bacterial respiratory carriage in French Hajj pilgrims and the effect of pneumococcal vaccine and other

- individual preventive measures: A prospective cohort survey. *Travel Med Infect Dis* 2019;31:101343.
9. Nik Zuraina NMN, Sarimah A, Suharni M, Hasan H, Suraiya S. High frequency of *Haemophilus influenzae* associated with respiratory tract infections among Malaysian Hajj pilgrims. *J Infect Public Health* 2018;11:878-83.
  10. Shirah BH, Al Nozha FA, Zafar SH, Kalumian HM. Mass Gathering Medicine (Hajj Pilgrimage in Saudi Arabia): The Outcome of Cardiopulmonary Resuscitation during Hajj. *J Epidemiol Glob Health* 2019;9:71-5.
  11. Hoang VT, Dao TL, Ly TDA, Belhouchat K, Larbi Chaht K, Gaudart J, et al. The dynamics and interactions of respiratory pathogen carriage among French pilgrims during the 2018 Hajj. *Emerg Microbes Infect* 2019. <https://doi.org/10.1080/22221751.2019.1693247>
  12. Memish ZA, Assiri A, Turkestani A, Yezli S, Al Masri M, Charrel R, et al. Mass gathering and globalization of respiratory pathogens during the 2013 Hajj. *Clin Microbiol Infect*. 2015;21:571.e1-8.
  13. Meats E, Feil EJ, Stringer S, Cody AJ, Goldstein R, Kroll JS, et al. Characterization of encapsulated and noncapsulated *Haemophilus influenzae* and determination of phylogenetic relationships by multilocus sequence typing. *J Clin Microbiol* 2003;41:1623-36.
  14. Nicola J. High, Feinan Fan, Joseph D. Schwartzman. Chapter 97 - *Haemophilus influenzae*. *Molecular Medical Microbiology* (Second Edition), 2015;3:1709-28.
  15. King P. *Haemophilus influenzae* and the lung (*Haemophilus* and the lung). *Clin Transl Med* 2012;1:10

16. Nix EB, Hawdon N, Gravelle S, Biman B, Brigden M, Malik S, et al. Risk of invasive *Haemophilus influenzae* type b (Hib) disease in adults with secondary immunodeficiency in the post-Hib vaccine era. *Clin Vaccine Immunol* 2012;19:766–71
17. Blain A, MacNeil J, Wang X, Bennett N, Farley MM, Harrison LH, et al. Invasive *Haemophilus influenzae* disease in adults  $\geq 65$  years, United States, 2011. *Open Forum Infect Dis* 2014;1:ofu044.
18. Adam HJ, Richardson SE, Jamieson FB, Rawte P, Low DE, Fisman DN. Changing epidemiology of invasive *Haemophilus influenzae* in Ontario, Canada: evidence for herd effects and strain replacement due to Hib vaccination. *Vaccine* 2010;28:4073–8.
19. Rubach MP, Bender JM, Mottice S, Hanson K, Weng HY, Korgenski K, et al. Increasing incidence of invasive *Haemophilus influenzae* disease in adults Utah, USA. *Emerg Infect Dis* 2011;17:1645–50.
20. Memish ZA, Almasri M, Turkestani A, Al-Shangiti AM, Yezli S. Etiology of severe community-acquired pneumonia during the 2013 Hajj—part of the MERS-CoV surveillance program. *Int J Infect Dis* 2014;25:186-90.
21. Memish ZA, Assiri A, Turkestani A, Yezli S, Al Masri M, Charrel R, et al. Mass gathering and globalization of respiratory pathogens during the 2013 Hajj. *Clin Microbiol Infect* 2015;21:571.e1-8.
22. Razavi SM, Sabouri-Kashani A, Ziaee-Ardakani H, Tabatabaei A, Karbakhsh M, Sadeghipour H, et al. Trend of diseases among Iranian pilgrims during five consecutive years based on a Syndromic Surveillance System in Hajj. *Med J Islam Repub Iran* 2013;27:179–85

23. Wilkinson TMA, Aris E, Bourne SC, Clarke SC, Peeters M, Pascal TG, et al. Drivers of year-to-year variation in exacerbation frequency of COPD: analysis of the AERIS cohort. *ERJ Open Res* 2019;5. pii:00248-2018.
24. Hoang VT, Nguyen TT, Belhouchat K, Meftah M, Sow D, Benkouiten S, et al. Antibiotic use for respiratory infections among Hajj pilgrims: A cohort survey and review of the literature. *Travel Med Infect Dis.* 2019;30:39-45.
25. Ridderberg W, Fenger MG, Nørskov-Lauritsen N. *Haemophilus influenzae* may be untypable by the multilocus sequence typing scheme due to a complete deletion of the fucose operon. *J. Med. Microbiol* 2010;59:740–2.
26. Shuel ML, Karlowsky KE, Law DK, Tsang RS. Nonencapsulated or nontypeable *Haemophilus influenzae* are more likely than their encapsulated or serotypeable counterparts to have mutations in their fucose operon. *Can. J. Microbiol* 2011;57:982–6.
27. De Gier C, Kirkham LA, Nørskov-Lauritsen N. Complete deletion of the fucose operon in *Haemophilus influenzae* is associated with a cluster in multilocus sequence analysis-based phylogenetic group II related to *Haemophilus haemolyticus*: implications for identification and typing. *J. Clin. Microbiol* 2015;53:3773–8.
28. Anderson R, Wang X, Briere EC, Katz LS, Cohn AC, Clark TA, Messonnier NE, et al. *Haemophilus haemolyticus* isolates causing clinical disease. *J. Clin. Microbiol* 2012;50:2462–5.
29. Price EP, Harris TM, Spargo J, Nosworthy E, Beissbarth J, Chang AB, et al. Simultaneous identification of *Haemophilus influenzae* and *Haemophilus haemolyticus* using real-time PCR. *Future Microbiol* 2017;12:585-93.




30. Hammitt LL, Crane RJ, Karani A, Mutuku A, Morpeth SC, Burbidge P et al. Effect of *Haemophilus influenzae* type b vaccination without a booster dose on invasive *H. influenzae* type b disease, nasopharyngeal carriage, and population immunity in Kilifi, Kenya: a 15-year regional surveillance study. *Lancet Glob Health*. 2016;4(3):e185-e194.

Figure 1: Study design of cohort survey on the dynamic of *Haemophilus influenzae* carriage among 121 French pilgrims during the 2018 Hajj season

Figure 2: Temporal distribution of 128 MLST positive samples among 75 pilgrims with at least one positive sample.

Figure 3: Number of MLST positive samples and sequences successfully assigned of seven MLST genes





Number

