Investigating the role of root exudates in the interaction between oak seedlings and purple moor grass in temperate forest

Marine Fernandez, Philippe Malagoli, Christiane Gallet, Catherine Fernandez, Antoine Vernay, Thierry Ameglio, Philippe Balandier

► To cite this version:

Marine Fernandez, Philippe Malagoli, Christiane Gallet, Catherine Fernandez, Antoine Vernay, et al.. Investigating the role of root exudates in the interaction between oak seedlings and purple moor grass in temperate forest. Forest Ecology and Management, 2021, 491 (119175), 10.1016/j.foreco.2021.119175 . hal-03196931

HAL Id: hal-03196931

<https://hal-amu.archives-ouvertes.fr/hal-03196931>

Submitted on 15 Apr 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives| 4.0 International License

Forest Ecology and Management

Investigating the role of root exudates in the interaction between oak seedlings and purple moor grass in temperate forest --Manuscript Draft--

Manuscript Number:	FORECO-D-20-01535R2
Article Type:	Full Length Article
Keywords:	interaction; competition; interference; allelopathy; forest regeneration
Corresponding Author:	Philippe Malagoli Université Clermont Auvergne AUBIERE, France
First Author:	Marine FERNANDEZ
Order of Authors:	Marine FERNANDEZ Philippe MALAGOLI Christiane GALLET Catherine FERNANDEZ Antoine VERNAY Thierry AMEGLIO Philippe BALANDIER
Abstract:	<p>Oak regeneration in temperate forests often fails in the presence of understory grass. Competition by resource exploitation between plants has been extensively studied. By contrast, competition by interference, especially chemical interference (allelopathy), has been much less thoroughly examined and its relative importance remains unclear. We investigated the influence of allelopathic interaction on plant performance (biomass production) in a pot experiment with sessile oak (<i>Quercus petraea</i>) and purple moor grass (<i>Molinia caerulea</i>), either sole- or mixed-grown. Plants were watered with either <i>Quercus</i> root exudates or <i>Molinia</i> root exudates. After 6 months of growth, oak biomass increment was significantly lowered by <i>Molinia</i> root exudates. The oak's root system was more strongly affected than its aerial part. <i>Quercus</i> root exudates favoured oak growth but did not affect moor grass. Conversely, <i>Molinia</i> root exudates had a small depressive effect on its own growth, but its biomass was favoured by the presence of oak grown in the same pot. Allelopathy reduced oak biomass by about 17%, and resource exploitation by an additional 33%, totalling 50%, assuming the two processes are strictly additive, which is debatable. Although untargeted metabolomic analysis by UHPLC failed to identify any potentially allelopathic substances involved, our study demonstrates a lower but critical contribution of chemical interference on oak seedling-moor grass competition compared to exploitation processes. To ensure oak regeneration, management of forest ecosystems should thus first focus on reducing moor grass close to oak seedlings to help decrease its allelopathic effect and ease resource competition.</p>
Suggested Reviewers:	Marie-Charlotte Nilsson marie-charlotte.nilsson@slu.se Expertise in N cycling within tree-graminoïds Lluís Coll lluis.coll@udl.cat Expertise in competition for soil nutrient root uptake into tree-graminoïds system Leslie Weston lewiston@csu.edu.au Expertise in belowground allelopathy Manuel Reigosa mreigosa@uvigo.es Expertise in plant physiology response to allelochemicals

1 **Highlights**

- 2 - Oak-moor grass interaction also relies on chemical interference
- 3 - Moor grass roots emit allelopathic compounds reducing oak root growth
- 4 - Moor grass interference critically contributes to reduce oak growth
- 5 - Oak regeneration management should first focus on reducing moor grass density

sessile oak (*Quercus petraea*) and purple moor grass (*Molinia caerulea*), either sole- or mixed-grown. Plants were watered with either *Quercus* root exudates or *Molinia* root exudates. After 6 months of growth, oak biomass increment was significantly lowered by *Molinia* root exudates. The oak's root system was more strongly affected than its aerial part. *Quercus* root exudates favoured oak growth but did not affect moor grass. Conversely, *Molinia* root exudates had a small depressive effect on its own growth, but its biomass was favoured by the presence of oak grown in the same pot. Resource exploitation had a more detrimental effect than allelopathy and both processes together decreasing oak biomass by 50%. Although untargeted metabolomic analysis by UHPLC failed to identify any potentially allelopathic substances involved, our study demonstrates a lower but critical contribution of chemical interference on oak seedling-moor grass competition compared to exploitation processes. To ensure oak regeneration, management of forest ecosystems should thus first focus on reducing moor grass close to oak seedlings to help decrease its allelopathic effect and ease resource competition.

43 Introduction

44 Identifying the balance of processes involved in plant-plant interactions has drawn much
45 attention. Numerous studies have focused on competition for light and soil resources, but the
46 mechanisms involved in soil resource competition (intrinsic root uptake capacity, monitoring
47 of resources uptake dynamic, roles of root exudation, etc.) are seldom investigated, especially
48 in the context of forest tree regeneration. Root competition (intra or inter-specific) has often
49 been described as one of the main interactions driving plant establishment in communities
50 (Casper and Jackson, 1997; Coomes and Grubb, 2000; Schenk, 2006). Two types of root
51 competition have been identified, based on two plant strategies, termed “scramble” and
52 “contest” by Schenk (2006). So far, most studies have focused on “scramble”, which is
53 competition by resource exploitation (Grime, 1974; Tilman, 1990; Wilson and Tilman, 1993).
54 This is when two species capture the same resources in the same finite space. Each species then
55 depletes the resource reservoir to the detriment of the other. “Contest”, or competition by
56 interference, includes mechanisms by which one species inhibits belowground resource access
57 to another species by active or passive release of chemicals from either roots or shoots. Plant
58 ecologists often use the term allelopathy to describe “contest” competition, but interference also
59 include other non-resource interactions, such as autotoxicity (Guretzky, 2020), space
60 competition (Casper and Jackson, 1997; Schenk, 2006), litter composition and thickness
61 (Foster, 1999; Michalet et al., 2017). Schenk (2006) defines allelopathy narrowly in terms of
62 resource access, but it can include all processes of chemically-mediated interference that alter
63 germination, growth, survival or reproduction (Reigosa et al., 1999; Rice, 1984).
64 Allelochemicals produced by plant organs are released into the environment through (i) release
65 of volatile organic compounds (Effah et al., 2019; Penuelas and Llusia, 1998; Santonja et al.,
66 2019), (ii) decomposition of fallen leaves and needles (Fernandez et al., 2016; Hashoum et al.,
67 2017; Nilsson, 1994), (iii) living root exudates (van Dam and Bouwmeester, 2016), and
68 (iv) decomposition of dead roots (Bertin et al., 2003; Fernandez et al., 2016; Mallik et al., 2016;
69 Tsunoda and van Dam, 2017). Phytotoxic substances in root exudates have been identified in
70 various species such as *Secale cereale* (Pérez and Ormeno-Núñez, 1991), *Avena fatua* (Pérez
71 and Ormeño-Núñez, 1991), *Cucumis sativus* (Yu et al., 2003; Yu and Matsui, 1994), *Oryza*
72 *sativa* (Kato-Noguchi, 2004), *Chenopodium murale* (Batish et al., 2007) and *Peperomia*
73 *argyreia* (Hao et al., 2010). Allelochemicals can alter neighbouring plant growth and
74 functioning either directly (e.g. by interfering with root system growth and development, root
75 nutrient uptake, or physiological processes) or indirectly through changes in belowground soil

properties, such as physicochemical conditions (pH, ions availability...) (Huang et al., 2013; Xuan et al., 2005; Zeng, 2014) or soil community diversity, including N₂-fixing bacteria and mycorrhizal associations). Seed germination and seedling establishment have been reported as life stages and processes often targeted by allelochemicals (Gallet and Pellissier, 2002). Allelopathy and competition by exploitation can act simultaneously. This makes determining the contributions of the two processes a challenge, especially under field conditions (Fernandez et al., 2016; Nilsson, 1994; Viard-Crtat et al., 2012). Evidence of allelopathic potential can be obtained by removing donor plants or organs, but this is not possible with roots (Fernandez et al., 2016; Fuerst and Putnam, 1983; Inderjit and Mallik, 2002; Olofsdotter et al., 1999; Ridenour and Callaway, 2001).

In most boreal and temperate forests, understorey vegetation has been shown to affect the establishment and growth of tree regeneration (Balandier et al., 2006; Davis et al., 1998; Pags and Michalet, 2003). Nilsson (1994) sought to separate and quantify contributions of exploitation competition and allelopathic effects of *Empetrum hermaphroditum* shoots on Scots pine (*Pinus sylvestris*) seedling growth. These authors pointed out both competition types were involved: *E. hermaphroditum* led to soil resource depletion but also exerted chemical inhibition on Scots pine seedling growth. Significant studies have been carried out on conifer regeneration failures in the presence of an ericaceous community (Mallik, 2003), underlining the potential role of allelochemicals released by ericaceous species that alter conifer seedling growth (Inderjit and Mallik, 2002, 1996; Mallik et al., 2016). More generally, interactions involving allelopathic substances in woodland have been identified in different tree species regeneration failures (Fisher, 1987) but are still poorly understood, especially in temperate deciduous forests.

Tree seedlings often show significant susceptibility to the presence of herbaceous neighbours. In particular, monopolistic perennial grasses, with a rapid vegetative development and a dense fasciculate root system, rapidly absorb soil resources such as inorganic nitrogen, competing by exploitation to the detriment of tree seedling growth and biomass (Balandier et al., 2006; Coll et al., 2003; Davies, 1987; Vernay et al., 2018a, 2016). At the opposite, with erectophile leaves (angle $\geq 65^\circ$, e.g. Balandier et al., 2006), perennial grasses generally induce not so strong competition for light (Gaudio et al., 2011; Picon-Cochard et al., 2006; Vernay et al., 2016). A common case is the regeneration of valuable oak species such as *Quercus petraea* in temperate forests, which is often jeopardized by the presence of interfering understorey species such as *Molinia caerulea*, a widespread monopolistic grass (Taylor et al., 2001). For instance, competition by exploitation for N-resources in favour of moor grass was described as

a major reason for oak regeneration failure (Vernay et al., 2018a, 2018b). However, the respective importance of allelopathy in oak-grass interactions remains to be unravelled (Becker and Lévy, 1982; Timbal et al., 1990). Some studies have demonstrated the allelopathic potential of some oak species (*Q. robur*, *Q. pubescens*, *Q. coccifera*) exerted both on model and cultivated species (lettuce, wheat, barley, lentil, chickpea, faba bean and flax) and native herbaceous species (Alrababah et al., 2009; Hashoum et al., 2017; Souto et al., 1994). Conversely, *Quercus* seedling susceptibility to tree allelochemicals has been demonstrated in very few studies (Fernandez et al. 2016; *Q. pubescens* and *Pinus halepensis*).

Studies generally focus on “one-way” allelopathic interaction, analysing effects of species A on species B but rarely testing effects of root exudates of species B on A, so that the resultant interaction between the two plants is often inaccurately assessed. Moreover, allelochemicals can have intraspecific allelopathic effects, also called auto-allelopathy or autotoxicity (Böttger et al., 2018; Friedman and Waller, 1985; Singh et al., 1999). Various grasses or crops are known to exert an allelopathic effect on themselves, such as wheat (Wu et al., 2007) or barley (Ben-Hammouda et al., 2002), which can influence the community dynamics. In an interspecific context, it is not known whether *Quercus* root exudates contain secondary metabolites, in addition to the nitrogen released in the soil, that can improve moor grass growth (Fernandez et al., 2020). Allelopathic potential of root exudates has received much attention but little is known about their facilitative potential (Li et al., 2016). Fernandez et al. (2020) concomitantly observed a negative effect of the presence of moor grass on oak seedling growth and a positive reverse effect largely due to rapid nitrogen transfers from tree seedling to grass. Interactions simultaneously involving both a competition effect of plant A on plant B and a facilitation effect of plant B on plant A are called antagonistic (Michalet et al., 2011; Pistón et al., 2018; Schöb et al., 2014, 2013; Zélé et al., 2018) and have been scantily described.

Based on two complementary experiments, the “two-way” allelopathic interaction between sessile oak seedling and purple moor grass was investigated and extended by a more specific focus on the “one-way” interaction to characterize the allelopathic effect of *Molinia* exudates on oak seedlings specifically on shoot and root biomass. In the “two-way” experiment, we hypothesized (i) that *Molinia* root exudates might contain chemicals adverse to oak seedling growth and (ii) oak seedling root exudates might contain chemicals facilitating moor grass growth. We also applied root exudates of *Quercus* and *Molinia* on oak and moor grass, respectively, to test inhibitory or facilitative potential in the context of intraspecific interaction.

Materials and Methods

Two experiments were conducted in pots under outdoor conditions in Clermont-Ferrand (Auvergne, France, 45°45'N 3°07'E, altitude 394 m a.s.l, temperate climate). Using pots allowed a better controlled irrigation and harvesting of the whole root system. A “two-way” experiment was managed to study both *Molinia* allelopathic effect on oak seedling and on conspecific *Molinia*, and the reciprocal, *Quercus* allelopathic effect on *Molinia* and conspecific oaks. This was done watering oak or *Molinia* grown alone with a water solution of root extracts of either oak or *Molinia*. In this experiment, we also considered the exploitative competition effects on *Molinia* and *Quercus* at the plant scale (total biomass), by growing the two species together in a same pot. In a “one-way” experiment, we only tested the allelopathic effect of *Molinia* on oak growth: here oak's shoot and root compartments were separately measured to focus on oak organs response to interference competition. One-year-old oak seedlings were sourced from a local nursery. *Molinia caerulea* (L.) Moench was collected in a local forest at Paray-le-Frésil (Auvergne Rhône-Alpes, France; 46°39'N 3°36'E). Oaks and moor grasses were planted in pots filled with natural forest soil (typical luvisol redoxisol pseudogley, sandy loam) collected in the same forest as the moor grasses. To avoid water stress, pots were irrigated to field capacity throughout the experiment. Humidity probes set up in some pots monitored soil water content in the middle of the pot. No fertilizer was added to the pot during the experiment.

“Two-way” experiment (**Figure 1A**)

The experiment lasted from March to September 2018. In March 2018, 30 1-year-old bare-root oak seedlings (*Quercus petraea* (Matt.) Liebl.) and 30 moor grass tufts were planted separately in 5 L plastic pots or together in 10 L plastic pots. Oaks were 51.7 ± 4.0 cm in height (mean $\pm$ SE), and 7.6 ± 1.3 mm in diameter on average. Moor grasses were 10.1 ± 1.0 g fresh weight. To prepare solutions of fresh root aqueous maceration, 12 donor oaks and 12 donor moor grasses planted in 5 L pots were collected at the rate of one plant per week (one solution of root exudate supply per week for 12 weeks). Intact fresh roots were washed, weighed and consistently macerated in distilled water for 48 h in darkness at room temperature. The concentration of these root exudates was 2% (root dry weight/water volume), which is the range commonly reported in literature (Fernandez et al., 2006: 2.5 and 5%; Hashoum et al., 2017: 1 and 2.5%; Nilsson et al., 1993: 0, 0.5, 1.5, 2.5 and 3.75%). The final volume of aqueous exudates was split and immediately poured into the pots containing the target oak seedling or moor grass. Concomitantly, sole-growing targets received an equivalent volume of distilled

water (control). Root exudates of *Quercus* and *Molinia* were then applied on soil immediately after maceration. The experimental design was as follows (Fig. 1):

(i) Six recipient oaks and six recipient moor grasses were planted alone in 5 L pots without addition of root exudates (“sole-grown”, SG), they received an equivalent volume of distilled water (control)

(ii) Six sole-growing recipient oaks received *Quercus* root exudates (“SG + *Quercus* exudates”) and six others received *Molinia* root exudates (“SG + *Molinia* exudates”),

(iii) Six sole-growing recipient-moor grasses received *Quercus* root exudates (“SG + *Quercus* exudates”) and six others received *Molinia* root exudates (“SG + *Molinia* exudates”),

(iv) One oak and one moor grass were planted in the same 10 L pot (“mixed-grown”) to analyse competition through physical interaction without addition of root exudates ($n = 6$).

Oak seedlings used in conditions (ii) and (iii) were on average 44.3 ± 13.6 g fresh weight. Oaks used in (i) and (iv) were 33.4 ± 10.5 g fresh weight. Because of this unintentional initial biomass difference, we could not compare final biomass of shoot and root between the different pot conditions. Alternatively, we compared response in total dry biomass by relative growth increment, i.e. considering the initial difference, in the four experimental conditions. Relative increment (RI) was calculated by the expression (South, 1995):

$$RI = \frac{DW_F - DW_I}{DW_I},$$

where DW_I was the initial dry weight of oak and moor grass when potting and DW_F was the final dry weight at harvest.

Plants were harvested in September 2018, i.e. after 6 months of growth. Shoots and roots were collected, washed and dried at 60 °C for 48 h before weighing.

One-way experiment (Figure 1B)

The experiment lasted from April 2017 to August 2017. In April 2017, 16 one-year-old bare-root oak seedlings (*Quercus petraea*) and 12 moor grass tufts were planted separately in 5 L plastic pots. In this experiment, moor grasses were used only as ‘donors’ to produce root exudates, and oaks were only targets. Oaks were 19.9 ± 6.7 g (mean $\pm$ SE) fresh weight, 40.8 ± 4.2 cm in height, and 4.9 ± 0.8 mm in diameter on average. Moor grasses were 10.3 ± 1.5 g fresh weight. In this experiment, special attention was paid to height and weight

homogenization of oaks to compare shoot and root biomass of controls and *Molinia* exudate receivers.

One donor moor grass tuft was harvested every week for 12 weeks from June 2017. Root exudates were obtained by the same method as described in the “two-way” experiment. The final volume of aqueous exudates was split weekly and immediately poured into the eight pots containing one target oak seedling each. Concomitantly, eight sole-growing target oak seedlings received an equivalent volume of distilled water (control).

Receiver oak seedlings were harvested in October 2017, *i.e.* after 6 months of growth. Oak shoots and roots were collected, roots were carefully separated from soil by washing, dried at 60 °C for 48 h, and weighed. Organ biomass was used as an integrated proxy of oak growth throughout the season.

Data analysis

Statistical analysis was performed using R software (Version 3.4.1.).

In the two-way experiment, data were means of $n = 6$ biological replicates. We used a linear model (lm function) to determine the response of relative dry biomass increment, the root/shoot (R/S) dry biomass ratio to competition (SG vs MG) and exudate exposure (SG + *Quercus* exudates, SG + *Molinia* exudates). A pairwise comparison was then applied to the different treatment combinations to assess the significant differences ($\alpha = 0.05$). Data were tested for normality using the Shapiro-Wilk test and for homoscedasticity using the Levene test before statistical analyses. Data were analysed using one-way ANOVA with Tukey’s *post hoc* test.

In the one-way experiment, data were the means of $n = 8$ biological replicates. We used a linear model (lm function) to determine the response of oak dry weight and oak relative dry biomass increment to the different pot conditions: SG and SG + *Molinia* exudates. A *t* test was used to examine the difference in shoot and root dry weight and RI between “sole-grown” and “SG + *Molinia* exudates”. Here, results were analysed with one level of significance ($\alpha = 0.05$).

Results

Response of relative biomass increment and root/shoot ratio to root exudate application (two-way experiment)

In the two-way experiment, *Molinia* root exudates (SG + *Molinia* exudates) had no significant effect on oak relative dry weight relative increment (RI) in comparison with SG (*p*

= 0.46, **Figure 2A, Tables S1 and S2**). Oak biomass was significantly lower in MG than in SG treatment ($p = 0.04$), and SG + *Molinia* exudate treatment was marginally higher than MG ($p = 0.09$). Globally, oak RI was ordered SG > SG + *Molinia* exudates > MG. Application of *Quercus* root exudates marginally increased oak RI compared to the sole-grown condition ($p = 0.08$).

For moor grasses, a marginally lower relative dry weight increment was recorded in the presence of its own root exudates compared to control ($p = 0.08$, **Figure 2B, Tables S1 and S2**), whereas neither *Quercus* root exudates (“SG + *Quercus* exudates”) nor oak physical presence (“mixed-grown”) had any effect on *Molinia* RI. However, a significantly higher *Molinia* RI was estimated between SG + *Molinia* exudates and MG ($p = 0.04$).

Oak root/shoot ratio (R/S) was unaffected by application of either root exudate (**Figure 3, Table S3 and S4**). By contrast, moor grass R/S ratio was dramatically reduced when mixed-grown with oak ($p = 0.02$, **Figure 3, Table S3 and S4**).

Response of oak root and shoot biomass, relative biomass increment and root/shoot ratio to Molinia root exudate application (one-way experiment)

In the one-way experiment, root dry weight of oak seedlings was significantly lower after weekly supply of *Molinia* root exudates ($p = 0.03$, **Figure 4 and Table S5**), whereas shoot dry weight reduction in response to *Molinia* root exudates was only marginally significant ($p = 0.06$, **Figure 4 and Table S5**). Root dry weight was reduced by 50% and shoot dry weight by 44% by *Molinia* exudates. No significant difference was observed in relative biomass increment (**Figure 5 and Table S5**) or in R/S ratio (**Figure 6 and Table S5**).

Discussion

Our results show that oak-moor grass interactions are more complex than hitherto described. Competition by exploitation is not the only explanation for the deleterious effect of moor grass on oak growth. *Molinia* root exudates generally had a negative effect on oak growth, especially on roots, suggesting involvement of allelochemicals. Another interesting result was the strong decrease in the moor grass R/S ratio in the presence of oak, revealing a change in biomass allocation. A higher biomass allocation in shoot purple moor grass without necessity to invest in the root compartment suggested that oak facilitated grass growth. Conversely, root exudates of *Quercus* had no significant effect on moor grass but they tended to have a positive intraspecific effect.

As reported by numerous studies in natural conditions, light competition occurs simultaneously with belowground resource and interference competition. Even if light competition by perennial grasses is often limited as highlighted in the introduction, we recognize that the net outcome of competition *in situ* would be the integration of light and belowground competition processes. However, the study and the following discussion focused on the allelopathic effect. Accordingly, light competition was removed from the experimental design to understand in more details the belowground allelopathic effect. Our aim was not to reproduce natural conditions but rather to lean on belowground processes. A more suitable design would be necessary to disentangle the combined contribution of light, belowground resource competition and allelopathy in the field.

The two-way experiment showed a consistent lower oak RI when mixed-grown with moor grass, and to a lesser extent after application of *Molinia* exudate solution. This is consistent with the simultaneous expression of the two types of competition, namely resource exploitation and chemical interference, respectively. When mixed-grown with moor grass, oak underwent both processes, whereas when sole-grown and watered with the *Molinia* exudate solution, oak was subjected only to allelopathic effects. On the assumption, which remains to be thoroughly tested, that effects of interference and competition by resource exploitation do not interact and can be distinguished from each other (Inderjit and Callaway, 2003; Inderjit and Del Moral, 1997; Uddin and Robinson, 2017; Weidenhamer, 2006), the difference between oak SG, SG + *Molinia* exudates and MG gave the relative contributions of the two competition types. Mixed-grown oak RI showed a 50% decrease compared to sole-grown, whereas allelopathy (SG + *Molinia* exudates) was responsible for a 17% decrease, although not statistically significant. Considering that allelopathy and resource competition are independent of each other, have additive effects, and the nature and quantity of root exudates were the same than in maceration, this means that resource competition could be responsible for about a $50\% - 17\% = 33\%$ decrease. This result suggested that in our pot experiment, resource exploitation had a greater effect than the allelopathic pathway. These results were consistent with other studies working on interactions between *P. sylvestris* and *E. hermaphroditum* (Nilsson et al., 1993) or between *Q. pubescens* and *P. halepensis* (Fernandez et al., 2016). They demonstrated that *P. sylvestris* and *Q. pubescens* biomass were more strongly reduced by a combination of allelopathy and competition than allelopathy alone. Moreover, treatments altered biomass allocation of *Quercus pubescens* with a lower R/S ratio in the case of allelopathy (Fernandez et al., 2016). Our results did not support this effect, oak R/S being non-

significantly affected by the treatment although the root system was more strongly altered than the shoot system, probably owing to a high variability of responses.

Root exudates of oak favour its own growth, but exudates of Molinia inhibit its own growth

Quercus root exudates tended to favour oak biomass but had no effect on moor grass biomass. This result was surprising because previous studies using different species compositions demonstrated a facilitating effect in interspecific interactions and not in intraspecific interactions. For example, Li et al. (2016) evidenced that root exudates of maize promoted N₂ fixation of faba bean (*Vicia faba* L.), whereas to our knowledge, whether root exudates can facilitate growth of conspecific species is still not known. Further studies should analyse the effects of species physical presence on the nature and intensity of root exudates rather than root exudates of sole-grown species. By contrast, in the present study, moor grass biomass was not significantly affected by *Quercus* root exudates. This is at variance with some studies reporting an allelopathic effect of *Quercus* sp from different regions on understorey species (Li et al., 2007). Callaway et al. (1991) identified an allelopathic property of adult *Quercus* root exudates that inhibited understorey productivity in woodlands. Alrababah et al. (2009) and Hashoum et al. (2017) demonstrated that aqueous extract of green and senescent mature *Q. coccifera* and *Q. pubescens* leaves and litter contained allelochemicals that reduced seed germination and controlled the dynamics of the herbaceous species. In our study, we were interested in the regeneration phase of the oak. At this stage young seedlings are characterized by small below- and aboveground biomasses, making them more vulnerable to grass understorey, and their allelochemical profile is probably different from that of a mature tree. *Molinia* own root exudates tended to decrease its total relative dry weight increment. Auto-allelopathy is a recognised mechanism in various plant species (Jamshidi et al., 2011; Mafeo and Mashela, 2010; Yu et al., 2003) that can lead to community stability by tempering the establishment of a single species. Specificity and intensity of plant interactions seem difficult to predict and probably depend on the species and on its biotic and abiotic environment.

As reported elsewhere (Vernay et al., 2018a, 2018b) grass dry weight was higher when mixed-grown with oak, although the difference was marginally significant in the present study. The most remarkable effect was recorded on the moor grass R/S ratio in the mixed-grown condition, which was significantly lower. Preferential biomass allocation to shoots can be interpreted as a strategy of a fast-growing species associated with a need to meet its C requirements when N availability allows it (Mardanov et al., 1998). Sufficient N availability

could be attributed to oak rhizodeposition, which provided nitrogen and facilitates moor grass shoot growth (Fernandez et al., 2020).

Molinia root exudates inhibit oak growth

Biomass analysis in our “one-way” experiment suggested that *Molinia* root exudates might contain chemicals that inhibit oak seedling growth. Biomass analysis of both root and shoot highlighted that *Molinia* root exudates had a greater effect on oak roots than shoots, as also observed in Nilsson's study (1994). This decrease was also observed in the relative total dry weight increment and R/S ratio, though not significantly, probably because of wide variabilities. However, the allelopathy effect on oak revealed a temporal variability, decreased growth appearing more marginal in our “two-way experiment” in 2018 than in our “one-way experiment” in 2017.

The one-way experiment focusing specifically on each organ allowed a better understanding of the action mechanisms of *Molinia* root exudates: inhibiting oak root growth, root exudates participated in strengthening competition for resources in favour of moor grass. Such inhibition of the oak root system is likely to have a negative impact on shoot growth in the following spring.

Zhang et al. (2020) demonstrated through a recent meta-analysis that allelopathy can reduce plant performance by 25% on average. In the “one-way” experiment, oak shoot and root biomasses were diminished by 47% and 50%, respectively, with *Molinia* root exudates. RI was reduced by 31% in the “one-way” experiment and to a lesser extent by 17% in the “two-way” experiment. Also, R/S ratio was decreased by 9.7% and 3.8% in the “one-way” and “two-way” experiments, respectively. In our study, we observed an annual variability of the reducing effect and a different intensity of the allelopathic effect in the considered organ. However, the mean of all these reductions due to allelopathy was 26%, which lies in the value range reviewed by Zhang et al. (2020)

Nature of root exudates

No specialized metabolite was evidenced in the aqueous exudates, although the difference in chemical composition of *Molinia* and *Quercus* was confirmed (Figure S1). To our knowledge, studies on root exudates are rare, as extraction from root extracts obtained after root grinding are commonly favoured. However, this did not account for what was actually released by the roots, since all root substances were supplied, including those that would not necessarily

be released into the soil. Root exudates were obtained by an artificial method (maceration) giving a proxy of root exudates, but this is a reliable method if root exudates could not be extracted from the soil (Oburger and Jones, 2018).

Despite a low concentration of root exudates in the analysed solution, continuous supply interestingly resulted in an inhibition of growth in oak along the monitored period. This striking result suggested a strong allelopathic effect of *Molinia* roots (SG + *Molinia* exudates). Moreover, the swift response suggested that allelochemicals likely originated from root exudation by moor grass roots rather than from release by moor grass root decomposition.

Perspectives

Overall, it is noteworthy that different chemicals originating from root aqueous maceration of different species can have positive or negative effects depending on that species and on the target species. Contrasting effects of root exudates of *Quercus* and *Molinia* on each other highlighted the absolute necessity to consider the “two-way” analysis in plant interactions, as well as the intraspecific interactions. Our experiment was conducted in pots so that we could test such effects *in vivo*. Although many substances are present in the root exudates, we did not identify them in this study. Whereas untargeted metabolomics has been successfully applied to profile root exudates of several species (van Dam and Bouwmeester, 2016), only very few compounds were identified (often less than 10%) and most of these were primary compounds (such as sugars or amino acids). Identifying specialized compounds in our root exudates would require a comprehensive study focused on the composition of exudates, needing a combination of platforms to succeed. Substances indicated by statistical analysis could be further identified, either by applying MS/MS on peaks in the extract, or by isolating the substance for additional structural analysis using NMR (Oburger and Jones, 2018; van Dam and Bouwmeester, 2016).

Another important point to consider are soil altering allelopathic effect *via* modifications of biotic and abiotic soil parameters. Root exudates are known to influence nutrient availability and physicochemical soil conditions and reciprocally (Mohammadkhani and Servati, 2018; Zeng, 2014). By altering the soil properties, it is possible that moor grass allelochemicals may have indirectly affected oak growth. We could not exclude this effect, but results suggested that moor grass exudate supply to SG moor grass did not significantly reduce moor grass RI.

Our study highlights the important role of interference processes in oak-grass interaction. Further research should pursue the characterization of the nature and quantity of

exudates (Oburger and Jones, 2018), given their critical role in plant-plant interaction dynamics (Ehlers et al., 2020).

From a forest management perspective, and specifically in relation to sessile oak regeneration, this study confirms that forest perennial grasses are detrimental to seedling establishment. The allelopathic pathway is one of the processes involved in this inhibition but it seems less important than competition by resource exploitation during the period of growth. Nevertheless, production of allelochemicals contributes to reduced oak performance (growth, soil exploration, efficiency in nutrient acquisition), accentuating competition for resources. Targets of allelopathic substances could be microbial communities associated with oak roots including mycorrhizae, but extensive studies are still needed to address this question. The density and proximity of moor grass around young oak should be reduced by various means to avoid allelopathic effects, which will also reduce resource exploitation. One solution would be to act on light availability in the understorey to modulate the presence and intensity of the competitive processes of moor grass (Gaudio et al., 2011). The decoupling of exploitation *vs* interference interaction may not be straightforward (Inderjit and Del Moral, 1997) and the potential indirect facilitative effect (Michalet et al., 2015; Siemann and Rogers, 2003) from mature trees could increase resource availability for seedlings and mitigate the production of allelopathic substances from understorey species.

Acknowledgements

UHPLC-MS was performed on the IMBE Support Service on Chemical Ecology and Metabolomics (funded by the CNRS, the Provence Alpes Côte d'Azur Region, the TOTAL Foundation and ANR) particularly the MALLABAR platform. The authors thank Stéphane Greff for its valuable help in metabolite extraction and chemical analyses of root exudates.

416 *Appendix 1*

417 *Metabolite extraction*

418 *Molinia* root exudate composition was determined with an UHPLC instrument (Dionex
419 Ultimate 3000 equipped with an RS Pump, an autosampler, a thermostated column
420 compartment and a UV diode array, Thermo Scientific, USA) coupled to an accurate mass
421 spectrometer (qToF) equipped with an ESI source (Impact II, Bruker Daltonics, Germany).
422 UHPLC separation was done on an Acclaim C18 column (2.1 mm × 150 mm, 2.2 µm, Thermo
423 Scientific, USA). Elution rate was set to 0.5 mL min⁻¹ at a constant temperature of 45°C. A
424 pooled sample combining 10 µL of each sample was used to determine the chromatographic
425 method and the injection volume. This pooled sample was also used as a quality control.

426 Injection was set to 10 µL after twofold dilution of all exudates with the same solvent as that
427 used for extraction. Chromatographic solvents were composed of A: water with 0.1% formic
428 acid and B: acetonitrile with the same additive. The chromatographic gradient was set as
429 follows: 2% of B for 2 min, then increase in the proportion of solvent B to reach 100% at
430 14 min. Each analysis was followed by a column cleaning phase with 100% B for 3 min, and
431 column equilibration for 3 min, giving a total runtime of 20 min. Samples of each condition
432 were randomly injected to integrate time-dependent changes in UHPLC-MS chromatographic
433 fingerprints. Pooled samples, injected at the beginning, at the end and every six samples, were
434 used for ion intensity normalization. Blanks were also injected to remove background signals.

435 MS parameters were set as follows: nebulizer gas, N₂ at 51 psi; dry gas, N₂ at 12 L min⁻¹,
436 capillary temperature 200°C and voltage 3000 V. The mass spectrometer was systematically
437 calibrated with formate/acetate solution forming clusters on the studied mass range before a full
438 set of analyses. The same calibration solution was automatically injected before each sample
439 for internal mass calibration. For the mass spectra, tests were performed in both negative and
440 positive modes. Mass spectra were in negative ionization mode in full scan mode from 50 to
441 1200 amu at 2 Hz. DDA-MS2 analyses were performed on three major ions detected at each
442 scan on a pooled sample for metabolite annotation.

443 *Root exudates*

444 Analyses were automatically recalibrated using internal calibration, ensuring a precision of *m/z*
445 below 2 ppm on the mass range, before exporting data in netCDF files (centroid mode) using
446 Bruker Compass DataAnalysis 4.3. Analysis files were then processed using the XCMS

447 package (Smith et al. 2006) of R software, using the different steps necessary to generate the
448 final data matrix: (i) peak picking for detection of different features, (ii) retention time
449 correction ("obiwarp" method), (iii) grouping, (iv) filling of peaks to integrate portions where
450 peaks were initially absent and (v) report and data matrix generation transferral to Excel. Each
451 individual ion of each analysis was then normalized according to the injection order as
452 described by Van Der Kloet et al. (2009). After the data set normalization, around 6000 features
453 were kept before the filtering steps. To ensure data quality and remove redundant signals, three
454 successive filtering steps were applied to pre-processed data using an in-house script on R. The
455 first was based on the signal/noise (S/N) ratio to remove signals observed in blanks (S/N set at
456 10 for features matching between pooled samples and blanks). The second allowed suppression
457 of signals based on the value of the coefficient of variation of ion intensities in pooled samples
458 (threshold at 0.4). The last step consisted in the deletion of all auto-correlated ions (threshold
459 at 0.8) to remove isotopes and adducts.

References

- Alrababah, M.A., Tadros, M.J., Samarah, N.H., Ghosheh, H., 2009. Allelopathic effects of *Pinus halepensis* and *Quercus coccifera* on the germination of Mediterranean crop seeds. *New For.* 38, 261–272. <https://doi.org/10.1007/s11056-009-9145-8>
- Balandier, P., Collet, C., Miller, J.H., Reynolds, P.E., Zedaker, S.M., 2006. Designing forest vegetation management strategies based on the mechanisms and dynamics of crop tree competition by neighbouring vegetation. *Forestry* 79, 3–27. <https://doi.org/10.1093/forestry/cpi056>
- Batish, D.R., Lavanya, K., Pal Singh, H., Kohli, R.K., 2007. Root-mediated allelopathic interference of nettle-leaved goosefoot (*Chenopodium murale*) on wheat (*Triticum aestivum*). *J. Agron. Crop Sci.* 193, 37–44. <https://doi.org/10.1111/j.1439-037X.2006.00243.x>
- Becker, M., Lévy, G., 1982. Le dépérissement du chêne en forêt de Tronçais : les causes écologiques. *Ann. des Sci. For.* 39, 439–444.
- Ben-Hammouda, M., Ghorbal, H., Kremer, R.J., Oueslatt, O., 2002. Autotoxicity of barley. *J. Plant Nutr.* 25, 1155–1161. <https://doi.org/10.1081/PLN-120004379>
- Bertin, C., Yang, X., Weston, L.A., 2003. The role of root exudates and allelochemicals in the rhizosphere. *Plant Soil* 256, 67–83. <https://doi.org/10.1023/A:1026290508166>
- Böttger, A., Vothknecht, U., Bolle, C., Wolf, A., 2018. Plant secondary metabolites and their general function in plants, in: *Lessons on Caffeine, Cannabis & Co.* pp. 3–17. <https://doi.org/10.1007/978-3-319-99546-5>
- Callaway, R.M., Nadkarni, N.M., Mahall, B.E., 1991. Facilitation and interference of *Quercus Douglasii* on understory productivity in Central California. *Ecology* 72, 1484–1499.
- Casper, B.B., Jackson, R.B., 1997. Plant competition underground. *Annu. Rev. Ecol. Syst.* 28, 545–570. <https://doi.org/10.1146/annurev.ecolsys.28.1.545>
- Coll, L., Balandier, P., Picon-Cochard, C., Prévosto, B., Curt, T., 2003. Competition for water between beech seedlings and surrounding vegetation in different light and vegetation composition conditions. *Ann. For. Sci.* 60, 593–600. <https://doi.org/10.1051/forest:2003051>
- Coomes, D.A., Grubb, P.J., 2000. Impacts of root competition in forests and woodlands: a theoretical framework and review of experiments. *Ecol. Monogr.* 70, 171–207.
- Davies, D.A., 1987. Long-term effects of improvement methods on *Molinia caerulea* dominant rough grazing on wet hill land. 2. Mineral composition of herbage and soil physical, chemical and biological characteristics. *J. Agric. Sci.* 109, 243. <https://doi.org/10.1017/S0021859600080655>
- Davis, M.A., Wrage, K.J., Reich, P.B., 1998. Competition between tree seedlings and herbaceous vegetation: support for a theory of resource supply and demand. *J. Ecol.* 86, 652–661. <https://doi.org/10.1046/j.1365-2745.1998.00087.x>
- Effah, E., Holopainen, J.K., McCormick, A.C., 2019. Potential roles of volatile organic compounds in plant competition. *Perspect. Plant Ecol. Evol. Syst.* 38, 58–63. <https://doi.org/10.1016/j.ppees.2019.04.003>

501 Ehlers, B.K., Berg, M.P., Staudt, M., Holmstrup, M., Glasius, M., Ellers, J., Tomiolo, S.,
502 Madsen, R.B., Slotsbo, S., Penuelas, J., 2020. Plant Secondary Compounds in Soil and
503 Their Role in Belowground Species Interactions. *Trends Ecol. Evol.* 35, 716–730.
504 <https://doi.org/10.1016/j.tree.2020.04.001>

505 Fernandez, C., Lelong, B., Vila, B., Mévy, J.P., Robles, C., Greff, S., Dupouyet, S., Bousquet-
506 Mélou, A., 2006. Potential allelopathic effect of *Pinus halepensis* in the secondary
507 succession: an experimental approach. *Chemoecology* 16, 97–105.
508 <https://doi.org/10.1007/s00049-006-0334-z>

509 Fernandez, C., Monnier, Y., Santonja, M., Gallet, C., Weston, L.A., Prévosto, B., Saunier, A.,
510 Baldy, V., Bousquet-Mélou, A., 2016. The impact of competition and allelopathy on the
511 trade-off between plant defense and growth in two contrasting tree species. *Front. Plant*
512 *Sci.* 7, 1–14. <https://doi.org/10.3389/fpls.2016.00594>

513 Fernandez, M., Malagoli, P., Vernay, A., Améglio, T., Balandier, P., 2020. Below-ground
514 nitrogen transfer from oak seedlings facilitates *Molinia* growth : ¹⁵N pulse-chase
515 labelling. *Plant Soil* 449, 343–356. <https://doi.org/10.1007/s11104-020-04473-9>

516 Fisher, R.F., 1987. Allelopathy: a potential cause of forest regeneration failure. *Allelochem.*
517 *Role Agric. For.* 176–184. <https://doi.org/doi:10.1021/bk-1987-0330.ch016>

518 Foster, B.L., 1999. Establishment, competition and the distribution of native grasses among
519 Michigan old-fields. *J. Ecol.* 87, 476–489.

520 Friedman, J., Waller, G.R., 1985. Allelopathy and autotoxicity. *Trends Biochem. Sci.* 10, 47–
521 50. [https://doi.org/10.1016/0968-0004\(85\)90224-5](https://doi.org/10.1016/0968-0004(85)90224-5)

522 Fuerst, E.P., Putnam, A.R., 1983. Separating the competitive and allelopathic components of
523 interference - Theoretical principles. *J. Chem. Ecol.* 9, 937–944.
524 <https://doi.org/10.1007/BF00982203>

525 Gallet, C., Pellissier, F., 2002. Interactions allélopathiques en milieu forestier. *Rev. For.*
526 *Française* 6, 567–576.

527 Gaudio, N., Balandier, P., Philippe, G., Dumas, Y., Jean, F., Ginisty, C., 2011. Light-
528 mediated influence of three understorey species (*Calluna vulgaris*, *Pteridium aquilinum*,
529 *Molinia caerulea*) on the growth of *Pinus sylvestris* seedlings. *Eur. J. For. Res.* 130, 77–
530 89. <https://doi.org/10.1007/s10342-010-0403-2>

531 Grime, J.P., 1974. Vegetation classification by reference to strategies. *Nature* 250, 26–31.
532 <https://doi.org/10.1038/252497a0>

533 Guretzky, J.A., 2020. Plant Interactions, in: *Forages: The Science of Grassland Agriculture*.
534 pp. 187–199.

535 Hao, W., Ren, L., Ran, W., Shen, Q.R., 2010. Allelopathic effects of root exudates from
536 watermelon and rice plants on *Fusarium oxysporum* f.sp. *niveum*. *Plant Soil* 336, 485–
537 497. <https://doi.org/10.1007/s11104-010-0505-0>

538 Hashoum, H., Santonja, M., Gauquelin, T., Saatkamp, A., Gavinet, J., Greff, S., Lecareux, C.,
539 Fernandez, C., Bousquet-Mélou, A., 2017. Biotic interactions in a Mediterranean oak
540 forest: role of allelopathy along phenological development of woody species. *Eur. J. For.*
541 *Res.* 136, 699–710. <https://doi.org/10.1007/s10342-017-1066-z>

542 Huang, L., Song, L., Xia, X., 2013. Plant-soil feedbacks and soil sickness: from mechanisms

543 to application in agriculture. J. Chem. Ecol. 39, 232–242.
544 <https://doi.org/10.1007/s10886-013-0244-9>

545 Inderjit, Callaway, R.M., 2003. Experimental designs for the study of allelopathy. Plant Soil
546 256, 1–11. <https://doi.org/10.1023/A:1026242418333>

547 Inderjit, Del Moral, R., 1997. Is separating resource competition from allelopathy realistic?
548 Bot. Rev. 63, 221–230. <https://doi.org/10.1007/BF02857949>

549 Inderjit, Mallik, A., 2002. Can *Kalmia angustifolia* interference to black spruce (*Picea*
550 *mariana*) be explained by allelopathy? For. Ecol. Manage. 160, 75–84.
551 [https://doi.org/10.1016/S0378-1127\(01\)00463-7](https://doi.org/10.1016/S0378-1127(01)00463-7)

552 Inderjit, Mallik, A.U., 1996. The nature of interference potential of *Kalmia angustifolia*. Can.
553 J. For. Res. 26, 1899–1904.

554 Jamshidi, S., Hashemizadeh, S., Shahrokhi, S., 2011. Assessment of auto-allelopathic potential
555 of broomcorn (*Sorghum vulgare* var. *technicum*) 13, 2–6.

556 Kato-Noguchi, H., 2004. Allelopathic substance in rice root exudates: Rediscovery of
557 momilactone B as an allelochemical. J. Plant Physiol. 161, 271–276.
558 <https://doi.org/10.1078/0176-1617-01188>

559 Li, B., Yu-Ying, L., Wu, H.-M., Zhang, F.-F., Li, C.-J., Li, X.-X., Lambers, H., Li, L., 2016.
560 Root exudates drive interspecific facilitation by enhancing nodulation and N₂ fixation.
561 Proc. Natl. Acad. Sci. 113, 6496–6501. <https://doi.org/10.1073/pnas.1523580113>

562 Li, G. De, Jia, L.M., Li, X.W., 2007. Research advances in allelopathy of *Quercus* L. For.
563 Stud. China 9, 287–294. <https://doi.org/10.1007/s11632-007-0046-7>

564 Mafeo, T.P., Mashela, P.W., 2010. Allelopathic inhibition of seedling emergence in
565 dicotyledonous crops by *Cucumis* bio-nematicide. African J. Biotechnol. 9, 8349–8354.
566 <https://doi.org/10.5897/AJB10.652>

567 Mallik, A.U., 2003. Conifer regeneration problems in boreal and temperate forests with
568 ericaceous understory: role of disturbance, seedbed limitation, and keystone species
569 change. CRC. Crit. Rev. Plant Sci. 22, 341–366.

570 Mallik, A.U., Biswas, S.R., Collier, L.C.S., 2016. Belowground interactions between *Kalmia*
571 *angustifolia* and *Picea mariana*: roles of competition, root exudates and ectomycorrhizal
572 association. Plant Soil 403, 471–483. <https://doi.org/10.1007/s11104-016-2819-z>

573 Mardanov, A., Samedovam, A., Shirvany, T., 1998. Root-shoot relationships in plant
574 adaptation to nitrogen deficiency 147–154.

575 Michalet, R., Chen, S.Y., An, L.Z., Wang, X.T., Wang, Y.X., Guo, P., Ding, C.C., Xiao, S.,
576 2015. Communities: are they groups of hidden interactions? J. Veg. Sci. 26, 1–12.
577 <https://doi.org/10.1111/jvs.12226>

578 Michalet, R., Maalouf, J., Hayek, P. Al, 2017. Direct litter interference and indirect soil
579 competitive effects of two contrasting phenotypes of a spiny legume shrub drive the forb
580 composition of an oromediterranean community. Oikos 126, 1090–1100.
581 <https://doi.org/10.1111/oik.03800>

582 Michalet, R., Xiao, S., Cavieres, L.A., Ragan, M., 2011. Phenotypic variation in nurse traits
583 and community feedbacks define an alpine community. Ecol. Lett. 14, 433–443.
584 <https://doi.org/10.1111/j.1461-0248.2011.01605.x>

585 Mohammadkhani, N., Servati, M., 2018. Nutrient concentration in wheat and soil under
586 allelopathy treatments. J. Plant Res. 131, 143–155. [https://doi.org/10.1007/s10265-017-](https://doi.org/10.1007/s10265-017-0981-x)
587 0981-x

588 Nilsson, M.-C., Högborg, P., Zackrisson, O., Fengyou, W., 1993. Allelopathic effects by
589 *Empetrum hermaphroditum* on development and nitrogen uptake by roots and
590 mycorrhizae of *Pinus silvestris*. Can. J. Bot. 71, 620–628. [https://doi.org/10.1139/b93-](https://doi.org/10.1139/b93-071)
591 071

592 Nilsson, M.C., 1994. Separation of allelopathy and resource competition by the boreal dwarf
593 shrub *Empetrum hermaphroditum* Hagerup. Oecologia 98, 1–7.
594 <https://doi.org/10.1007/BF00326083>

595 Oburger, E., Jones, D.L., 2018. Sampling root exudates – Mission impossible? Rhizosphere 6,
596 116–133. <https://doi.org/10.1016/j.rhisph.2018.06.004>

597 Olofsdotter, M., Navarez, D., Rebulanan, M., Streibig, J.C., 1999. Weed-suppressing rice
598 cultivars - Does allelopathy play a role? Weed Res. 39, 441–454.
599 <https://doi.org/10.1046/j.1365-3180.1999.00159.x>

600 Pagès, J.-P., Michalet, R., 2003. A test of the indirect facilitation model in a temperate
601 hardwood forest of the Northern French Alps. J. Ecol. 91, 932–940.

602 Penuelas, J., Llusia, J., 1998. Influence of intra- and inter-specific interference on terpene
603 emission by *Pinus halepensis* and *Quercus ilex* seedlings. Biol. Plant. 41, 139–143.

604 Pérez, F.J., Ormeno-Nuñez, J., 1991. Difference in hydroxamic acid content in roots and root
605 exudates of wheat (*Triticum aestivum* L.) and rye (*Secale cereale* L.): Possible role in
606 allelopathy. J. Chem. Ecol. 17, 1037–1043. <https://doi.org/10.1007/BF01402932>

607 Pérez, F.J., Ormeño-Nuñez, J., 1991. Root exudates of wild oats: allelopathic effect on spring
608 wheat. Phytochemistry 30, 2199–2202. [https://doi.org/10.1016/0031-9422\(91\)83614-Q](https://doi.org/10.1016/0031-9422(91)83614-Q)

609 Picon-Cochard, C., Coll, L., Balandier, P., 2006. The role of below-ground competition
610 during early stages of secondary succession: the case of 3-year-old Scots pine (*Pinus*
611 *sylvestris* L.) seedlings in an abandoned grassland. Oecologia 148, 373–383.
612 <https://doi.org/10.1007/s00442-006-0379-2>

613 Pistón, N., Michalet, R., Schöb, C., Macek, P., Armas, C., Pugnaire, F.I., 2018. The balance
614 of canopy and soil effects determines intraspecific differences in foundation species’
615 effects on associated plants. Funct. Ecol. 32, 2253–2263. [https://doi.org/10.1111/1365-](https://doi.org/10.1111/1365-2435.13139)
616 2435.13139

617 Reigosa, M.J., Sánchez-moreiras, A., González, L., 1999. Ecophysiological approach in
618 allelopathy. Crit. Rev. Biotechnol. 18, 577–608.
619 <https://doi.org/10.1080/07352689991309405>

620 Rice, E.L., 1984. Allelopathy, in: Rice, E.L. (Ed.), . Academic Press, Orlando.
621 <https://doi.org/https://doi.org/10.1016/B978-0-08-092539-4.50019-8>

622 Ridenour, W.M., Callaway, R.M., 2001. The relative importance of allelopathy in
623 interference: the effects of an invasive weed on a native bunchgrass. Oecologia 126,
624 444–450. <https://doi.org/10.1007/s004420000533>

625 Santonja, M., Bousquet- Mélou, A., Greff, S., Ormeño, E., Fernandez, C., 2019. Allelopathic
626 effects of volatile organic compounds released from *Pinus halepensis* needles and roots.

627 Ecol. Evol. 8201–8213. <https://doi.org/10.1002/ece3.5390>

628 Schenk, H.J., 2006. Root competition: beyond resource depletion. J. Ecol.
629 <https://doi.org/10.1111/j.1365-2745.2006.01124.x>

630 Schöb, C., Callaway, R.M., Anthelme, F., Brooker, R.W., Cavieres, L.A., Kikvidze, Z.,
631 Lortie, C.J., Michalet, R., Pugnaire, F.I., Xiao, S., Cranston, B.H., García, M.C., Hupp,
632 N.R., Llambí, L.D., Lingua, E., Reid, A.M., Zhao, L., Butterfield, B.J., 2014. The
633 context dependence of beneficiary feedback effects on benefactors in plant facilitation. J.
634 Physiol. 204, 386–396. <https://doi.org/10.1111/nph.12908>

635 Schöb, C., Prieto, I., Armas, C., Pugnaire, F.I., 2013. Consequences of facilitation: one plant's
636 benefit is another plant's cost. Funct. Ecol. 28, 500–508. [https://doi.org/10.1111/1365-](https://doi.org/10.1111/1365-2435.12185)
637 2435.12185

638 Siemann, E., Rogers, W.E., 2003. Changes in light and nitrogen availability under pioneer
639 trees may indirectly facilitate tree invasions of grasslands. J. Ecol. 91, 923–931.
640 <https://doi.org/10.1046/j.1365-2745.2003.00822.x>

641 Singh, H.P., Batish, D.R., Kohli, R.K., 1999. Autotoxicity: Concept, Organisms, and
642 Ecological Significance. CRC. Crit. Rev. Plant Sci. 18, 757–772.
643 <https://doi.org/10.1080/07352689991309478>

644 South, D.B., 1995. Relative Growth Rate: a critique. South African For. J. 173, 1–6.

645 Souto, X.C., Gonzales, L., Reigosa, M.J., 1994. Comparative analysis of allelopathic effects
646 produced by four forestry species during decomposition process in their soils in Galicia
647 (NW Spain). J. Chem. Ecol. 20, 3005–3015. <https://doi.org/10.1007/BF02098405>

648 Taylor, K., Rowland, A.P., Jones, H.E., 2001. *Molinia caerulea* (L.) Moench. J. Ecol. 89,
649 126–144. <https://doi.org/10.1046/j.1365-2745.2001.00534.x>

650 Tilman, D., 1990. Constraints and tradeoffs: toward a predictive theory of competition and
651 succession. Oikos 58, 3–15. <https://doi.org/10.2307/3565355>

652 Timbal, J., Gelpe, J., Garbaye, J., Courrier, G., 1990. Étude préliminaire sur l'effet dépressif
653 de la molinie (*Molinia caerulea*) sur la croissance et l'état mycorhizien de semis de chêne
654 rouge (*Quercus rubra*). Ann. For. Sci. 21, 643–649.

655 Tsunoda, T., van Dam, N.M., 2017. Root chemical traits and their roles in belowground biotic
656 interactions. Pedobiologia (Jena). 65, 58–67.
657 <https://doi.org/10.1016/j.pedobi.2017.05.007>

658 Uddin, M.N., Robinson, R.W., 2017. Allelopathy and resource competition: the effects of
659 *Phragmites australis* invasion in plant communities. Bot. Stud. 58.
660 <https://doi.org/10.1186/s40529-017-0183-9>

661 van Dam, N.M., Bouwmeester, H.J., 2016. Metabolomics in the Rhizosphere: Tapping into
662 Belowground Chemical Communication. Trends Plant Sci. 21, 256–265.
663 <https://doi.org/10.1016/j.tplants.2016.01.008>

664 Vernay, A., Balandier, P., Guinard, L., Améglio, T., Malagoli, P., 2016. Photosynthesis
665 capacity of *Quercus petraea* (Matt.) saplings is affected by *Molinia caerulea* (L.) under
666 high irradiance. For. Ecol. Manage. 376, 107–117.
667 <https://doi.org/10.1016/j.foreco.2016.05.045>

668 Vernay, A., Malagoli, P., Fernandez, M., Perot, T., Améglio, T., Balandier, P., 2018a.

- Improved *Deschampsia cespitosa* growth by nitrogen fertilization jeopardizes *Quercus petraea* regeneration through intensification of competition. *Basic Appl. Ecol.* 31, 21–32. <https://doi.org/10.1016/j.baae.2018.06.002>
- Vernay, A., Malagoli, P., Fernandez, M., Perot, T., Améglio, T., Balandier, P., 2018b. Carry-over benefit of high internal N pool on growth and function of oak seedlings (*Quercus petraea*) competing with *Deschampsia cespitosa*. *For. Ecol. Manage.* 419–420, 130–138. <https://doi.org/10.1016/j.foreco.2018.03.039>
- Viard-Crétat, F., Baptist, F., Secher-Fromell, H., Gallet, C., 2012. The allelopathic effects of *Festuca paniculata* depend on competition in subalpine grasslands. *Plant Ecol.* 213, 1963–1973. <https://doi.org/10.1007/s11258-012-0143-0>
- Weidenhamer, J.D., 2006. Distinguishing allelopathy from resource competition: The role of density. *Allelopath. A Physiol. Process with Ecol. Implic.* 85–103. https://doi.org/10.1007/1-4020-4280-9_4
- Wilson, S.D., Tilman, D., 1993. Plant competition and resource availability in response to disturbance and fertilization. *Ecology* 74, 599–611.
- Wu, H., Pratley, J., Lemerle, D., An, M., Liu, D.L., 2007. Autotoxicity of wheat (*Triticum aestivum* L.) as determined by laboratory bioassays. *Plant Soil* 296, 85–93. <https://doi.org/10.1007/s11104-007-9292-7>
- Xuan, T.D., Tawata, S., Khanh, T.D., Chung, I.M., 2005. Decomposition of allelopathic plants in soil. *J. Agron. Crop Sci.* 191, 162–171.
- Yu, J.Q., Matsui, Y., 1994. Phytotoxic substances in root exudates of cucumber (*Cucumis sativus* L.). *J. Chem. Ecol.* 20, 21–31. <https://doi.org/10.1007/BF02065988>
- Yu, J.Q., Ye, S.F., Zhang, M.F., Hu, W.H., 2003. Effects of root exudates and aqueous root extracts of cucumber (*Cucumis sativus*) and allelochemicals, on photosynthesis and antioxidant enzymes in cucumber. *Biochem. Syst. Ecol.* 31, 129–139. [https://doi.org/10.1016/S0305-1978\(02\)00150-3](https://doi.org/10.1016/S0305-1978(02)00150-3)
- Zélé, F., Magalhães, S., Kéfi, S., Duncan, A.B., 2018. Ecology and evolution of facilitation among symbionts. *Nat. Commun.* 9, 1–12. <https://doi.org/10.1038/s41467-018-06779-w>
- Zeng, R. Sen, 2014. Allelopathy - The solution is indirect. *J. Chem. Ecol.* 40, 515–516. <https://doi.org/10.1007/s10886-014-0464-7>
- Zhang, Z., Liu, Y., Yuan, L., Weber, E., van Kleunen, M., 2020. Effect of allelopathy on plant performance: a meta-analysis. *Ecol. Lett.* <https://doi.org/10.1111/ele.13627>

Figure 1. Experimental design of “two-way” experiment (A) and “one-way” experiment (B) with the different treatments tested on oak seedlings and *Molinia* tufts: sole-grown (SG), SG + *Quercus* exudates, SG + *Molinia* exudates, and mixed-grown (competition by exploitation + allelopathy).

Figure 2. Oak (a) and purple moor grass (b) relative dry biomass increment in the “two-way” experiment for the four different treatments: sole-grown (white bars), SG + *Quercus* exudates (light blue bars), SG + *Molinia* exudates (blue bars) and mixed-grown (dark blue bars). Values are reported as means $\pm$ SE (standard error, $n = 6$). Different letters indicate a significant difference between treatments.

Figure 3. Oak (a) and purple moor grass (b) root/shoot dry biomass ratio in the “two-way” experiment for the four different treatments: sole-grown (white bars), SG + *Quercus* exudates (light blue bars), SG + *Molinia* exudates (blue bars) and mixed-grown (dark blue bars). Values are reported as means $\pm$ SE ($n = 6$). Different letters indicate a significant difference between treatments.

Figure 4. Above- and belowground dry weights (g) of oak in the “one-way” experiment for the two different treatments: sole-grown (white bars) and SG + *Molinia* exudates (blue bars). Values are reported as means $\pm$ SE ($n = 8$). * corresponds to $p < 0.05$.

Figure 5. Oak relative dry biomass increment in the “one-way” experiment for the two different treatments: sole-grown (white bars) and SG + *Molinia* exudates (blue bars). Values are reported as means $\pm$ SE ($n = 8$).

Figure 6. Oak root/shoot dry biomass ratio in the “one-way” experiment for the two different treatments: sole-grown (white bars) and SG + *Molinia* exudates (blue bars). Values are reported as means $\pm$ SE ($n = 8$).

727 Appendix

728

729

Statistical tables

Abbreviations

Ctrl	Control
ME	<i>Molinia</i> root exudates
OE	Oak root exudates
Comp	Competition (exploitation competition + allelopathy)

Table S1. (Figure 2). Means $\pm$ SE of oak and moor grass relative dry biomass increment (two-way experiment) in sole-grown (SG), *Quercus* root exudates (SG + *Quercus* exudates), *Molinia* root exudates (SG + *Molinia* exudates) and mixed-grown (MG) treatment ($n = 6$).

	Oak				<i>Molinia</i>			
	SG	SG + <i>Q</i>	SG + <i>M</i>	MG	SG	SG + <i>Q</i>	SG + <i>M</i>	MG
Mean $\pm$ SE	1.71 $\pm$ 0.29	3.06 $\pm$ 0.60	1.42 $\pm$ 0.24	0.85 $\pm$ 0.19	30.47 $\pm$ 2.23	28.94 $\pm$ 5.25	23.54 $\pm$ 2.68	34.37 $\pm$ 3.68

Table S2. (Figure 2). Test statistic (F -value), statistical significance (p -value), and degrees of freedom (DF) assessing the effect of **treatment** on oak and moor grass relative dry biomass increment ($n = 6$).

		SG- SG+ <i>Q</i>	SG- SG+ <i>M</i>	SG- MG	SG+ <i>Q</i> - SG+ <i>M</i>	SG+ <i>Q</i> - MG	SG+ <i>M</i> - MG
Oak	p -value	0.08	0.46	0.04	0.04	0.01	0.09
	F -stat				6.61		
	DF				21		
<i>Molinia</i>	p -value	0.79	0.08	0.39	0.38	0.42	0.04
	F -stat				1.28		
	DF				20		

Table S3. (Figure 3). Means $\pm$ SE of oak and moor grass R/S ratio (two-way experiment) in sole-grown (SG), *Quercus* root exudates (SG + *Q*), *Molinia* root exudates (SG + *M*) and mixed-grown (MG) treatment ($n = 6$).

	Oak				Purple moor grass			
	SG	SG + <i>Q</i>	SG + <i>M</i>	MG	SG	SG + <i>Q</i>	SG + <i>M</i>	MG
Mean $\pm$ SE	0.80 $\pm$ 0.09	0.94 $\pm$ 0.06	0.77 $\pm$ 0.09	0.86 $\pm$ 0.14	1.58 $\pm$ 0.33	1.70 $\pm$ 0.56	1.64 $\pm$ 0.39	0.45 $\pm$ 0.06

Table S4. (Figure 3). Test statistic (F -value), statistical significance (p -value), and degrees of freedom (DF) assessing the effect of **treatment** on oak and moor grass R/S ratio ($n = 6$).

		SG- SG+ <i>Q</i>	SG- SG+ <i>M</i>	SG- MG	SG+ <i>Q</i> - SG+ <i>M</i>	SG+ <i>Q</i> - MG	SG+ <i>M</i> - MG
Oak	p -value	0.22	0.80	0.72	0.13	0.59	0.58
	F -stat						
	DF						
<i>Molinia</i>	p -value	0.86	0.91	0.02	0.93	0.07	0.02
	F -stat				2.35		
	DF				21		

Table S5. (Figure 4, 5 and 6). Means $\pm$ SE (g) of oak shoot and root dry weight (one-way experiment) in sole-grown (SG) and *Molinia* root exudates (SG + *M*) treatment. Test statistic (*t*-value), statistical significance (*p*-value), and degrees of freedom (DF) assessing the effect of **treatment** on shoot and root dry weight of oak (*n* = 8).

		Biomass		Relative increment		Root/shoot ratio	
		SG	SG+ <i>M</i>	SG	SG+ <i>M</i>	SG	SG+ <i>M</i>
Shoot	Mean	16.36 $\pm$ 2.78	9.37 $\pm$ 1.79	3.34 $\pm$ 1.58	2.29 $\pm$ 1.31	1.75 $\pm$ 0.16	1.58 $\pm$ 0.09
	<i>p</i> -value		0.06		0.17		0.35
	<i>t</i> -value		-2.11		1.44		-0.98
	DF		10.22		13.54		9.39
Root	Mean	28.84 $\pm$ 4.84	14.48 $\pm$ 2.43				
	<i>p</i> -value		0.03				
	<i>t</i> -value		-2.65				
	DF		8.84				

Figure 1

Figure 2. Two-way experiment

Figure 3. *Two-way experiment*

Figure 4. *One-way experiment*

Figure 5.
One-way experiment

Figure 6.
One-way experiment

Figure S1

Declaration of conflict of interest

None.

MF: Conceptualization; Data curation; Formal analysis; Methodology; Writing - original draft; review & editing.

PM, TA, PB: Conceptualization; Formal analysis; Funding acquisition; Methodology; Project administration; Supervision; Validation; Writing - review & editing.

CF, CG, AV: Formal analysis; Methodology; Software; Validation; Writing - review & editing.