



HAL
open science

Mapping the Interactions between *Escherichia coli* Tol Subunits

Xiang Y.-Z. Zhang, Emilie Goemaere, Rémi Thomé, Marthe Gavioli, E. Cascales, Roland Lloubés

► **To cite this version:**

Xiang Y.-Z. Zhang, Emilie Goemaere, Rémi Thomé, Marthe Gavioli, E. Cascales, et al.. Mapping the Interactions between *Escherichia coli* Tol Subunits. *Journal of Biological Chemistry*, 2009, 284 (7), pp.4275-4282. 10.1074/jbc.M805257200 . hal-03274816

HAL Id: hal-03274816

<https://amu.hal.science/hal-03274816>

Submitted on 14 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Mapping the Interactions between *Escherichia coli* Tol Subunits

ROTATION OF THE TolR TRANSMEMBRANE HELIX*

Received for publication, July 10, 2008, and in revised form, November 21, 2008. Published, JBC Papers in Press, December 15, 2008, DOI 10.1074/jbc.M805257200

Xiang Y.-Z. Zhang, Emilie L. Goemaere, Rémi Thomé, Marthe Gavioli, Eric Cascales, and Roland Lloubès¹

From the Laboratoire d'Ingénierie des Systèmes Macromoléculaires, Institut de Biologie Structurale et Microbiologie, Centre National de la Recherche Scientifique, UPR 9027, 13402 Marseille, France

The TolQRA proteins of *Escherichia coli* form an inner membrane complex involved in the maintenance of the outer membrane stability and in the late stages of cell division. The TolQR complex uses the proton motive force to regulate TolA conformation and its interaction with the outer membrane Pal lipoprotein. It has been proposed that an ion channel forms at the TolQR transmembrane helix (TMH) interface. This complex assembles with a minimal TolQ:TolR ratio of 4–6:2 and therefore involves 14–20 TMHs. To define the organization of the transmembrane helices in the membrane within the TolQR complex, we initiated a cysteine scanning study. In this study, we report results for the systematic replacement of each residue of the TolR TMH. Phenotypic analyses first showed that most of the mutants are functional. Three mutants, TolR L22C, D23C, and V24C, were shown to affect TolQR functioning. Disulfide bond complex formation further showed that two TolR anchors are close enough to interact. Two substitutions, L22C and V24C, form high level of dimers, suggesting that the TolR helix rotates as molecular gears between these two positions and that disulfide bond formation between these residues blocked the rotary motion. Mutations of critical residues located within the TolQ TMH2 and TMH3 and the TolR TMH and proposed to form the ion pathway prevent rotation between these two residues. TolR anchors may form molecular gears that oscillate in response to proton motive force to regulate channel activity.

The Tol proteins form a complex involved in the maintenance of cell envelope integrity (1, 2) and in the late stage of the cell division process (3). Three of these proteins, TolQ, TolR, and TolA interact through their transmembrane (TM)² segments in the inner membrane (4–8). In the periplasm, TolB,

the outer membrane-anchored peptidoglycan-associated lipoprotein Pal, and the C-terminal globular domain of TolA bind to each other to form a network linking the inner and outer membranes and the peptidoglycan layer (9–11). Previous studies showed that the TolQ, TolR, and TolA proteins form a complex with a 4–6:2:1 stoichiometry. The proton motive force (PMF), the TolQ and TolR proteins, and the TMH of TolA regulate the conformation of the periplasmic region of TolA and probably its subsequent interaction with Pal (10, 12, 13). The TolQ-R complex thus appears to form a molecular motor that converts chemical energy derived from the PMF to mechanical movements (12). This appears possible through the passage of ions in an aqueous lumen formed by the single TMH of TolR and two helices (TMH2 and TMH3) of TolQ (12, 14). Several residues, including the critical Asp²³ residue that lies within the TolR TM segment, have been shown to be necessary for the TolQR complex to function as a molecular motor. Because of the hydrophilic behavior of these residues, it has been suggested that they probably form the aqueous ion pathway. The proline residue at position 187 of TolQ (TMH3) is also essential and has been proposed to regulate conformational modifications of the TolQR complex in response to ion transit through the aqueous channel (14). Recently, a cysteine scanning study demonstrated that the C-terminal periplasmic domain of TolR forms head-to-tail homo-dimers (15), a result that has been confirmed by the NMR structure of the *Haemophilus influenzae* TolR periplasmic domain (16). Cysteine accessibility indicates that the same domain changes conformation in response to PMF and to critical residues suspected to form the ion pathway. Consequently, it has been proposed that the C-terminal domain of TolR may regulate ion entrance or ion flows through the TolQ-R TM channel (15).

The TolQ-TolR complex is homologous to the ExbB-ExbD complex that functions in membrane transport and to the MotA-MotB complex that forms the stator of the flagellar motor (12, 17, 18). Biochemical analyses on the homologous TolR and ExbD proteins using chemical cross-linkers indicated that TolR forms homo-dimers (8), whereas ExbD forms dimers and trimers (19). Determinants for TolR dimer formation mainly localize in the periplasmic domain of the protein. The three-dimensional structure of the *H. influenzae* TolR periplasmic domain confirmed the formation of homodimers with a head-to-tail conformation (16), whereas the three-dimensional structure of ExbD obtained under acid pH conditions corresponds to a monomer (20). No structural information is avail-

* This work was supported by the Centre National de la Recherche Scientifique and a grant from Agence National de la Recherche Projet Blanc SODATOL. This work was also supported by funds from SODATOL (to X. Y.-Z. Z.) and by a Fondation pour la Recherche Médicale fellowship (to E. L. G.). The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ To whom correspondence should be addressed: Laboratoire d'Ingénierie des Systèmes Macromoléculaires, Institut de Biologie Structurale et Microbiologie, Centre National de la Recherche Scientifique, UPR 9027, 31 chemin Joseph Aiguier, 13402 Marseille Cedex 20, France. Tel.: 33-491-16-46-63; Fax: 33-491-71-21-24; E-mail: lloubes@ibsm.cnrs-mrs.fr.

² The abbreviations used are: TM, transmembrane; TMH, transmembrane helix; PMF, proton motive force; WT, wild type; DTT, dithiothreitol; HA, hemagglutinin; Tricine, N-[2-hydroxy-1,1-bis(hydroxymethyl)ethyl]glycine.

TolR Transmembrane Helix Forms Homodimers

able for MotB; however, extensive mutagenesis has been performed using targeted disulfide cross-linking and evidenced that the TMHs from two MotB molecules are close enough to interact (21).

Using the whole batch of available data, an arrangement of four molecules of MotA and two MotB has been proposed in the flagellar motor (22). Based on the stoichiometry of one TolA molecule, a model for the arrangement of the TolQRA complex containing four to six TolQ and two TolR molecules has been proposed (12). Similar ratios were measured for the ExbB-D-TonB complex (23). This will form a minimal complex composed of 15–21 helices. To understand how the TolQRA TM segments interact in the membrane and how they mechanically respond to PMF or ion transit, we initiated a cysteine scanning approach. Herein, we performed a cysteine substitution mutagenesis of TolR TMH residues. The phenotypic consequences of residue replacements and the patterns of disulfide cross-linking obtained are consistent with an organization of two TolR TM segments in close contact through two distinct faces, suggesting helix rotation. Further experiments support the model in which ion transit at the TolQR TM helices interface induce a rotary motion of the TolR TM segment between the two faces.

EXPERIMENTAL PROCEDURES

Strain, Plasmid Construction, and Growth Conditions—GM1 *Escherichia coli* K12 strain (WT), and its derivatives, TPS300 and TPS13 (*tolR::Cm* and *tolQR*, respectively) (24), were used in this study. pOK-QR was constructed by insertion of a EcoRI-BamHI fragment from pTPS304 (a ColE1 pUC19 derivatives encoding *tolQ-tolR*) (24) into the P15A derivative pOK12 vector (25). pUC-R and pOK-R constructs were obtained by deletion of the *tolQ* gene from plasmid pTPS304 and pOK-QR, respectively, using recombinant PCR (26) and oligonucleotides 5'-TTAAACTCCGCGACAATAGACTTGGGAAGCGCAGAGGCTT and 5'-GTCTATTGTTCGCGGAGTTTAAGCAATGGCCAGAGCGCGTGGACGA. In these constructs, the *tolQ* Shine Dalgarno is followed by *tolR* gene. *tolR* mutants were engineered as described previously (12), using the pUC-R or the pOK-R plasmids as templates. Site-directed substitutions were introduced by QuikChange mutagenesis PCR using complementary pairs of oligonucleotides (sequences available upon request). Plasmid pOK-Q_{HA}, encoding a functional TolQ protein carrying a C-terminal hemagglutinin (HA) tag and compatible with the pUC-R derivatives, will be described elsewhere. pOK-Q_{HA} carrying point mutations (T145A, T178A, and P187V TolQ mutant proteins) were constructed by QuikChange mutagenesis. All of the constructs were verified by restriction analyses and further DNA sequencing. Routinely, cells were grown aerobically in LB medium at 37 °C supplemented with ampicillin (100 μg/ml; pUC derivatives), kanamycin (50 μg/ml; pOK derivatives), or chloramphenicol (40 μg/ml).

Assays for Outer Membrane Stability—Outer membrane permeability was assessed by measuring the level of detergent susceptibility, as well as the level of periplasmic RNase I release in the extracellular medium. Detergent susceptibility was measured by the level of survival of the strain on 1% deoxy-




FIGURE 1. Conservation of TolR anchor. *A*, alignment of TolR, ExbD, and MotB N-terminal consensus sequences (from Ref. 12) (*uppercase letter*, residue conserved in >90% of the sequences; *lowercase letter*, residue conserved in >60% of the sequences; *dot*, not conserved residue). The *E. coli* TolR N-terminal sequence is indicated (*bottom row*). The frame represents the position of the TM domain. The residues mutated in this study are *underlined* in the *E. coli* TolR sequence. *B*, steady state levels of WT and mutants TolR proteins used in this study. 0.2×10^8 cells were boiled in Laemmli buffer supplemented with β -mercaptoethanol, loaded onto 12.5% acrylamide Tricine SDS-PAGE, and immunodetected with anti-TolR polyclonal antibody. The molecular weight markers are indicated on the left.

cholate supplemented LB plates or on LB liquid medium containing various concentrations of SDS (as described previously) (27). In the latter case, survival is reported as the concentration of SDS sufficient to inhibit 50% of the cell growth after 3 h of cultures (LD_{50}). RNase I leakage was measured on RNA-containing plates, as described previously (27).

Colicin Tolerance Assay—Colicin activities were checked by the presence of halos on a lawn of the strain to be tested, as described previously (28). The data are reported as the maximal dilution of the colicin stock sufficient for inhibiting cell growth.

In Vivo Disulfide Bond Formation and Immunodetection—*In vivo* disulfide bond formation was essentially carried out as described previously (15) with slight modifications. 8×10^8 exponentially growing cells were harvested and resuspended in 1 ml of 20 mM sodium phosphate buffer, pH 6.8, and then treated for 10 min with 2.5 mM *N*-ethylmaleimide (Sigma) to block reduced thiol groups. When required, the cells were treated with the oxidative catalyst copper (II) orthophenanthroline 0.3 mM (CuOP; Sigma) prior washing in sodium phosphate buffer and blocking with *N*-ethylmaleimide. After centrifugation, the cell pellets were resuspended in Laemmli loading buffer in the presence or absence of the reducing agent β -mercaptoethanol. *In vivo* formaldehyde cross-linking was performed as described previously (29). Proteins were separated on a 15% SDS-PAGE, and immunodetections were performed using the anti-TolR polyclonal antibody (8), secondary goat antibodies coupled to alkaline phosphatase and 5-bromo-4-chloro-3-indolylphosphate (Sigma), and nitroblue tetrazolium (Sigma).

RESULTS

Sequence Conservation—The TolR, ExbD, and MotB proteins share similarities at the sequence level and probably at the structural and functional levels. Fig. 1*A* shows a sequence conservation consensus of all TolR, ExbD, and MotB TM anchors available on the protein data base. The general consensus obtained from the TolR, ExbD, and MotB consensus shows

that the aspartate residue at position 23, two leucine residues at positions 28 and 29, and a phenylalanine residue at position 32 are conserved, whereas many other positions present residue with similar hydrophobic properties (positions relative to the *E. coli* TolR sequence; Fig. 1A). However, because of the nature of a transmembrane helix and the general hydrophobicity of the residues, these residues probably act as structural determinants. The functional role of the Asp²³ residue has been documented in the TolR protein, as well as in the ExbD and MotB homologues. It has been proposed that this aspartate plays a central role in the PMF-dependent process and may couple ion or proton transport to mechanical movement within the TolQR-ExbBD-MotAB or PomAB complexes (12, 14, 30–33). A recent accessibility study showed that it is indeed the case because TolR C-terminal residues display different solvent accessibilities depending on PMF and residues important for channel formation, including the TolR Asp²³ residue (15). To understand how transmembrane helices organize together in the membrane and how PMF and hydrophilic ion channel residues regulate structural transitions, we initiated a cysteine scanning approach using site-directed mutagenesis.

Cysteine Scanning: Mutagenesis and Expression Levels of Cys Substitutions—To gain information about the structural organization and dynamic features of the TolR transmembrane anchor, each amino acid residue of the helix (residues 18–37) (34, 35) was individually replaced with cysteine. Mutations were introduced into the pUC-R plasmid, which expresses *tolR* at WT levels, by a recombinant PCR technique. Steady state levels of cysteine substituted TolR proteins were determined after introduction of the mutated plasmids into the *tolR* TPS300 strain. Initial studies using the polyclonal anti-TolR antibodies showed that the substituted TolR proteins migrate at the same position and accumulate at comparable levels as native TolR (Fig. 1B). However, as shown previously (5, 12), mutations of the Asp²³ or the Pro¹⁸⁷ residue led to migration defects on denaturing gel electrophoresis.

Phenotypic Characterization of Single Cysteine Mutants—The 20 singly substituted TolR mutants were tested for outer membrane stability, reflected by their capacity to grow on deoxycholate-containing plates as well as the level of periplasmic components release. Like most transmembrane helices, the TolR transmembrane segment was shown to be remarkably tolerant to cysteine substitution because all but three mutants displayed WT phenotype (Table 1). As expected, the strain producing the D23C TolR protein released RNase I in the medium and did not grow in presence of deoxycholate. Similar phenotypes were observed when the L22C and V24C *tolR* alleles were expressed. The TolR protein is also parasitized by bacterial toxins, called colicins, to penetrate and kill the target cell (for a review, see Ref. 36). We then tested the collection of substituted TolR mutants for group A colicin uptake, a process independent of PMF (36). All the strain producing the TolR mutants, except D23C, were killed by the Tol-dependent colicins (Table 1). We further tested the *tolR* substitutions for the uptake of group B colicins after introduction of the mutated plasmids into the *exbBD-tolQR* strain expressing *tolQ* (from the pOK-Q_{HA} plasmid). In contrast to group A colicins, group B colicins, except colicins 5 and 10, require energy from the ExbBD-TonB

TABLE 1
Phenotypes of strains producing Cys-substituted TolR proteins

TPS300 (*tolR*) cells expressing WT *tolR* or *tolR* cysteine mutants were tested for level of import of group A colicins A, N, and E2 (*tolR*-dependent colicins) and group B colicin D (*tolR*-independent), susceptibility to deoxycholate (DOC), and leakage of the periplasmic RNase I. Colicin sensitivity was tested on bacterial lawns by the spot dilution assay. The number indicated represents the maximal 10-fold dilution of the colicin stock (1 mg/ml) still able to kill the strain tested (R, resistant; S, sensitive; 4, sensitive to 10⁴ dilution). Deoxycholate resistance was tested by 100-fold dilutions of the strain on 1% DOC LB plates. The number indicated represents the maximal dilution for which the strain still grows. RNase I leakage was estimated on RNA containing LB plates (–, no leakage; +, low levels of leakage; ++, high levels of leakage). Identical results were obtained using *tolR* cysteine mutants expressed from the pOK-R plasmid (data not shown).

	Colicin				DOC	RNase
	A	N	E2	D		
None	R	R	R	S	S	++
WT	4	4	4	S	4	–
I18C	3	3	3	S	4	–
V19C	3	3	3	S	4	–
P20C	3	3	4	S	4	–
L21C	3	3	4	S	4	–
L22C	3	3	3	S	S	++
D23C	R	R	R	S	S	++
V24C	3	3	3	S	S	++
L25C	4	4	4	S	4	–
L26C	4	4	4	S	4	–
V27C	3	3	4	S	4	–
L28C	4	3	4	S	2	–
L29C	3	3	4	S	4	–
L30C	3	3	4	S	2	+
I31C	3	3	4	S	4	–
F32C	3	3	4	S	4	–
M33C	4	3	4	S	4	–
A34C	3	3	4	S	3	–
T35C	3	3	4	S	3	–
A36C	3	3	4	S	4	–
P37C	3	3	4	S	3	–

or the chimera TolQR-TonB complex (a phenomenon known as “cross-complementation”), to penetrate the cell (36).³ Production of TolQ and the TolR_{D23C} protein in the *exbBD-tolQR* strain renders cells resistant to all group B colicins tested (B, D, Ia, M, 5, and 10), whereas the L22C and V24C mutations prevented cell killing by group B colicins except colicins 5 and 10 (data not shown). These results confirmed that the D23C mutation abolishes all TolR functions (outer membrane stability and colicin import), whereas the L22C and V24C mutations specifically blocked the PMF-dependent TolR function (outer membrane stability) but not the energy-independent colicin import process. Taken together, the phenotype analysis suggests that the TolR Cys substitutions lie within the three previously characterized classes (Ref. 14 and Fig. 2): functional mutants (class I; all except L22C, D23C, and V24C), nonfunctional mutants (class II: D23C), and discriminative mutants (class III: L22C and V24C). This latter category includes mutations affecting TolQR channel activities, either at the level of channel formation (14) or prevention of conformational changes through disulfide bond formation between cysteine residues (15). Because of the nature of these substitutions, we suspected the L22C and V24C TolR proteins to dimerize. However, as seen on the helical wheel projection (Fig. 2), the Leu²² and Val²⁴ residues are located on opposite faces, suggesting that if the TolR TMH dimerizes, helix rotation should occur. Alternatively, these

³ E. L. Goemaere, L. Journet, I. Schalk, R. Lloubes, and E. Cascales, unpublished results.




FIGURE 3. TolR residues involved in homodimer formation. A, 0.2×10^8 cells of the indicated strain were boiled in Laemmli buffer in presence of β -mercaptoethanol (+ β ME, upper panel) or in absence of reducing agent ($-\beta$ ME, lower panel), loaded onto 12.5% acrylamide SDS-PAGE, and immunodetected with anti-TolR polyclonal antibody. The positions of TolR and TolR dimers are indicated on the right. *, low levels of dimerization; **, high levels of dimerization. The molecular weight markers are indicated on the left. B, representation of results from A on a helical wheel projection. The interface of dimerization is indicated by the dotted line. Arrows indicate positions of substitutions that form dimers (solid arrows, high levels; dotted arrows, low levels).

and that rotation between the two helices is induced by torque generation by the TolQR ion channel.

DISCUSSION

The TolQ, TolR, and TolA proteins form a complex at the inner membrane, interacting through their transmembrane segments, as evidenced using chemical cross-linking or isolation of allele-specific suppressive mutations (5–8, 12, 14). These led to the conclusion that the transmembrane helix of TolR is at the proximity of the helices II and III of TolQ, whereas TolQ transmembrane III contacts the transmembrane I, which is in interaction with the transmembrane anchor of the TolA protein. Stoichiometry analyses showed that the ratio of the TolQ-TolR-TolA complex is 4–6:2:1 (4, 12), adding further complexity in helix organization. However, trying to reconcile all the results and the faces of transmembrane segments in interaction has been shown to be difficult, raising the possibility that structural modifications and movements of helices during functioning may occur. The idea that the TolQRA complex undergoes conformational changes was first suggested by the observation that TolA interacts with the outer membrane-anchored peptidoglycan-associated lipoprotein Pal in a PMF- and TolQR-dependent manner (10, 12). Structural modifications of TolA dependent on PMF and on the TolQ and TolR proteins were demonstrated using protease accessibilities (13). Recently, we demonstrated using cysteine accessibilities that the C-terminal periplasmic domain of the TolR protein is subjected to structural modifications (15).

The observation that the TolQ and TolR helices share similarities, not only with the homologous ExbB and ExbD proteins

but also with the flagella stator components MotA and MotB, led to the idea that these three couples of proteins may share a common mechanism of energy transfer (12, 17, 18). This was exemplified by the presence of a conserved aspartate residue with TolR, ExbD, and MotB, which is critical for function (12, 30–32, 38). Following these studies, it has been proposed that the three helices (TolR, TolQ-II, and TolQ-III) may delimit a channel by which ions transit and that a “signal” is then transmitted to the TolA protein via the TolQ-I segment. An analysis of TolQ transmembrane residues further demonstrated that short length side chain residues are involved in helix packing, whereas hydrophilic residues may participate in the energy transduction mechanism (14). In the current model, which is actively tested in our laboratory, a dimer of TolR anchors is surrounded by four to six TolQ molecules, thus allowing the formation of four to six ion path-

ways, for which channel activity may be regulated by mechanical movements.

To determine how the TolQRA helices interact and how they undergo structural modifications, we performed a systematic cysteine scanning of the TolQRA helices. Here, we reported results obtained for the TolR TM helix. Our results clearly showed that cysteines from two distinct monomers are close enough to efficiently form a disulfide bond dimer without the aid of an oxidative agent. Similar results were obtained in a systematic cysteine scanning study of the flagellar MotB protein TMH (21). In the case of the TolR protein, this is clearly shown for the L22C and V24C mutants. Disulfide bond formation blocks TolR dimers, which prevents function of the Tol system and growth in presence of detergent. Interestingly, this phenotype is reversed by the addition of reducing agent in the medium, as shown for cysteine mutations occluding the SecY protein channel (39). On a perfect helical wheel projection of the TolR TMH, the Leu²² and Val²⁴ residues are separated by an angle of $\sim 160^\circ$. In between, most substitutions form dimers, albeit at lower levels. These results raise the hypotheses (i) that the TolR transmembrane helix rotates between the two “extreme” positions (Leu²² and Val²⁴) using the “intermediate” positions such as molecular ratchet gears or cogwheels or (ii) that the transmembrane helix is subject to physical torsions or constraints in such a way that all these residues (intermediate and extreme) are located on a reduced, unique face. The first model has been proposed in light of similar results using the MotB transmembrane anchor as model system (21). In the case of MotB, projection on a similar wheel than the one used in Fig. 2 shows that the angle between the two extreme residues is

TolR Transmembrane Helix Forms Homodimers




FIGURE 4. TolR functioning is specifically prevented by disulfide bond formation. 10^9 bacteria of the indicated strain were inoculated in 10 ml of medium in presence of 0.25% SDS, and growth was monitored by measuring absorbance at 600 nm every 45 min. After 90 min of culture, DTT was added (closed circles) or not (closed squares) to the final concentration of 5 mM (arrow).

~110°. However, these results cannot discriminate between the two hypotheses. One may hypothesize that in the absence of movement, the TolR dimer rotation will be stopped in both of the extreme positions (hypothesis (i) above), whereas all extreme and intermediate substitutions will still promote helix dimerization (hypothesis (ii) above). Our results using mutations that affect residues that regulate ion transit favor the first hypothesis. It seems thus likely that the two TolR helices from a dimer oscillate between two extreme conformations (Fig. 6). Alternatively, because the 4–6:2 stoichiometry of the TolQR




FIGURE 5. TolR anchor rotations depend on key residues of the ion pathway. 0.2×10^8 cells producing the indicated TolR Cys substitution (lanes) in combination with the TolR or TolQ mutations (panels), treated with the oxidative agent copper (II) orthophenanthroline to increase dimer formation, were boiled in Laemmli buffer in absence of reducing agent, loaded onto 12.5% acrylamide SDS-PAGE, and immunodetected with anti-TolR polyclonal antibody. The positions of TolR and TolR dimers are indicated on the right. The molecular weight markers are indicated on the left. The mutant names on the top and on the left indicate positions of primary and secondary mutations, respectively.

complex represents the “minimal complex,” one may hypothesize that TolR form tetramers. In this case, both L22C and V24C substitutions may form disulfide bonds independently of helix rotation. However, this hypothesis can be ruled out because the L22C/V24C TolR double mutant only forms dimers as single substitutions do (data not shown).

The Asp²³ residue, which has been proposed to be critical for ion conduction, is located ~100° from either Leu²² or Val²⁴ on a helical wheel projection (Fig. 2). Using the rotary motion, the Asp²³ residue may thus alternate between two different ion channels, as suggested for MotB. Such a rotation may thus place the Thr³⁵ residue at the position previously occupied by the Asp²³ residue in the channel. The presence of a hydrophilic residue at that position may thus stabilize the putative aqueous channel. The TolR T35C mutant displayed a WT phenotype, but mutation of the Thr³⁵ residue by an alanine confers cell sensitivity toward deoxycholate, but cells remain sensitive to




FIGURE 6. A model for the TolR anchor rotary motion. The TolR anchors from two distinct TolR molecules are represented at the extreme positions of the dimerization interface (corresponding to high levels of dimerization) and thus oscillate between conformations 1 and 2 (red arcs). This may involve passage through a number of intermediate conformations (corresponding to low levels of dimerization) such as molecular gears. The aspartate residue then oscillates between two channels, regulating opening (green areas, open or active channels) or closure (blue areas, closed or inactive channels).

colicins (data not shown), classifying such a mutation as discriminative (class III), suggesting a functional role of this residue in response to PMF. This implies that rotation of the TolR anchor is functionally important, which is confirmed by the fact that disulfide bond formation with TolR L22C or V24C lead to cells that are impaired in membrane integrity but still sensitive for colicin import (Tables 1 and 2).

Because the level of dimerization of the TolR L22C and V24C is higher than for the intermediate position, one may hypothesize that the time necessary to reach one extreme position from the other is fast and that the rotary motion stops transiently at the extreme positions. However, the TolR F32C and T35C mutant do not dimerize, even if these residues are located between the two extreme positions. This may suggest that the two TolR helices from a dimer are not close together on the whole length but rather are closer at the cytoplasmic side than at the periplasmic side of the inner membrane. This may account for the strong dimerization of Leu²² and Val²⁴ residues, which are situated at the cytoplasmic side of the membrane; however, the Ile¹⁸ and Leu²¹ residues, which are located at the cytoplasmic side of the TolR helix, have intermediate dimerization properties. Our results showed that, similarly to TolR C-terminal domain movement (15), TolR helix rotation is regulated by three channel residues, the TolR Asp²³ residue and the Thr¹⁴⁵ and Thr¹⁷⁸ residues located on helices II and III of TolQ, respectively. In the case of the MotAB complex, it has been shown using protease accessibilities that MotA conformation is regulated by the MotB Asp residue (17). The discriminative mutation P187V does not abolish rotation, a result consistent with the observation that this residue does not regulate TolR C-terminal conformational modification (15). The TolQ Pro¹⁸⁷ residue has been shown to be required for the Tol complex functioning and has been suggested to couple ion transit to conformational changes (14). Our results suggest that TolR

anchors rotate independently of Pro¹⁸⁷, which thus may regulate conformational modifications of TolQ.

Overall, our results showed that two helices from two different TolR molecules are close to each other in the TolQR complex. These two TolR TMHs rotate as molecular gears between two positions, L22C and V24C. This rotation is functionally important for activity of the TolQR ion channel.

Acknowledgments—We thank Denis Duché, Philippe Roche, Christophe Meynier, and members of the Llobès, Sturgis, and Bouveret teams for helpful discussions Denis Duché and Laure Journet for critical reading of the manuscript; and Cécile Ancieux for encouragement.

REFERENCES

- Webster, R.E. (1991) *Mol. Microbiol.* **5**, 1005–1011
- Llobès, R., Cascales, E., Walburger, A., Bouveret, E., Lazdunski, C., Bernadac, A., and Journet, L. (2001) *Res. Microbiol.* **152**, 523–529
- Gerding, M. A., Ogata, Y., Pecora, N. D., Niki, H., and de Boer, P. A. (2007) *Mol. Microbiol.* **63**, 1008–1025
- Guihard, G., Boulanger, P., Benedetti, H., Llobès, R., Besnard, M., and Letellier, L. (1994) *J. Biol. Chem.* **269**, 5874–5880
- Lazzaroni, J. C., Vianney, A., Popot, J. L., Benedetti, H., Samatey, F., Lazdunski, C., Portalier, R., and Geli, V. (1995) *J. Mol. Biol.* **246**, 1–7
- Déroutiche, R., Bénédetti, H., Lazzaroni, J. C., Lazdunski, C., and Llobès, R. (1995) *J. Biol. Chem.* **270**, 11078–11084
- Germon, P., Clavel, T., Vianney, A., Portalier, R., and Lazzaroni, J. C. (1998) *J. Bacteriol.* **180**, 6433–6439
- Journet, L., Rigal, A., Lazdunski, C., and Bénédetti, H. (1999) *J. Bacteriol.* **181**, 4476–4484
- Bouveret, E., Déroutiche, R., Rigal, A., Llobès, R., Lazdunski, C., and Bénédetti, H. (1995) *J. Biol. Chem.* **270**, 11071–11077
- Cascales, E., Gavioli, M., Sturgis, J. N., and Llobès, R. (2000) *Mol. Microbiol.* **38**, 904–915
- Walburger, A., Lazdunski, C., and Corda, Y. (2002) *Mol. Microbiol.* **44**, 695–708
- Cascales, E., Llobès, R., and Sturgis, J. N. (2001) *Mol. Microbiol.* **42**, 795–807
- Germon, P., Ray, M. C., Vianney, A., and Lazzaroni, J. C. (2001) *J. Bacteriol.* **183**, 4110–4114
- Goemaere, E. L., Cascales, E., and Llobès, R. (2007) *J. Mol. Biol.* **366**, 1424–1436
- Goemaere, E. L., Devert, A., Llobès, R., and Cascales, E. (2007) *J. Biol. Chem.* **282**, 17749–17757
- Parsons, L. M., Grishaev, A., and Bax, A. (2008) *Biochemistry* **47**, 3131–3142
- Kojima, S., and Blair, D. F. (2001) *Biochemistry* **40**, 13041–13050
- Zhai, Y. F., Heijne, W., and Saier, M. H., Jr. (2003) *Biochim. Biophys. Acta* **1614**, 201–210
- Higgs, P. I., Myers, P. S., and Postle, K. (1998) *J. Bacteriol.* **180**, 6031–6038
- García-Herrero, A., Peacock, R. S., Howard, S. P., and Vogel, H. J. (2007) *Mol. Microbiol.* **66**, 872–889
- Braun, T. F., and Blair, D. F. (2001) *Biochemistry* **40**, 13051–13059
- Braun, T. F., Al-Mawsawi, L. Q., Kojima, S., and Blair, D. F. (2004) *Biochemistry* **43**, 35–45
- Higgs, P. I., Larsen, R. A., and Postle, K. (2002) *Mol. Microbiol.* **44**, 271–281
- Sun, T. P., and Webster, R. E. (1987) *J. Bacteriol.* **169**, 2667–2674
- Vieira, J., and Messing, J. (1991) *Gene (Amst.)* **100**, 189–194
- Ansaldi, M., Lepelletier, M., and Méjean, V. (1996) *Anal. Biochem.* **234**, 110–111
- Cascales, E., and Llobès, R. (2004) *Mol. Microbiol.* **51**, 873–885
- Journet, L., Bouveret, E., Rigal, A., Llobès, R., Lazdunski, C., and Bénédetti, H. (2001) *Mol. Microbiol.* **42**, 331–344
- Cascales, E., Bernadac, A., Gavioli, M., Lazzaroni, J. C., and Llobès, R.

ToIR Transmembrane Helix Forms Homodimers

- (2002) *J. Bacteriol.* **184**, 754–759
30. Braun, V., Gaisser, S., Herrmann, C., Kampfenkel, K., Killmann, H., and Traub, I. (1996) *J. Bacteriol.* **178**, 2836–2845
31. Zhou, J., Sharp, L. L., Tang, H. L., Lloyd, S. A., Billings, S., Braun, T. F., and Blair, D. F. (1998) *J. Bacteriol.* **180**, 2729–2735
32. Kojima, S., Shoji, T., Asai, Y., Kawagishi, I., and Homma, M. (2000) *J. Bacteriol.* **182**, 3314–3318
33. Kojima, S., and Blair, D. F. (2004) *Int. Rev. Cytol.* **233**, 93–134
34. Kampfenkel, K., and Braun, V. (1993) *J. Bacteriol.* **175**, 4485–4491
35. Muller, M. M., Vianney, A., Lazzaroni, J. C., Webster, R. E., and Portalier, R. (1993) *J. Bacteriol.* **175**, 6059–6061
36. Cascales, E., Buchanan, S. K., Duché, D., Kleanthous, C., Lloubès, R., Postle, K., Riley, M., Slatin, S., and Cavard, D. (2007) *Microbiol. Mol. Biol. Rev.* **71**, 158–229
37. Duché, D., Baty, D., Chartier, M., and Letellier, L. (1994) *J. Biol. Chem.* **269**, 24820–24825
38. Blair, D. F., Kim, D. Y., and Berg, H. C. (1991) *J. Bacteriol.* **173**, 4049–4055
39. Saparov, S. M., Erlandson, K., Cannon, K., Schaletzky, J., Schulman, S., Rapoport, T. A., and Pohl, P. (2007) *Mol. Cell.* **26**, 501–509