

## Pituitary Neoplasm Nomenclature Workshop: Does Adenoma Stand the Test of Time?

Ken Ho, Maria Fleseriu, Ursula Kaiser, Roberto Salvatori, Thierry Brue, M Beatriz Lopes, Pamela Kunz, Mark Molitch, Sally Camper, Mônica Gadelha, et al.

► **To cite this version:**

Ken Ho, Maria Fleseriu, Ursula Kaiser, Roberto Salvatori, Thierry Brue, et al.. Pituitary Neoplasm Nomenclature Workshop: Does Adenoma Stand the Test of Time?. Journal of the Endocrine Society, Oxford University Press, 2021, 5 (3), 10.1210/jendso/bvaa205 . hal-03310331

**HAL Id: hal-03310331**

**<https://hal-amu.archives-ouvertes.fr/hal-03310331>**

Submitted on 30 Jul 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Pituitary Neoplasm Nomenclature Workshop: Does Adenoma Stand the Test of Time?

Ken Ho<sup>1</sup>, Maria Fleseriu<sup>2</sup>, Ursula Kaiser<sup>3</sup>, Roberto Salvatori<sup>4</sup>, Thierry Brue<sup>5</sup>, M. Beatriz Lopes<sup>6</sup>, Pamela Kunz<sup>7</sup>, Mark Molitch<sup>8</sup>, Sally A. Camper<sup>9</sup>, Mônica Gadelha<sup>10</sup>, Luis V. Syro<sup>11</sup>, Edward Laws<sup>3</sup>, Martin Reincke<sup>12</sup>, Hiroshi Nishioka<sup>13</sup>, Ashley Grossman<sup>14</sup>, Ariel Barkan<sup>9</sup>, Felipe Casanueva<sup>15</sup>, John Wass<sup>16</sup>, Adam Mamelak<sup>17</sup>, Laurence Katznelson<sup>18</sup>, A.J. van der Lely<sup>19</sup>, Sally Radovick<sup>20</sup>, Martin Bidlingmaier<sup>12</sup>, Margaret Boguszewski<sup>21</sup>, Jens Bollerslev<sup>22</sup>, Andrew R. Hoffman<sup>18</sup>, Nelson Oyesiku<sup>23</sup>, Gerald Raverot<sup>24</sup>, Anat Ben-Shlomo<sup>17</sup>, Rob Fowkes<sup>25</sup>, Ilan Shimon<sup>26</sup>, Hidenori Fukuoka<sup>27</sup>, Alberto M. Pereira<sup>28</sup>, Yona Greenman<sup>29</sup>, Anthony P. Heaney<sup>30</sup>, Mark Gurnell<sup>31</sup>, Gudmundur Johannsson<sup>32</sup>, Robert Y. Osamura<sup>33</sup>, Michael Buchfelder<sup>34</sup>, Maria Chiara Zatelli<sup>35</sup>, Marta Korbonits<sup>36</sup>, Philippe Chanson<sup>37</sup>, Nienke Biermasz<sup>28</sup>, David R. Clemmons<sup>38</sup>, Niki Karavitaki<sup>39</sup>, Marcello D. Bronstein<sup>40</sup>, Peter Trainer<sup>41</sup>, Shlomo Melmed<sup>17</sup>

<sup>1</sup>The Garvan Institute of Medical Research, Sydney, Australia; <sup>2</sup>Oregon Health & Science University, Portland, Oregon, USA; <sup>3</sup>Brigham and Women's Hospital, Boston, Massachusetts, USA; <sup>4</sup>The Johns Hopkins University School of Medicine, Baltimore, Maryland, USA; <sup>5</sup>Aix-Marseille University, Marseille, France; <sup>6</sup>University of Virginia School of Medicine, Charlottesville, Virginia, USA; <sup>7</sup>Yale School of Medicine, New Haven, Connecticut, USA; <sup>8</sup>Northwestern University Feinberg School of Medicine, Chicago, Illinois, USA; <sup>9</sup>University of Michigan Medical School, Ann Arbor, Michigan, USA; <sup>10</sup>Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil; <sup>11</sup>Hospital Pablo Tobon Uribe and Clínica Medellín – Grupo Quirónsalud, Medellín, Colombia; <sup>12</sup>Klinikum der Universität, Ludwig-Maximilians-Universität, München, Germany; <sup>13</sup>Toranomon Hospital, Tokyo, Japan; <sup>14</sup>University of Oxford, Oxford, and Barts and the London School of Medicine, London, UK;

© The Author(s) 2021. Published by Oxford University Press on behalf of the Endocrine Society.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial reproduction and distribution of the work, in any medium, provided the original work is not altered or transformed in any way, and that the work is properly cited. For commercial re-use, please contact [journals.permissions@oup.com](mailto:journals.permissions@oup.com)

<sup>15</sup>Santiago de Compostela University, Santiago de Compostela, Spain; <sup>16</sup>Churchill Hospital, Oxford, UK; <sup>17</sup>Cedars-Sinai Medical Center, Los Angeles, California, USA; <sup>18</sup>Stanford University School of Medicine, Stanford, California, USA; <sup>19</sup>Erasmus University Medical Center, Rotterdam, The Netherlands; <sup>20</sup>Rutgers-Robert Wood Johnson Medical School, New Brunswick, New Jersey, USA; <sup>21</sup>Federal University of Parana, Curitiba, Brazil; <sup>22</sup>University of Oslo, Oslo, Norway; <sup>23</sup>Emory University School of Medicine, Atlanta, Georgia, USA; <sup>24</sup>Hospices Civils de Lyon and Lyon 1 University, Lyon, France; <sup>25</sup>Royal Veterinary College, University of London, London, UK; <sup>26</sup>Rabin Medical Center, Beilinson Hospital, Petah-Tikva, Israel; <sup>27</sup>Kobe University Hospital, Kobe, Japan; <sup>28</sup>Leiden University Medical Center, Leiden, The Netherlands; <sup>29</sup>Tel Aviv-Sourasky Medical Center, Tel Aviv University, Tel Aviv, Israel; <sup>30</sup>David Geffen School of Medicine University of California, Los Angeles, California, USA; <sup>31</sup>University of Cambridge & Addenbrooke's Hospital, Cambridge, UK; <sup>32</sup>Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden; <sup>33</sup>Nippon Koukan Hospital Kawasaki & Keio University School of Medicine, Tokyo, Japan; <sup>34</sup>University Hospital Erlangen, Erlangen, Germany; <sup>35</sup>University of Ferrara, Ferrara, Italy; <sup>36</sup>Queen Mary University of London, UK; <sup>37</sup>University Paris-Saclay & Assistance Publique Hôpitaux de Paris, Hôpital Bicêtre, Le Kremlin-Bicêtre, France; <sup>38</sup>University of North Carolina School of Medicine, Chapel Hill, North Carolina, USA; <sup>39</sup>University of Birmingham, Birmingham, UK; <sup>40</sup>University of São Paulo Medical School, São Paulo, Brazil; <sup>41</sup>University of Manchester, Manchester, UK.

**Funding:** Supported by unrestricted educational grants to the Pituitary Society from Chiasma, Inc., Corcept Therapeutics, Ipsen Biopharmaceuticals, Novo Nordisk, Pfizer Inc., Strongbridge Biopharma, and Tiburio Therapeutics. Funding sources had no role in data collection, analysis, and interpretation, decision to publish, or preparation of the manuscript.

**Corresponding author:**

Ken Ho, MD

The Garvan Institute of Medical Research

384 Victoria Street

Sydney, NSW 2010

Australia

**Disclosure Summary:** The authors have declared that no conflict of interest exists.

Accepted Manuscript

## Abstract

The *WHO Classification of Endocrine Tumours* designates pituitary neoplasms as adenomas. A proposed nomenclature change to neuroendocrine tumors (PitNETs) has been met with concern by some stakeholder groups. The Pituitary Society coordinated the Pituitary Neoplasm Nomenclature (PANOMEN) Workshop to address the topic. Experts in pituitary developmental biology, pathology, neurosurgery, endocrinology, and oncology, including representatives nominated by the Endocrine Society, European Society of Endocrinology, European Neuroendocrine Association, Growth Hormone Research Society, and International Society of Pituitary Surgeons. Clinical epidemiology, disease phenotype, management, and prognosis of pituitary adenomas differ from that of most NETs. The vast majority of pituitary adenomas are benign and do not adversely impact life expectancy. A nomenclature change to PitNET does not address the main challenge of prognostic prediction, assigns an uncertain malignancy designation to benign pituitary adenomas, and may adversely affect patient. Due to pandemic restrictions, the workshop was conducted virtually, with audio-visual lectures and written précis on each topic provided to all participants. Feedback was collated and summarized by Content Chairs and discussed during a virtual writing meeting moderated by Session Chairs, which yielded an evidence-based draft document sent to all participants for review and approval. There is not yet a case for adopting the PitNET nomenclature. The PANOMEN Workshop recommends that the term adenoma be retained and that the topic be revisited as new evidence on pituitary neoplasm biology emerges.

**Keywords:** Pituitary neoplasm, pituitary adenoma, tumor, neuroendocrine

## Introduction

Pituitary neoplasms are classified according to size, location, secretory function, cell type, and neoplastic behavior. Histologically, classification is defined by the *WHO Classification*, a series that is updated regularly based on advances in research that impact diagnostic pathology. The current 2017 *WHO Classification* designates pituitary neoplasms as “adenomas,” recognizing that the vast majority of these lesions are benign and only a small subset exhibit aggressive, and exceptionally rarely, malignant characteristics (1).

In 2017, the International Pituitary Pathology Club proposed that pituitary adenomas be termed pituitary neuroendocrine tumors (PitNETs) because pituitary hormone-producing cells are neuroendocrine cells and invasive pituitary adenomas share some behavioral similarities with NETs (2). The Pituitary Society highlighted significant concerns for the proposed change (3), which generated further discussion (4,5).

To enable contributions from key stakeholders, the Pituitary Society convened an international multidisciplinary workshop, Pituitary Neoplasm Nomenclature (PANOMEN), designed to address whether categorizing pituitary adenomas as NETs is supported by scientific evidence, whether NET terminology is more appropriate and representative of pituitary neoplastic behavior compared with the term “adenoma,” and whether changing to the NET nomenclature benefits patient care.

## Workshop Planning and Structure

Guiding principles that determined PANOMEN workshop planning were inclusiveness, stakeholder participation, transparency, editorial independence, and unbiased selection of speakers and discussants. Invitations were extended by the Pituitary Society to relevant professional societies including the Endocrine Society, European Society of Endocrinology, European Neuroendocrine Association, Growth Hormone Research Society, and International Society of Pituitary Surgeons, as well as to expert pituitary opinion leaders to ensure balanced multidisciplinary representation across developmental biology, pathology, neurosurgery, endocrinology, and oncology. An invitation was

also extended to leaders of the International Agency for Research on Cancer (IARC), responsible for development of the WHO Classification series.

PANOMEN focused on clinical epidemiology, taxonomy, classification, and aggressive neoplasms; because of pandemic restrictions, it was conducted entirely virtually. Speakers delivered remote audio-visual lectures and disseminated written lecture précis for review by all participants. Content Chairs of each session, with complete editorial independence, integrated speaker précis and participants' comments into assigned topic summaries, which formed the backbone of a document developed for a virtual meeting moderated by respective Session Chairs.

From this virtual meeting, a Writing Group comprising Speakers, Content Chairs, Session Chairs, and Workshop Chairs developed a balanced and accurate evidence-based draft document, which was then circulated to all participants for further feedback. An edited draft was then recirculated for final review and acceptance. Workshop sponsors played no role in this process and did not review the draft before or after the virtual meeting.

## **Epidemiology and Clinical Outcomes**

### ***Pituitary Neoplasms***

Pituitary tumors account for about 15% of all intracranial masses. The overwhelming majority of these are benign adenomas arising from adenohypophyseal cells (6,7). Population surveys show that pituitary adenomas affect health outcomes due to consequences of hormone excess or deficiency, or less commonly, from a mass effect in about 1 in 1000 persons in the community (6). However, pituitary neoplasms are much more common than clinically apparent. This is because autopsy and imaging studies of persons without a known history of pituitary disease show that 10-15% of the population harbor undiagnosed pituitary neoplasms, most of which are small (8-10). Just over half of clinically significant pituitary adenomas undergo surgical resection; of these, about 15% are locally invasive (5,7). Thus, invasive behavior is uncommon even among neoplasms requiring surgery, and is

very rarely encountered clinically because surgery is not indicated for the overwhelming majority of pituitary neoplasms. Pituitary carcinomas are exceptionally rare, accounting for <0.5% of invasive lesions (11).

Subclinical adenomas, which account for more than 99% of pituitary neoplasms, do not affect life expectancy. Among the 1% of pituitary clinically significant neoplasms, the order of prevalence is PRL-secreting (50-60%), non-functioning (20-40%), GH-secreting (10-15%) and ACTH-secreting (5-8%) adenomas (6,7,12).

Prolactinomas are usually successfully treated with dopamine agonists, and surgery is required in <10% of patients (6,7). No standardized mortality ratio (SMR) has been reported for prolactinomas. Up to 25% of untreated non-secreting adenomas enlarge under observation (6); any growth is generally very slow, ranging from 0.4 mm to 1.0 mm/year. For symptomatic, surgically resected, non-secreting adenomas, about one-third recur or progress over time (6). The SMR for patients with invasive recurrent macroadenomas reported in studies from the United Kingdom, Denmark and Sweden range from 1.1 to 3.6, with excess deaths due mainly to circulatory, respiratory, and infectious causes (13). Acromegaly is usually treated first with surgery, and medical treatment is commonly required for those incompletely controlled. The SMR is 2.0 to 3.0 in untreated acromegaly but falls to that of the general population with effective hormonal control (14). Over 90% of patients with Cushing disease have microadenomas and up to 80% of these can be controlled by surgery (6). Disease control reduces hypercortisolemia-driven SMR from 3.7 to 1.2 (15,16).

Reduced quality of life and morbidity are influenced by hormonal hypersecretion or hyposecretion and treatment effects, and rarely by adenoma mass effects (17).



## ***Neuroendocrine Tumors***

NETs were first described in 1907 by Oberndorfer who coined the term “Karzinoide,” meaning cancer-like, to describe tumorlets found in the small intestine (18). NETs, widely acknowledged as potentially cancerous, are epithelial neoplasms derived from neuroendocrine cells and most commonly originate in the gastrointestinal tract, followed by the lungs and other sites. Although of epithelial origin, pituitary adenomas are not classified as NETs (19). Primary pituitary NETs (also referred to as carcinoids in the literature) are extremely rare (see section on Aggressive Neoplasms) and should be differentiated from metastatic NETs of other organs (20).

While NETs are rare, the age-adjusted incidence has risen over the past decades, and increased 6-fold to 6.98 per 100,000 in 2012 (21,22). A recent autopsy study reported a NET prevalence of about 0.5% (23). Median survival is 9.3 years, with the highest rates seen in localized disease (>30 years), Grade 1 NETs (16.2 years), and NETs of the appendix (30 years) (21).

Treatment decisions for NETs are based on functional status, extent of disease, grade and differentiation, rate of growth, primary site, and somatostatin receptor (SST) expression status. NETs are extremely heterogeneous, and confusing nomenclature hampers accurate diagnosis, treatment selection, and epidemiologic studies.

### **Taxonomy**

#### ***Human Pituitary Development and Cellular Differentiation***

The anterior pituitary lobe is derived from embryonic oral ectoderm while the posterior lobe is derived from neural ectoderm (24,25). At 5 weeks gestation, oral ectoderm invaginates to form Rathke’s pouch, which then separates, and the pituitary stalk and posterior lobe form an evagination of neural ectoderm. By 13 weeks, the overall pituitary structure is established.

Signaling pathways important for early pituitary growth and development include those that regulate development of other organs from cranial placodes, as well as pathways common to pituitary and gastrointestinal neuroendocrine cells (25). Transcription factors determine hormone-specific

pituitary stem cell development. Immunostaining (IHC) for pituitary hormones and cell-specific transcription factors enables classification of differentiated pituitary adenomas based on pituitary cell lineage ([Table 1](#)).

### ***Anterior Pituitary Cell Type Markers***

Most functioning adenomas can be simply and reliably classified by immunohistochemistry (IHC) of pituitary hormones. The 2017 WHO classification is based on histological markers with pituitary hormones, low-molecular-weight cytokeratin (LMWCK, or CAM5.2), and transcription factors ([Table 1](#)). In adenomas that are either immunonegative or only faintly positive for pituitary hormones, transcription factor staining determines lineage classification (26). Null cell adenomas are immunonegative for both pituitary hormones and specific transcription factors.

Transcription factor IHC also helps differentiate non-secreting pituitary adenomas from non-adenomatous pituitary neoplasms, including craniopharyngiomas, meningiomas, and paragangliomas, as well as metastatic NETs that also express neuroendocrine markers such as synaptophysin, chromogranin A, and SSTs. These latter markers are also positive in most pituitary tumors, although they are not used for diagnosis as they are not pituitary-specific. Synaptophysin and SSTs, but not chromogranin A, are also expressed in follicular thyroid (27) and adrenal cortical adenomas (28), neither of which are classified as NETs. The question of whether pituitary adenomas are biologically distinct from extra-pituitary neuroendocrine neoplasms (29) requires further rigorous comparative genomic and molecular single-cell analyses within and between tissue types.

## Classifications

### *Surgical Classifications*

Cushing, in his 1912 monograph “The Pituitary Body and Its Disorders” (30), coined the term *pituitary adenomas* after analysis of 47 patients with pituitary disease.

Subsequently, Hardy’s classification is based on the size and stage of pituitary adenomas, and describes “microadenomas” that are resectable with preservation of the normal pituitary gland (31). Further sub-classifications proposed by Kovacs and Horvath were based on adenoma electron microscopy ultrastructure (32). Cavernous sinus invasion by pituitary adenomas forms the basis of the systematic imaging classification by Knosp (33).

The phenomenon of invasive pituitary adenomas remains poorly understood (34). In a series of 354 surgically resected macroadenomas, histologic dural invasion was present in 45% of cases and increased in frequency with increasing adenoma size, but did not affect adenoma recurrence rates (35). As invasiveness does not necessarily imply aggressiveness, there is a need for better understanding as to why some adenomas invade but are so rarely malignant. Indeed, malignancy, which is exceptionally rare among pituitary adenomas, is a strikingly differentiating characteristic from the natural history of most NETs.

### **2017 WHO Classification of Pituitary Neoplasms**

The 2017 WHO classification is based on immunoexpression of pituitary hormones, pituitary-specific transcription factors, and other cell differentiating co-factors (1). Pituitary cell lineage-based classification has been validated by specific genomic, epigenetic, and methylation signatures (36).

The WHO solely classifies pituitary neoplasms as either *adenomas* or *carcinomas* and abandoned the 2004 category of *atypical adenoma* (19) due to a lack of clinical evidence that tumor behavior differed between typical and atypical adenomas (1). Rather, the 2017 classification

incorporated proliferation (mitotic count and Ki-67 labeling index) and tumor invasion in histological evaluation, both of which correlate with more aggressive tumor behavior. It also recognized histological variants of “high-risk” pituitary adenomas that behave more aggressively, represented mainly by sparsely granulated somatotroph adenomas, silent corticotroph adenomas, Crooke’s cell adenomas, the newly termed pluri-hormonal PIT-1 positive adenomas, and lactotroph adenomas in men. However, it acknowledged a lack of histopathological prognosticators for these clinically aggressive pituitary adenomas.

*What are the consequences of changing classification nomenclature?*

Any suggestion to change a disease name should address biological relevance as well as practicality, acceptability, and nomenclature principles. When considering a change of pituitary neoplasm nomenclature from adenoma to NET, it is imperative to rigorously consider whether the distinction between endocrine and neuroendocrine neoplasm is conceptual or histological, and whether NETs and adenomas are morphologically distinct. Importantly, unlike NETs, the overwhelming majority of pituitary neoplasms do not require biopsy for histological definition, including those that are treated medically.

Rationales supporting a nomenclature change include recognizing that some pituitary tumors may not behave in a benign clinical manner. Thus, it has been argued that clinically aggressive tumors should not be termed “adenomas,” a terminology implying a benign clinical behavior. However, a change to NET nomenclature does not provide additive guidance for distinguishing between tumors that remain benign and those that behave more aggressively. Furthermore, a proposed change to NET nomenclature also implies including relatively benign pituitary neoplasms in the very sizable category of neuroendocrine neoplasms arising in other organs/systems. This approach could potentially provide an integrated classification whereby benign pituitary adenomas are classified with the entire group of neuroendocrine neoplasms (carcinoid tumors).



## Aggressive Pituitary Neoplasms

Pituitary adenomas account for more than 95% of the many tumor types that arise in the sella (e.g., craniopharyngiomas, chordomas) (6,7). Of the adenomas, <0.1% of all clinically significant resected and nonresected lesions exhibit aggressive behavior (3). In this context, the term *invasive* is inappropriately used as if synonymous with *aggressive*. Invasiveness may best be considered as an imaging diagnosis, sometimes found on pathological examination, and describes tumors that infiltrate adjacent structures, such as the cavernous sinuses, bone, and sphenoid sinus. Tumor invasion of the cavernous sinus based on imaging criteria is a strong predictor of recurrence after surgery (33, 37,38). A tumor is regarded as *aggressive* if there is unusually rapid growth rate, earlier and more frequent recurrences requiring repeated surgeries, and clinically relevant growth despite optimal standard therapies (43). Thus, the terms *invasive* and *non-invasive* should refer only to imaging or morphological findings, and *aggressive* and *non-aggressive* pituitary adenomas to their clinical behavior (44).

The 2017 WHO classification recognizes pituitary tumors as having a low or high probability of recurrence (1). Of pituitary tumors requiring surgical treatment, 15% are 'aggressive' (37) but only 0.1-0.2% of these progress to become true carcinomas, as defined by evidence of cerebrospinal and/or systemic metastases (45-49). Histopathological and molecular markers are unable to identify the very rare tumors that exhibit aggressive growth or very rarely malignancy. In a study of 166 aggressive pituitary tumors, including 40 pituitary carcinomas, a classical proliferation index, such as Ki-67, did not distinguish aggressive pituitary tumors from pituitary carcinomas, and there was no significant difference in clinical parameters (50). No marker reliably seems to predict tumor behavior. Metalloproteinase 9 (MMP9) and PTTG correlate only with proliferative behavior (51). Whether tumor dedifferentiation represents an early pathogenic event or is a consequence of the carcinogenic process or intrapituitary signaling dysfunction is unknown (52,53).

NETs (referred to as carcinoids in the literature) of the pituitary are exceedingly rare. Few cases of primary pituitary NETs have been reported (54-57), as have eight cases of NET metastasis to the pituitary, most frequently of broncho-pulmonary origin (20). They can be distinguished from adenohypophyseal adenomas and other pituitary tumors by hormonal and transcription factor profiles.

## **Summation and Conclusions**

### ***Summary***

#### *Epidemiology*

Pituitary adenomas are common, predominantly indolent neoplasms, more than 99% of which do not affect life expectancy. By contrast, NETs are uncommon, apart from small tumors in appendiceal and rectal locations, and are potentially cancerous and reduce life expectancy.

#### *Taxonomy*

The developmental biology and regulatory function of the pituitary gland is that of a neuroendocrine gland. Anterior pituitary cells and cell types are identified by immunohistochemistry of hormones, transcription factors, and cytokeratin markers. Neuroendocrine markers commonly found in pituitary tumors are also expressed in other endocrine neoplasms including thyroid follicular and adrenal cortical adenomas that are pathologically distinct and not classified as NETs.

#### *Classification*

Pituitary adenomas can be classified by size, radiological grade, function, ultrastructure, cell type, and lineage. The 2017 WHO classification is based on immune detection of pituitary hormones, pituitary-specific transcription factors, and other cell differentiating co-factors. Renaming adenomas to NETs does not change prognostic prediction of pituitary neoplasms. Terminology change carries connotations that may affect patient disease perception and management, assigns an uncertain

malignancy designation to the vast majority of benign pituitary adenomas, and can adversely affect patient anxiety and overall disease management.

### *Aggressive Pituitary Neoplasms*

Presently validated histological markers do not reliably predict high-risk behavior of pituitary neoplasms.

### *Conclusions*

There is not yet a convincing argument for adopting the PitNET nomenclature but the question requires ongoing study and further discussion as new evidence emerges.

### *Epidemiology*

The clinical epidemiology, disease phenotype, management, and prognosis of pituitary neoplasms differ from most NETs.

### *Taxonomy*

Referring to a pituitary adenoma as a NET may be accurate from a developmental perspective. Studies of the molecular and genomic landscape to distinguish tissue types are required. Specific histological markers are required to distinguish neuroendocrine from endocrine neoplasms.

### *Classification*

Changes in classification should be justified by histological, pathological, and clinical evidence. A change to NET nomenclature does not guide identification of pituitary neoplasms that behave in a benign or aggressive manner. The proposed change



disproportionately portrays mostly benign pituitary neoplasms as aggressive or malignant, which may adversely affect patient well-being.

### *Aggressive Pituitary Neoplasms*

Further research is required to identify the very small number of indolent tumors that acquire aggressive and metastatic behavior.

### **Recommendations**

Following the PANOMEN workshop and review of the draft document, participants were asked: (1) whether the term tumor confers any advantage as a collective label for all pituitary neoplasms or to the subset of invasive pituitary adenomas; and (2) in the absence of rigorous pathological markers of high-risk behavior, whether imaging grades of invasiveness should be incorporated into a comprehensive pituitary classification and grading.

Seventy-nine percent (38/48) of authors recommend that the term “pituitary adenoma” not be replaced by “pituitary tumor,” and 58% (28/48) do not favor using the term “tumor” to designate the very small subset of invasive adenomas.

Given the absence of rigorous pathological markers of high-risk behavior, 65% (31/48) recommend that imaging grades be incorporated into a classification of pituitary neoplasms. Integrating imaging grade with pathological classification would likely benefit prognostication for clinical management.

It is recommended that the outcome of the PANOMEN workshop be communicated to relevant stakeholder professional societies, including those represented at this workshop and to the IARC/WHO for consideration in future classifications of endocrine tumors. It is recommended that this topic be revisited and the question of NET nomenclature be discussed

further as new evidence emerges. It is recommended that patient feedback be obtained before potential future nomenclature changes are considered.

### **Open Questions**

The workshop revealed controversy on issues which merit addressing in future workshops as new information emerges, including:

(1) Do primary pituitary NETs occur as an entity? Some participants were skeptical as to the validity of published reports supporting a diagnosis of primary pituitary neuroendocrine tumor.

(2) Is the difference between an endocrine and neuroendocrine cell conceptual or histological? Several participants questioned the specificity of traditional neuroendocrine markers (synaptophysin, chromogranin, neuron-specific enolase and somatostatin receptors) which are also expressed in follicular thyroid and adrenal cortical adenomas, neoplasms not regarded as of neuroendocrine origin.

Accepted Manuscript

## **Acknowledgements**

Drs. Chiara Villa, Olivera Casar-Borota, and Federico Roncaroli participated in the workshop and have declined authorship as they disagreed with the outcome.

The Pituitary Neoplasm Nomenclature Workshop was supported by unrestricted educational grants to the Pituitary Society from Chiasma, Inc., Corcept Therapeutics, Ipsen Biopharmaceuticals, Novo Nordisk, Pfizer Inc., Strongbridge Biopharma, and Tiburio Therapeutics. Funding sources had no role in data collection, analysis, and interpretation, decision to publish, or preparation of the manuscript. The authors thank Bari Laner and the Pituitary Society for coordinating the workshop.

## **Author contributions**

KH and SM served as Workshop Chairs. MM, PK, SC, HN, MBL, EL, LVS, TB, AG served as Speakers; MF, UK, RS, PT as Content Chairs; and MG, JW, FC, AB, MR as Session Chairs. All authors researched, reviewed, and commented on the data prior to and/or during the workshop. KH and SM prepared the manuscript drafts and all authors reviewed and approved the final manuscript.

## References

1. Lopes MBS. The 2017 World Health Organization classification of tumors of the pituitary gland: a summary. *Acta Neuropathol.* 2017; 134:521-535.
2. Asa SL, Casar-Borota O, Chanson P, Delgrange E, Earls P, Ezzat S, Grossman A, Ikeda H, Inoshita N, Karavitaki N, Korbonits M, Laws ER, Jr., Lopes MB, Maartens N, McCutcheon IE, Mete O, Nishioka H, Raverot G, Roncaroli F, Saeger W, Syro LV, Vasiljevic A, Villa C, Wierinckx A, Trouillas J, attendees of 14th Meeting of the International Pituitary Pathology Club AFN. From pituitary adenoma to pituitary neuroendocrine tumor (PitNET): an International Pituitary Pathology Club proposal. *Endocr Relat Cancer.* 2017; 24:C5-C8.
3. Ho KKY, Fleseriu M, Wass J, van der Lely A, Barkan A, Giustina A, Casanueva FF, Heaney AP, Biermasz N, Strasburger C, Melmed S. A tale of pituitary adenomas: to NET or not to NET : Pituitary Society position statement. *Pituitary.* 2019; 22:569-573.
4. Asa SL, Asioli S, Bozkurt S, Casar-Borota O, Chinezu L, Comunoglu N, Cossu G, Cusimano M, Delgrange E, Earls P, Ezzat S, Gazioglu N, Grossman A, Guaraldi F, Hickman RA, Ikeda H, Jaffrain-Rea ML, Karavitaki N, Kraljevic I, La Rosa S, Manojlovic-Gacic E, Maartens N, McCutcheon IE, Messerer M, Mete O, Nishioka H, Oz B, Pakbaz S, Pekmezci M, Perry A, Reiniger L, Roncaroli F, Saeger W, Soylemezoglu F, Tachibana O, Trouillas J, Turchini J, Uccella S, Villa C, Yamada S, Yarman S. Pituitary neuroendocrine tumors (PitNETs): nomenclature evolution, not clinical revolution. *Pituitary.* 2020; 23:322-325.
5. Ho KKY, Fleseriu M, Wass J, van der Lely A, Barkan A, Giustina A, Casanueva FF, Heaney AP, Biermasz N, Strasburger C, Melmed S. The tale in evolution: clarity,

- consistency and consultation, not contradiction and confusion. *Pituitary*. 2020; 23:476-477.
6. Molitch ME. Diagnosis and treatment of pituitary adenomas: a review. *JAMA*. 2017; 317:516-524.
  7. Melmed S. Pituitary-tumor endocrinopathies. *N Engl J Med*. 2020; 382:937-950.
  8. Hall WA, Luciano MG, Doppman JL, Patronas NJ, Oldfield EH. Pituitary magnetic resonance imaging in normal human volunteers: occult adenomas in the general population. *Ann Intern Med*. 1994; 120:817-820.
  9. Sanno N, Oyama K, Tahara S, Teramoto A, Kato Y. A survey of pituitary incidentaloma in Japan. *Eur J Endocrinol*. 2003; 149:123-127.
  10. Buurman H, Saeger W. Subclinical adenomas in postmortem pituitaries: classification and correlations to clinical data. *Eur J Endocrinol*. 2006; 154:753-758.
  11. Syro LV, Rotondo F, Ortiz LD, Kovacs K. 65 YEARS OF THE DOUBLE HELIX: Treatment of pituitary tumors with temozolomide: an update. *Endocr Relat Cancer*. 2018; 25:T159-T169.
  12. Agustsson TT, Baldvinsdottir T, Jonasson JG, Olafsdottir E, Steinhorsdottir V, Sigurdsson G, Thorsson AV, Carroll PV, Korbonits M, Benediktsson R. The epidemiology of pituitary adenomas in Iceland, 1955-2012: a nationwide population-based study. *Eur J Endocrinol*. 2015; 173:655-664.
  13. Tampourlou M, Fountas A, Ntali G, Karavitaki N. Mortality in patients with non-functioning pituitary adenoma. *Pituitary*. 2018; 21:203-207.
  14. Bolfi F, Neves AF, Boguszewski CL, Nunes-Nogueira VS. Mortality in acromegaly decreased in the last decade: a systematic review and meta-analysis. *Eur J Endocrinol*. 2018; 179:59-71.

15. Clayton RN, Jones PW, Reulen RC, Stewart PM, Hassan-Smith ZK, Ntali G, Karavitaki N, Dekkers OM, Pereira AM, Bolland M, Holdaway I, Lindholm J. Mortality in patients with Cushing's disease more than 10 years after remission: a multicentre, multinational, retrospective cohort study. *Lancet Diabetes Endocrinol.* 2016; 4:569-576.
16. Graversen D, Vestergaard P, Stochholm K, Gravholt CH, Jorgensen JO. Mortality in Cushing's syndrome: a systematic review and meta-analysis. *Eur J Intern Med.* 2012; 23:278-282.
17. Papakokkinou E, Olsson DS, Chantzichristos D, Dahlqvist P, Segerstedt E, Olsson T, Petersson M, Berinder K, Bensing S, Hoybye C, Eden-Engstrom B, Burman P, Bonelli L, Follin C, Petranek D, Erfurth EM, Wahlberg J, Ekman B, Akerman AK, Schwarcz E, Bryngelsson IL, Johannsson G, Ragnarsson O. Excess morbidity persists in patients with Cushing's disease during long-term remission: a Swedish nationwide study. *J Clin Endocrinol Metab.* 2020; 105:dga291.
18. Oberndorfer S. Karzinoide Tumoren des Dünndarms. *Frankf Z Pathol.* 1907; 1:425-429.
19. Osamura R, Grossman A, Korbonits M, Kovacs K, Lopes M, Matsuno A, Trouillas J. Pituitary gland: pituitary adenoma. In: Lloyd RV, Osamura RY, Klöppel G, J R, eds. *WHO Classification of Tumours of Endocrine Organs.* Vol 10. 4th ed. Lyon: IARC; 2017:14-18.
20. Moshkin O, Rotondo F, Scheithauer BW, Soares M, Coire C, Smyth HS, Goth M, Horvath E, Kovacs K. Bronchial carcinoid tumors metastatic to the sella turcica and review of the literature. *Pituitary.* 2012; 15:160-165.

21. Dasari A, Shen C, Halperin D, Zhao B, Zhou S, Xu Y, Shih T, Yao JC. Trends in the incidence, prevalence, and survival outcomes in patients with neuroendocrine tumors in the United States. *JAMA Oncol.* 2017; 3:1335-1342.
22. Yao JC, Hassan M, Phan A, Dagohoy C, Leary C, Mares JE, Abdalla EK, Fleming JB, Vauthey JN, Rashid A, Evans DB. One hundred years after "carcinoid": epidemiology of and prognostic factors for neuroendocrine tumors in 35,825 cases in the United States. *J Clin Oncol.* 2008; 26:3063-3072.
23. Eriksson J, Norlen O, Ogren M, Garmo H, Ihre-Lundgren C, Hellman P. Primary small intestinal neuroendocrine tumors are highly prevalent and often multiple before metastatic disease develops. *Scand J Surg.* 2019:1457496919874484.
24. Ikeda H, Suzuki J, Sasano N, Niizuma H. The development and morphogenesis of the human pituitary gland. *Anat Embryol (Berl).* 1988; 178:327-336.
25. Cheung LYM, Camper SA. Hypophysis development and disease in humans. *Reference Module in Biomedical Sciences.* 2018; doi: <https://doi.org/10.1016/B978-0-12-801238-3.65809-9>.
26. Nishioka H, Inoshita N, Mete O, Asa SL, Hayashi K, Takeshita A, Fukuhara N, Yamaguchi-Okada M, Takeuchi Y, Yamada S. The complementary role of transcription factors in the accurate diagnosis of clinically nonfunctioning pituitary adenomas. *Endocr Pathol.* 2015; 26:349-355.
27. Satoh F, Umemura S, Yasuda M, Osamura RY. Neuroendocrine marker expression in thyroid epithelial tumors. *Endocr Pathol.* 2001; 12:291-299.
28. Haak HR, Fleuren GJ. Neuroendocrine differentiation of adrenocortical tumors. *Cancer.* 1995; 75:860-864.

29. Zhang D, Hugo W, Donohue W, Girgis M, Bergsneider M, Wang MB, Heaney AP. Pituitary and neuroendocrine tumors exhibit distinct transcriptomes on single cell RNA sequencing. *J Endocr Soc.* 2020; 4:Abstract SAT-314.
30. Cushing H. *The Pituitary Body and its Disorders*. Philadelphia, PA: Lippincott; 1912.
31. Hardy J. Transphenoidal microsurgery of the normal and pathological pituitary. *Clin Neurosurg.* 1969; 16:185-217.
32. Horvath E, Kovacs K. Fine structural cytology of the adenohypophysis in rat and man. *J Electron Microsc Tech.* 1988; 8:401-432.
33. Knosp E, Steiner E, Kitz K, Matula C. Pituitary adenomas with invasion of the cavernous sinus space: a magnetic resonance imaging classification compared with surgical findings. *Neurosurgery.* 1993; 33:610-618.
34. Zada G, Woodmansee WW, Ramkissoon S, Amadio J, Nose V, Laws ER, Jr. Atypical pituitary adenomas: incidence, clinical characteristics, and implications. *J Neurosurg.* 2011; 114:336-344.
35. Meij BP, Lopes MB, Ellegala DB, Alden TD, Laws ER, Jr. The long-term significance of microscopic dural invasion in 354 patients with pituitary adenomas treated with transsphenoidal surgery. *J Neurosurg.* 2002; 96:195-208.
36. Neou M, Villa C, Armignacco R, Jouinot A, Raffin-Sanson ML, Septier A, Letourneur F, Diry S, Diedisheim M, Izac B, Gaspar C, Perlemoine K, Verjus V, Bernier M, Boulin A, Emile JF, Bertagna X, Jaffrezic F, Laloe D, Baussart B, Bertherat J, Gaillard S, Assie G. Pangenomic classification of pituitary neuroendocrine tumors. *Cancer Cell.* 2020; 37:123-134.
37. Trouillas J, Roy P, Sturm N, Dantony E, Cortet-Rudelli C, Viennet G, Bonneville JF, Assaker R, Auger C, Brue T, Cornelius A, Dufour H, Jouanneau E, Francois P, Galland F, Mougel F, Chapuis F, Villeneuve L, Maurage CA, Figarella-Branger D,



- Raverot G, members of H, Barlier A, Bernier M, Bonnet F, Borson-Chazot F, Brassier G, Caulet-Maugendre S, Chabre O, Chanson P, Cottier JF, Delemer B, Delgrange E, Di Tommaso L, Eimer S, Gaillard S, Jan M, Girard JJ, Lapras V, Loiseau H, Passagia JG, Patey M, Penfornis A, Poirier JY, Perrin G, Tabarin A. A new prognostic clinicopathological classification of pituitary adenomas: a multicentric case-control study of 410 patients with 8 years post-operative follow-up. *Acta Neuropathol.* 2013; 126:123-135.
38. Raverot G, Dantony E, Beauvy J, Vasiljevic A, Mikolasek S, Borson-Chazot F, Jouanneau E, Roy P, Trouillas J. Risk of recurrence in pituitary neuroendocrine tumors: a prospective study using a five-tiered classification. *J Clin Endocrinol Metab.* 2017; 102:3368-3374.
39. Asioli S, Righi A, Iommi M, Baldovini C, Ambrosi F, Guaraldi F, Zoli M, Mazzatenta D, Faustini-Fustini M, Rucci P, Giannini C, Foschini MP. Validation of a clinicopathological score for the prediction of post-surgical evolution of pituitary adenoma: retrospective analysis on 566 patients from a tertiary care centre. *Eur J Endocrinol.* 2019; 180:127-134.
40. Doust JA, Bell KJL, Glasziou PP. Potential Consequences of Changing Disease Classifications. *JAMA.* 2020; 323:921-922.
41. Dixon PR, Tomlinson G, Pasternak JD, Mete O, Bell CM, Sawka AM, Goldstein DP, Urbach DR. The role of disease label in patient perceptions and treatment decisions in the setting of low-risk malignant neoplasms. *JAMA Oncol.* 2019; 5:817-823.
42. Chadha NK, Repanos C. Patients' understanding of words used to describe lumps: a cross-sectional study. *J Laryngol Otol.* 2006; 120:125-128.
43. Raverot G, Burman P, McCormack A, Heaney A, Petersenn S, Popovic V, Trouillas J, Dekkers OM, European Society of E. European Society of Endocrinology clinical

- practice guidelines for the management of aggressive pituitary tumours and carcinomas. *Eur J Endocrinol*. 2018; 178:G1-G24.
44. Di Ieva A, Rotondo F, Syro LV, Cusimano MD, Kovacs K. Aggressive pituitary adenomas--diagnosis and emerging treatments. *Nat Rev Endocrinol*. 2014; 10:423-435.
  45. Dekkers OM, Karavitaki N, Pereira AM. The epidemiology of aggressive pituitary tumors (and its challenges). *Rev Endocr Metab Disord*. 2020; 21:209-212.
  46. Heaney AP. Clinical review: Pituitary carcinoma: difficult diagnosis and treatment. *J Clin Endocrinol Metab*. 2011; 96:3649-3660.
  47. Kasuki L, Raverot G. Definition and diagnosis of aggressive pituitary tumors. *Rev Endocr Metab Disord*. 2020; 21:203-208.
  48. Saeger W, Ludecke DK, Buchfelder M, Fahlbusch R, Quabbe HJ, Petersenn S. Pathohistological classification of pituitary tumors: 10 years of experience with the German Pituitary Tumor Registry. *Eur J Endocrinol*. 2007; 156:203-216.
  49. Yoo F, Kuan EC, Heaney AP, Bergsneider M, Wang MB. Corticotrophic pituitary carcinoma with cervical metastases: case series and literature review. *Pituitary*. 2018; 21:290-301.
  50. McCormack A, Dekkers OM, Petersenn S, Popovic V, Trouillas J, Raverot G, Burman P, ESE Survey Collaborators. Treatment of aggressive pituitary tumours and carcinomas: results of a European Society of Endocrinology (ESE) survey 2016. *Eur J Endocrinol*. 2018; 178:265-276.
  51. Melmed S. Pathogenesis of pituitary tumors. *Nat Rev Endocrinol*. 2011; 7:257-266.
  52. Ben-Shlomo A, Deng N, Ding E, Yamamoto M, Mamelak A, Chesnokova V, Labadzhyan A, Melmed S. DNA damage and growth hormone hypersecretion in pituitary somatotroph adenomas. *J Clin Invest*. 2020 Jul 16; 138:540.

53. Srirangam Nadhamuni V, Korbonits M. Novel insights into pituitary tumorigenesis: genetic and epigenetic mechanisms. *Endocr Rev.* 2020; 41:bnaa006.
54. Liu H, Wang H, Qi X, Yu C. Primary intracranial neuroendocrine tumor: two case reports. *World J Surg Oncol.* 2016; 14:138.
55. Nasi D, Perano D, Ghadirpour R, Iaccarino C, Servadei F, Romano A. Primary pituitary neuroendocrine tumor: Case report and literature review. *Surg Neurol Int.* 2017; 8:101.
56. Lynggard LA, Nielsen EH, Laurberg P. Carcinoid syndrome caused by a serotonin-secreting pituitary tumour. *Eur J Endocrinol.* 2014; 170:K5-9.
57. Bhambhani A, Dugar M, Rao J, Prasad R. Pituitary carcinoid coexisting with systemic lupus erythematosus: a rare combination. *Natl Med J India.* 2016; 29:209-211.

Accepted Manuscript



