1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Increasing the longitudinal bending stiffness of runners’ habitual shoes: an appropriate choice for improving running performance?

Submission type: Original investigation

Full names of the authors: Nicolas Flores^{ab*}, Guillaume Rao^b, Eric Berton^b, Nicolas Delattre^a

Institutional/corporate affiliations: ^aDecathlon SportsLab, Movement Sciences Department, Villeneuve d’Ascq, France; ^bAix Marseille Univ, CNRS, ISM, Marseille, France

Contact details for the corresponding author: *Nicolas Flores, Decathlon SportsLab, nicolas.flores@decathlon.com, +33(0)760726430

Preferred running head: Bending stiffness and prolonged running

Abstract word count: 221 words

Text-only word count: 5785 words

Number of figures: 4

Number of tables: 2

Abstract

Purpose: This study analysed the effects of increasing the longitudinal bending stiffness (LBS) of runners’ habitual shoes on the metabolic energetic demand, lower limb muscle activations, and stride spatiotemporal parameters during a prolonged running session through classical group investigation and a more individualised approach.

Methods: Eleven recreational male participants ran overground for 40 min at 95% of their ventilatory anaerobic threshold with low (participants' own shoes) or high LBS shoes (stiff carbon plate inserted under insole). The net energetic cost of running, lower leg muscle activations, and spatiotemporal parameters were measured during the prolonged running. The variables of interest were analysed for one minute in seven time intervals, which were five minutes apart.

Results: There were no main effects of LBS or interaction effects with running duration on the group averaged variables of interest. Overall, the participant-specific metabolic effects induced by an increased shoe LBS were not beneficial. Beneficial metabolic effects were more able to occur when the increased LBS decreased (or did not increase) the ground contact time relative to their habitual shoes, and for the taller runners.

Conclusions: Increasing the LBS in runners' habitual shoes did not induce systematic metabolic effects to all the runners. This study suggested that a shoe LBS intervention was likely not beneficial for performance purposes if the runners' shoe habits were too much disrupted.

Keywords: footwear, running shoes, shoe flexibility, longitudinal bending stiffness, running economy, subject-specific, performance.

Introduction

Since the first published scientific investigation twenty years ago,¹ the shoe longitudinal bending stiffness (LBS) has recently experienced a resurgence of interest in the scientific and industrial community, and sports competitions due to its potential beneficial effects for endurance running. The turning point in the footwear community occurred in 2017 when large benefits on the metabolic energetic demand to cover a given distance at a given speed (i.e.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

improved running economy) was shown with Nike VaporFly 4% shoes.^{2–5} These shoes were designed with a substantial energy return midsole foam, and stiff curved carbon plates inserted into the midsoles to increase the LBS. Several personal and world race records have been broken in runners wearing these shoes,^{6,7} thus generating a lot of debates about potential unfair advantages brought by the shoes.^{8–11} This has recently led the World Athletics to change shoe regulations. In road events, the maximum sole thickness allowed is actually 40 mm and only one rigid plate can be embedded in the sole. Nowadays, almost every running footwear company proposes one performance shoe model equipped with a stiff carbon plate. Taken together, these events suggest that the presence of stiff carbon plates in the shoes really makes beneficial differences on running economy and performance. Do the scientific studies support this suggestion?

The initial study that paved the way for the presence of carbon plate in running shoes has reported an improvement of running economy (expressed in consumed millilitre of oxygen per kilogram per minute) by 1% on average in a group of runners with an intermediate shoe LBS compared to more flexible and stiffer shoes.¹² Since then no systematic group differences were found in the subsequent studies. Five studies did not show any significant effect on running economy of an increased shoe LBS compared to more flexible shoes^{13–15} or over a range of different LBS,^{16,17} and one study showed a significant poorer running economy with an increased LBS compared to more flexible shoes¹⁸ (see Ortega et al.¹⁹ for an exhaustive review on this topic). These recent results suggest that there is no unique shoe LBS that induces running economy benefits systematically to all the runners, and the effects may even be detrimental.

1 The initial study of shoe LBS was also a pioneer in suggesting participant-specific
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 The initial study of shoe LBS was also a pioneer in suggesting participant-specific
2 optimisations of the LBS. Indeed, this study showed that heavier runners benefited more from
3 an intermediate increase of shoe LBS.¹² This result has led the following studies to consider
4 the participant-specific effects in their analyses. Oh and Park²⁰ showed an individual optimal
5 LBS for the running economy according to the stiffness of the runner's metatarsophalangeal
6 (MTP) joint, thus several LBS were suitable to several groups of runners with similar MTP
7 stiffnesses. McLeod et al.²¹ and Čigoja et al.¹⁷ identified an individual optimal LBS based on
8 running economy measurements but without highlighting any explanatory variables. Madden
9 et al.¹³ and Flores et al.²² reported participant-specific benefits of an increased shoe LBS
10 associated with biomechanical changes induced by an increased LBS. It was previously
11 reported that reduced distal lower limb muscle activation during the braking phase were
12 associated with greater ankle dorsiflexion.²² This study suggested that this may enhance the
13 energy storage in the elastic components that can be returned economically during push-off.
14 In line with this, an increased LBS have been shown to induce lower ankle plantarflexion
15 amplitude, slower ankle plantarflexion velocity, and greater Achilles tendon energy return
16 during push-off,^{13,17,22,23} which could contribute to increase performance. While the
17 participant-specific benefits induced by an increased shoe LBS are well acknowledged, the
18 individual characteristics and/or the running biomechanical changes associated with metabolic
19 improvements remain disputed and not systematic. Indeed, the above-mentioned studies did
20 not find the same variables explaining the metabolic benefits induced by an increased LBS
21 (for example, the runners' mass has never been highlighted as an explanatory variable in the
22 subsequent studies).

23 These previous results have exclusively been shown during short running durations.
24 Whether the effects of the shoe LBS are metabolically beneficial or not during a prolonged

1
2
3
4 1 run remain unknown. This question was recently partially addressed,²⁴ by showing that an
5
6 2 increased shoe LBS induced a delay in the distal-to-proximal work redistribution during
7
8 3 prolonged running. These authors suggested that using proximal muscles with greater volume
9
10 4 (i.e. *quadriceps*) should be more detrimental than using distal muscles with lower volume (i.e.
11
12 5 *triceps surae*).²⁵ Because the running economy was not measured in this previous study,²⁴
13
14 6 the metabolic benefits attributed to an increased shoe LBS during prolonged running remain
15
16 7 speculative.

17
18
19
20
21 8 While a lot of useful and relevant information is available regarding the LBS effects
22
23 9 on the metabolic demand and the running biomechanics during short duration,¹⁹ only
24
25 10 assumptions can be made concerning the metabolic effects of the LBS during longer running
26
27 11 exposure where only the running biomechanics was measured.²⁴ The current study aimed to
28
29 12 investigate these previous independent findings in the same study. The purpose was to analyse
30
31 13 the effects of increasing the longitudinal bending stiffness of runners' habitual shoes on
32
33 14 metabolic demand, lower limb muscle activations, and stride spatiotemporal parameters
34
35 15 during a prolonged running session through classical group investigation and a more
36
37 16 individualised approach. Based on the assumption from Cigoja et al.²⁴, it was hypothesised
38
39 17 that an increased shoe LBS would limit and/or delay the increase in metabolic energetic
40
41 18 demand. As highlighted in numerous studies,^{12,13,17,20–22} it was expected that the benefits
42
43 19 induced by an increased shoe LBS would also be participant-specific, especially according to
44
45 20 individual characteristics and/or biomechanical changes induced by an increased LBS relative
46
47 21 to their shoe habits.

48
49
50
51
52
53
54
55
56
57
58
59 23 **Methods**
60

Participants

Eleven male novice runners (Table 1), stating running at least once a week and being able to run for one hour at least at 10 km/h, volunteered to participate in the experiment. All participants gave their written informed consent to all the procedures performed during the experiment that were in accordance with the university ethical rules. The participants stated being healthy and free from injury in the previous year.

Shoe conditions

Two shoe conditions were used in this study. The control shoe condition was the participants' own running shoes (Table 1 and Supplementary material A), that they used during their usual running activity (low LBS). The use of participants' own running shoes should permit observation of their habitual running biomechanics without being influenced by unfamiliar shoe features²⁶ other than the LBS investigated in the study. The second shoe condition was obtained by the addition of thin (0.9 mm) stiff straight carbon plates under the insoles of the participant's own running shoes (high LBS). One size of carbon plates has been used, and we ensured that the plates were functionally well inserted in all the participants' shoes to not induce discomfort during running. The shoe LBS (Table 1) was measured with a mechanical manual bending test (see Supplementary material B for full details about the procedure) based on the study of Worobets and Wannop²⁷. The high LBS condition was on average $263 \pm 89\%$ stiffer than the low LBS. To answer to the current study's purpose concerning the choice to increase the LBS in runners' habitual shoes (with associated advantages and disadvantages), the shoe mass was not equalised between both shoe conditions. Thus the high LBS conditions (317 g on average) were 20 g per shoe heavier due to the carbon plate than the low LBS conditions (297 g on average).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 *** Table 1 near here ***

2 *Experimental protocol*

3 The experiment was composed of three sessions performed on three different days with at
4 least 48 hours rest between each session. From the welcome of the participant to the end of
5 the session, the first session lasted 1 hour and both subsequent sessions each lasted 2 hours.
6 On average, the three sessions were performed within 11 days.

7 In the first session, an incremental test was performed. The test started with a 5-min
8 warm-up of running at 1.94 m/s with participants' own running shoes on a treadmill (Saturn
9 300/100r, h/p/cosmos, Germany) with 1% slope gradient. The speed was then increased by
10 0.28 m/s every minute. An open-circuit expired-gas analysis system (K5, Cosmed, Italy)
11 recorded breath-by-breath the rate of oxygen uptake ($\dot{V}O_2$) and the rate of carbon dioxide
12 production ($\dot{V}CO_2$). The device was calibrated before each test with a reference gas
13 containing known concentrations of oxygen and carbon dioxide, using a 3.0 L syringe to
14 ensure an accurate volume air measurement. The incremental running test aimed to identify
15 the ventilatory anaerobic threshold (VAT) for each participant. The VAT was identified
16 through the onset increase of ventilation versus oxygen uptake ratio ($\dot{V}_E/\dot{V}O_2$) and end-tidal
17 partial pressure of oxygen ($P_{ET}O_2$), in the presence of still decreasing or constant ventilation
18 versus carbon dioxide production ratio ($\dot{V}_E/\dot{V}CO_2$) and $P_{ET}CO_2$.²⁸ The speed equal to 95% of
19 the speed that elicited VAT was kept for each participant and was imposed in both subsequent
20 sessions of prolonged running.

21 The subsequent sessions were performed on the field to consider realistic road running
22 conditions while keeping as much as possible of the data quality. The participants ran for a
23 prolonged duration with either the low LBS or the high LBS shoe condition, in a randomised
24 order. Due to potential fatigue accumulation, a prolonged running session was dedicated to

only one shoe condition, thus the low and high LBS shoe conditions were performed during separate sessions. Each participant performed both sessions at the same time of day. In each session, the experimental protocol was identical. The participants were equipped with the expired-gas analysis system, which was calibrated before each session as described above, and surface electromyograms (EMG; Trigno Avanti, Delsys, Massachusetts, USA), which were used to record muscle activations at 1926 Hz. After skin preparation, the EMG electrodes were located according to the recommendations of surface electromyography for the non-invasive assessment of muscles (SENIAM) at the level of superficial muscles *vastus lateralis* (VL), *gastrocnemius medialis* (GM), and *tibialis anterior* (TA). The SENIAM procedure for sensor placement was reproduced as accurately as possible by the same researcher between both sessions. The three sensors were firmly secured with a medical adhesive strap to avoid any artifacts due to sweating, vibrations, displacement, or detachment of the sensors throughout the entire duration of the experiments. The EMG sensors were connected to a datalogger (Trigno Personal Monitor, Delsys, Massachusetts, USA) located in an armband, thus allowing wireless EMG acquisitions during the outdoor prolonged run. A watch (Fenix 3, Garmin, Kansas, USA) paired with an accelerometer mounted on a heart rate strap (HRM-Run, Garmin, Kansas, USA) was used to measure accelerations of the upper body from which were computed the stride cadence, the ground contact time, and the vertical oscillation of the upper body.²⁹ To obtain day-specific respiratory reference values, the participants first performed 5 min sitting at rest on a chair without moving or speaking. Finally, the participants ran continuously for 40 min outdoor over a loop with hard concrete surface (Supplementary material C) at the targeted speed identified in the first session (imposed speed visually controlled with ground marks in each 20 m and a metronome). Gas exchanges data, EMG data, and spatiotemporal data were recorded during the whole duration of the experiment.

Data analysis

1
2
3 1 The VO_2 , the VCO_2 , the respiratory exchange ratio (RER), and the ambient temperature were
4
5 2 measured from the open-circuit expired-gas analysis system. For each subject, the averaged
6
7 3 VO_2 and the VCO_2 during the last minute of the sitting rest were subtracted from those
8
9 4 measured during the prolonged run to offset the day-specific respiratory reference values of
10
11 5 the participants between both sessions. Then, the net metabolic energetic cost of running was
12
13 6 computed from caloric equivalent based on the Péronnet and Massicotte³⁰ equation and
14
15 7 expressed in net metabolic power (W/kg). The net energetic cost of running, the RER, and the
16
17 8 ambient temperature were averaged during 1 min for seven time intervals: from 9 to 10 min
18
19 9 (baseline measurement), from 14 to 15 min, from 19 to 20 min, from 24 to 25 min, from 29 to
20
21 10 30 min, from 34 to 35 min, and from 39 to 40 min.
22
23
24
25
26

27 11 The EMG data were band-pass filtered [10-400 Hz] with a zero time-lag fourth order
28
29 12 Butterworth filter to remove signal noise and artifacts. Then, these data were rectified and
30
31 13 low-pass filtered at 15 Hz with a critically damped filter to smooth the signal.³¹ For each
32
33 14 muscle, activations were normalised by the maximal activation recorded in the seven time
34
35 15 intervals during the session. For each time interval, this normalised EMG signal was
36
37 16 integrated during the entire minute. Because of EMG sensor location (on muscles, thus on
38
39 17 wobbling masses) and the range of accelerometry measurements from EMG sensors (± 2 g),
40
41 18 the accelerometer signal was too noisy to accurately identify heel strike and toe off events.
42
43 19 The integration of EMG activations thus took into account both aerial and stance phases. Data
44
45 20 of GM activations is missing for participant 3 in the low LBS condition and participant 10 in
46
47 21 both LBS conditions, likely due to the fact that the left foot in the aerial phase hit the
48
49 22 supporting right leg where the GM sensor was located, thus disconnecting this sensor. A
50
51 23 firmer protection than the medical strap or the recording of another *triceps surae* muscle
52
53 24 would enable avoiding such an issue.
54
55
56
57
58
59
60

Raw data measured by the torso strap sensor were saved in the watch and then imported on Garmin Connect software. This software automatically computed the stride cadence, ground contact time, and vertical oscillation of the upper body from raw data, which were saved on '.fit' files. These '.fit' files were imported on Golden Cheetah software (v3.3.0) to extract the computed data at each second of the prolonged running in a more conventional '.txt' format. The flight time (in seconds) was computed from the measured stride cadence and ground contact time (in seconds), and the vertical and leg stiffnesses were also computed based on ground contact and flight times.³² The stride cadence, the ground contact time, the vertical oscillation of upper body, the vertical stiffness, and the leg stiffness were averaged during 1 min for the seven time intervals. Data of stride spatiotemporal parameters is missing for participant 1 in the low LBS condition and participant 4 in the high LBS condition due to an issue while saving raw data.

Individual data were then reorganised to allow the investigation of the individual beneficial or detrimental metabolic effects induced by the increase of the LBS in the participant's habitual shoes during a prolonged run. For each participant, the metabolic, EMG, and stride spatiotemporal data induced by the high LBS condition were subtracted to those induced by the low LBS condition. This procedure has been done for the seven time intervals for all the variables. At a given time interval, negative or positive values thus respectively indicated that an increased LBS in the participant's habitual shoes induced decreased (i.e. better running economy) or increased (i.e. poorer running economy) net energetic cost of running, and decreased or increased muscle activations and spatiotemporal responses, compared to a lower habitual LBS.

Statistical analysis

1
2
3 1 Due to some issues during the experiments, statistical analyses were performed on 9
4
5 2 participants for the GM activations and spatiotemporal parameters, and on 11 participants for
6
7 3 all other variables.
8
9

10
11 4 *Classical group-based investigation*
12
13

14 5 ANOVA assumptions (normality and homogeneity of residuals) were met for the VL
15
16 6 activations, the cadence, and the ground contact time, thus a two-way repeated measures
17
18 7 ANOVA ($\alpha=0.05$) was used to evaluate the main effects of 'LBS' and 'running duration'
19
20 8 factors, and the interaction of these factors, on these variables. All other variables were non-
21
22 9 parametrically analysed with a permutation procedure ($\alpha=0.05$), which was performed
23
24 10 through the Matlab open-source code (v.0.4.3, <http://www.spm1d.org>). The permutation
25
26 11 procedure consists of calculating the test statistic F values for a large number of re-orderings
27
28 12 (i.e. permutations) of the observations (i.e. data). For example in our data, one permutation
29
30 13 consists in permuting an observation induced by a given shoe condition for a given participant
31
32 14 at a given time interval with another observation induced by the other shoe condition for a
33
34 15 given participant at a given time interval. For each permutation, the test statistic F value is
35
36 16 computed. This results in a number of F values equal to the total number of permutations,
37
38 17 which tends to generate a Gaussian distribution of the frequency of F values. Then, the critical
39
40 18 test statistic threshold (F^*) is computed as the 95th percentile of this distribution to ensure that
41
42 19 only $\alpha=5\%$ of all permutations exceed the F^* . If the F value computed from the original
43
44 20 observations exceed F^* , this provides sufficient evidence to reject the null hypothesis (i.e.
45
46 21 observing a significant effect).³³ The partial omega squared (ω_p^2) effect size was finally
47
48 22 computed and interpreted as: trivial effect $\leq 0.02 <$ small effect $\leq 0.13 <$ medium effect ≤ 0.26
49
50 23 $<$ large effect.³⁴
51
52
53
54
55
56
57
58
59
60

When a significant main effect of running duration or the interaction between the LBS and the running duration was observed, post-hoc pairwise comparisons were made between the baseline and subsequent time intervals through the 'emmeans' package on RStudio (v.3.4.1, Massachusetts, USA). This resulted in six post hoc comparisons, thus the alpha level was corrected as $\alpha=0.05/6=0.008$.

All the statistical results (two main factors and interaction) are presented in the results section, but for the purpose of the study, only those concerning the LBS main factor and the interaction between the LBS and the running duration are discussed.

Individual approach

A correlation approach was used in an exploratory way by screening for potential candidate variables partially involved in the variation of the net energetic cost of prolonged running when the shoe LBS was investigated. This enabled to understand in which conditions a given individual was more able to benefit from a shoe LBS intervention.

Pearson correlations were performed, at the first and the last time interval of the prolonged run, between the individual variations of the net energetic cost of running induced by an increased LBS and ten potential explanatory variables. These potential explanatory variables were the individual variations of the cadence, the body vertical oscillation, and the ground contact time induced by an increased LBS, the individual variations of the TA, the GM, and the VL activations induced by an increased LBS, the participants' mass, height, speed and the relative increase of the shoe LBS.

The correlations kept for results and discussion sections should combine $p<0.1$ with a large effect size. A large effect size was identified by a Pearson regression coefficient (r) lower than -0.5 or greater than 0.5.³⁵

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

The speed equal to 95% of the speed that elicited VAT in the incremental test, which was imposed during the prolonged running, was 3.11 ± 0.19 m/s on average (Table 1). The RER was lower than 1 for all the participants indicating that the aerobic metabolism was the main contributor of the energetic cost of running. The ambient temperature was not significantly different between both sessions (low LBS session: $26.9\pm0.2^{\circ}\text{C}$; high LBS session: $27.3\pm0.2^{\circ}\text{C}$; $p=0.704$).

*** Figure 1 near here ***

Classical group-based investigation

No significant main effect of LBS or interaction effect of LBS and running duration was observed on any of the variables ($p>0.05$; Figure 1, Figure 2, and Figure 3).

The running duration induced a significant decrease in TA activations ($F_{6,60}=4.18$; $p=0.003$; $\omega_p^2=0.22$, medium effect; Figure 2), where TA activations were significantly lower from 39 to 40 min compared to baseline from 9 to 10 min ($p=0.006$). No significant main effect of the running duration was observed on all other variables ($p>0.05$; Figure 1, Figure 2, and Figure 3).

*** Figure 2 near here ***

*** Figure 3 near here ***

Individual approach

The simplest view of the individual metabolic results (both last lines in Table 2) showed that an increased LBS in the runners' habitual shoes was detrimental for most of the participants (7 vs. 4) in two time intervals at the beginning and in two time intervals at the end

of the run, whereas in three middle time intervals of the run the number of participants benefiting from an increased LBS was slightly greater (6 vs. 5). Furthermore, 2 runners always benefited from an increased LBS of their habitual shoes (participants 2 and 4), 4 runners always benefited from their habitual shoes (participants 8, 9, 10, and 11), and 5 runners randomly benefited either from their habitual shoes or an increased LBS throughout the prolonged running duration (right column in Table 2). This non-homogeneous participants' responses was revealed by a non-significant Chi-Square test ($\chi^2=0.09$; $p>0.05$) indicating that the number of participants benefiting from the low or high LBS ($n=6$) or those unclear ($n=5$) was not different than what would be expected if there was no effect of shoe LBS (50% of beneficiaries and 50% of unclear in the sample of participants). This result strengthens the individual approach to highlight in which conditions a given individual was more able to benefit from a shoe LBS intervention.

*** Table 2 near here ***

Based on the criteria set in the statistical section, four significant correlations were identified during the prolonged running (Figure 4; see Supplementary Material D for more results).

At baseline (from 9 to 10 min), the individual variation of the net energetic cost of running induced by an increased LBS was correlated with the individual variation of the ground contact time ($p=0.067$; $r=0.63$), the individual variation of the body vertical oscillation ($p=0.070$; $r=-0.63$), and the individual variation of TA activations ($p=0.094$; $r=-0.53$) induced by an increased LBS.

At the last time interval (from 39 to 40 min), the individual variation of the net energetic cost of running induced by an increased LBS was correlated with the height of the participants ($p=0.019$; $r=-0.69$).

*** Figure 4 near here ***

Discussion

For the first time, this study investigated the effects of increasing the LBS of runners’ habitual shoes on metabolic demand, lower limb muscle activations, and stride spatiotemporal parameters during a prolonged running session, through classical group investigation and a more individualised approach. Over the mean group of participants, the shoe LBS had no effect on the metabolic energetic demand. Nevertheless, participant-specific metabolic energetic responses to an increased LBS have been highlighted. Depending on the time interval of the prolonged run, these participant-specific metabolic responses were correlated with the variations of spatiotemporal parameters or muscle activations induced by an increased LBS, and with the participant’s height.

Classical group-based investigation

An increase of the LBS in runners’ habitual shoes did not induce a significant effect on the net energetic cost during the prolonged running. This result was not in line with the initial hypothesis, which was based on the only one study²⁴ that has investigated the shoe LBS effect during prolonged running. By measuring a delay in the distal-to-proximal joint work redistribution during a prolonged submaximal run under the anaerobic threshold, this previous study suggested that an increased shoe LBS might have beneficial repercussions on the metabolic demand. At similar running intensity (95% of the VAT), we did not observe such supposed metabolic benefits on the mean group of participants. This running intensity may have been too low to increase the net energetic cost of prolonged running (i.e. inducing fatigue). As previously recommended,³⁶ the used running intensity allowed an appropriate

1 calculation of the energetic cost of running, while taking account the aerobic metabolism as
2 the main contributor to the metabolic demand ($RER < 1$). If an increased shoe LBS really acts
3 in a beneficial way by limiting or delaying the onset of the energetic cost increase, this
4 putative beneficial effect could be observed in presence of fatigue. Future studies using
5 similar running intensity than in the current one should thus consider longer running duration
6 (> 40 min) to potentially induce VO_2 drift and thus fatigue. As it is, the lack of average effect
7 of the shoe LBS on the metabolic demand was in line with almost all the previous studies
8 using shorter duration of running.^{13–17}

9 All the variables were analysed from a baseline set to 9 min of running (about 1.7 km
10 based on the imposed speed of 3.11 m/s on average). This baseline was set to only measure an
11 eventual effect of the LBS in the duration of the prolonged running, while avoiding measuring
12 the adaptation effects due to reaching a steady-state VO_2 ,³⁷ a stable biomechanical
13 pattern,^{38,39} a stable neuromuscular system,⁴⁰ or stable mechanical shoe properties.⁴¹
14 Likely stabilised following this adaptation period, all the measured variables were unaffected
15 by the shoe LBS. This is different from what can be expected from previous studies as an
16 increased shoe LBS was known to have several influences on the running biomechanics.¹⁹
17 Therefore, at the group level, further assumptions can be made from the current prolonged
18 running. The physiological and biomechanical effects of an increased LBS may not occur at
19 all, or occur too early during this adaptation period, or be too subtle to be measured with
20 enough resolution on the field, or may not be systematic between the participants.

21 ***Individual approach***

22 The time-courses of all the measured variables during the prolonged running were
23 highly participant-specific within-session and between-session (Figure 1, Figure 2, and Figure

1
2
3 1 3). This lack of non-systematic individual responses might be one explanation to the non-
4
5 2 significant effects of the shoe LBS at the group level. The observed individual variability was
6
7 3 in line with several studies highlighting participant-specific effects of the shoe LBS according
8
9 4 to the task characteristics,¹⁸ the participants' characteristics^{12,20} and biomechanics.^{13,21,22}
10
11
12 5 In the current study, correlation analyses were used on individual variations induced by an
13
14 6 increased shoe LBS to highlight such potential associations. It can be noted that in our sample
15
16 7 of participants, an increased LBS mainly induced unclear (5 out of 11 runners) and
17
18 8 detrimental (4 out of 11 runners) metabolic effects. The correlations were thus interpreted in
19
20 9 terms of « less detrimental metabolic effect » rather than « more beneficial metabolic effect »
21
22
23 10 induced by an increased shoe LBS.
24
25
26

27 11 At baseline of the prolonged run, the correlations showed that an increased shoe LBS
28
29 12 was less able to induce detrimental metabolic effect if this shoe intervention decreased (or did
30
31 13 not increase) the ground contact time, increased (or did not decrease) the body vertical
32
33 14 oscillation, or increased (or did not decrease) the TA activations relative to the runners'
34
35 15 habitual shoes with lower LBS. These results were surprising as an increased shoe LBS was
36
37 16 reported to induce longer ground contact time,¹⁹ and because the metabolic demand has been
38
39 17 shown to be inversely proportional to the ground contact time⁴² (regardless of any shoe
40
41
42 18 intervention). Willwacher et al.⁴³ suggested that an increased ground contact time with an
43
44 19 increased shoe LBS may reflect a weakness of strength capability of the ankle plantarflexors.
45
46 20 On this basis, the current runners might not be strong enough to benefit from such LBS
47
48 21 increase in their habitual shoes. On the other hand, the change of LBS may be functionally too
49
50 22 different from the habitual shoes of some runners to maintain their habitual biomechanical
51
52 23 pattern. As a result, unusual adjustments in the stride spatiotemporal parameters may have
53
54 24 occurred to meet the requirements of the experiment, while inducing a greater energetic cost.
55
56
57
58
59
60

1 The ground contact time is generally inversely associated with the body vertical
2 oscillation and the aerial time.⁴⁴ This may explain why the variations of the ground contact
3 time and the body vertical oscillations were simultaneously identified as explanatory variables
4 in the correlations. This may explain why the variation of TA activations was also an
5 explanatory variable. Indeed, with the TA muscle being mainly activated during the aerial
6 phase (Supplementary material E), the greater TA activations identified as less detrimental in
7 the correlations should be due to a consequence of longer activation durations during longer
8 aerial phases (due to shorter ground stance phases) rather than greater activation magnitudes.

9 At the last time interval of the prolonged run, the greater the participants' height, the
10 lower the detrimental metabolic effects induced by an increased shoe LBS. Why the
11 participants' height may potentiate the metabolic effects of an increased shoe LBS was
12 unknown, but some hypotheses are presented. Greater height may imply greater mass and/or
13 greater body segment lengths. Additional correlation analyses confirmed this for the mass
14 ($p=0.002$; $r=0.82$) and the shoe size used as surrogate indicator of foot length ($p=0.034$;
15 $r=0.64$), although both these variables did not directly correlate with the variation of the net
16 energetic cost induced by an increased shoe LBS in the current study. It has been shown that
17 the greater the participants' mass, the greater the metabolic benefits induced by an
18 intermediate increase of shoe LBS.¹² The taller runners, and also the heavier, were thus less
19 affected by an increased LBS. Greater shoe size may imply greater forefoot bone length,
20 likely resulting in a greater lever arm of the ground reaction force relative to MTP and ankle
21 joint centers (i.e., gearing effect) during push-off.⁴⁵ This gearing effect, amplified by the high
22 LBS shoes,⁴³ may favour lower excursion of muscle fibres and slower shortening velocity of
23 the *triceps surae* muscle.^{17,46} These muscle responses have been associated with lower

1
2
3 1 metabolic energetic cost,⁴⁷ especially in participants benefiting from an increased shoe LBS
4
5
6 2 when surrogate measures were used.^{13,22}
7
8
9

10 3 According to the instant of the prolonged run, different explanatory variables were
11
12 4 involved in the participant-specific metabolic effects induced by an increased shoe LBS. This
13
14 5 could be due to the observed variability in the time-courses of the physiological and
15
16 6 biomechanical variables between the participants. Similar variability has previously been
17
18 7 reported during prolonged running where spatiotemporal parameters can either increase,
19
20
21 8 decrease, or remain constant over time depending on the runner.^{48,49} Because there was no
22
23
24 9 unique time-course of the biomechanical pattern over time (with or without shoe
25
26 10 intervention), a given correlation observed at an initial time interval of the run may not remain
27
28 11 at a later time interval because the correlated variables did not evolve similarly for each
29
30
31 12 runner. Furthermore, an increased shoe LBS may affect runners differently in a fatigue state
32
33 13 than in a non-fatigue state. This induced even more variability in the time-courses of the
34
35 14 physiological and biomechanical variables between the participants. This assumption seemed
36
37 15 to be corroborated with the number of participants benefiting from an increased shoe LBS
38
39 16 throughout the prolonged run. This number of participants followed an inverted “U-shape”
40
41 17 where the greatest number of participants benefiting from an increased LBS (n=6) occurred at
42
43 18 the half of the prolonged running. Before and after, a lower number of participants benefited
44
45 19 from an increased shoe LBS (n=4).
46
47
48
49

50 20 This study had some limitations highlighted for readers. While the current protocol
51
52 21 included a prolonged running session close to the VAT, the running duration and/or intensity
53
54 22 were not important enough to induce an increase in the net energetic cost (i.e. inducing
55
56 23 fatigue) in any of the shoe LBS conditions. The habitual shoe of the runners have been used,
57
58 24 which may differ in several features not measured, such as midsole foam viscoelasticity, sole
59
60

1 geometry, or shoe mileage. Thus, the beneficial (or detrimental) effects of an increased LBS
2 for some runners may have been potentiated by the appropriate (or not) interaction with these
3 other shoe features.⁵⁰ The difference in mass between both shoe LBS conditions (20 g on
4 average) may have slightly influenced the metabolic results by about 0.2% based on the
5 relationship between shoe mass and metabolic demand.⁵¹ A point of awareness was that the
6 results came from two different prolonged running sessions performed in two different days
7 with low or high LBS shoes. In order to offset the results obtained from two different days,
8 physiological data were normalised by the day-specific respiratory reference values of the
9 participants recorded during a sitting rest before each running session. Nevertheless, it cannot
10 be ensured that the current results, while reflecting individual variations in responses to LBS
11 interventions, were not influenced by other unwanted effects. Furthermore, EMG results may
12 have been influenced by EMG placement between sessions thus it cannot be definitively
13 ensured that the sensors were exactly placed in the same location between both sessions.
14 Although firmly secured with medical straps, the sensors' abilities to record muscle
15 activations could have been affected by the accumulation of sweat. This study had a small
16 sample size resulting in low statistical power regarding the net energetic cost of prolonged
17 running ($1-\beta=0.28$ for the LBS main effect; $1-\beta=0.40$ for the interaction effect). A *post hoc*
18 power analysis (G*Power v3.1.9.4) showed that, for achieving at least 0.80 power, 45 and 24
19 participants would have been needed to rule-out the Type II error possibilities for the LBS
20 main effect and the interaction effect, respectively. In correlation analyses, considering large
21 correlations ($r<-0.5$ or $r>0.5$) on 11 participants with an alpha level set to 0.1 resulted in 0.50
22 statistical power (while 23 participants would have been needed to achieve at least 0.80
23 power).³⁵ Therefore, findings from this study should be considered with caution. We would
24 suggest the future studies to include the explanatory variables highlighted in the current study
25 (i.e. ground contact time and height). This would enable to investigate shoe LBS effects with

1
2
3 1 more powerful experimental designs on the explanatory variables highlighted in the current
4
5 2 study. As it is, the current findings are suitable for male novice/recreational runners. As the
6
7
8 3 height (in this study) and the mass¹² are explanatory variables of the metabolic demand, it is
9
10 4 hypothesised that female runners may require a shoe LBS tuning different from males due to
11
12
13 5 their morphological specificities.
14
15
16 6

17
18
19 7 **Conclusion**
20
21

22 8 To the question: “is it an appropriate choice to increase the longitudinal bending stiffness in
23
24 9 habitual shoes of runners for improving performance?”, the answer from this study would be:
25
26 10 “it depends”. At a mean group level, an increased shoe LBS did not limit and/or delay the
27
28
29 11 increase of the net metabolic energetic cost during a prolonged running. However, participant-
30
31 12 specific metabolic (beneficial or detrimental) effects induced by an increased LBS have been
32
33
34 13 highlighted. These participant-specific metabolic effects were associated with the variation of
35
36 14 the ground contact time at the beginning and the height of the participants at the end of the
37
38 15 prolonged run. The results of this study increased the body of knowledge concerning the
39
40 16 potential explanatory variables associated with specific metabolic effects of shoe LBS.
41
42
43 17 However, future studies are still needed to be able to recommend, with more certainty, the
44
45 18 appropriate increase of shoe LBS to a given runner who needs it, based on his individual
46
47 19 characteristics and/or biomechanics. Indeed, the metabolic effects of an increased LBS were
48
49 20 more detrimental than the use of habitual shoes with lower LBS for most of the participants in
50
51 21 the current study. This study thus suggested that a shoe LBS intervention was likely not
52
53
54 22 beneficial for performance purposes if the runners’ shoe habits were too much disrupted.
55
56

57 23 **Acknowledgements**
58
59
60

Authors would thank Romain Hardouin for his help during the configuration of EMG record and during EMG data export. Authors also thank all the participants for taking part to these three sessions of running. Finally, authors thank Dr. Eleni Grimpampi for her English review of the manuscript.

Declaration of interest statement

NF and ND are members of the company Decathlon™ in the research department. NF, GR, EB, and ND declared that the results of the study are presented clearly, honestly, and without fabrication, falsification, nor inappropriate data manipulation.

References

1. Stefanyshyn DJ, Nigg BM. Influence of midsole bending stiffness on joint energy and jump height performance. *Med Sci Sports Exerc* 2000; 32: 471–476.
2. Hoogkamer W, Kipp S, Frank JH, et al. A Comparison of the Energetic Cost of Running in Marathon Racing Shoes. *Sport Med* 2018; 48: 1009–1019.
3. Barnes KR, Kilding AE. A Randomized Crossover Study Investigating the Running Economy of Highly-Trained Male and Female Distance Runners in Marathon Racing Shoes versus Track Spikes. *Sport Med* 2019; 49: 331–342.
4. Hunter I, McLeod A, Valentine D, et al. Running economy, mechanics, and marathon racing shoes. *J Sports Sci* 2019; 37: 2367–2373.
5. Hébert-Losier K, Finlayson SJ, Driller MW, et al. Metabolic and performance responses of male runners wearing 3 types of footwear: Nike Vaporfly 4%, Saucony

- 1
2
3 1 Endorphin racing flats, and their own shoes. *J Sport Heal Sci*. Epub ahead of print
4
5 2 2020. DOI: 10.1016/j.jshs.2020.11.012.
6
7
8 3 6. Quealy K, Katz J. Nike Says Its \$250 Running Shoes Will Make You Run Much
9
10 4 Faster. What if That's Actually True? - The New York Times,
11
12
13 5 [https://www.nytimes.com/interactive/2018/07/18/upshot/nike-vaporfly-shoe-](https://www.nytimes.com/interactive/2018/07/18/upshot/nike-vaporfly-shoe-strava.html)
14
15 6 [strava.html](https://www.nytimes.com/interactive/2018/07/18/upshot/nike-vaporfly-shoe-strava.html) (2018, accessed 16 December 2020).
16
17
18 7 7. Bermon S, Garrandes F, Szabo A, et al. Effect of Advanced Shoe Technology on the
19
20 8 Evolution of Road Race Times in Male and Female Elite Runners. *Front Sport Act*
21
22 9 *Living* 2021; 3: 653173.
23
24
25 10 8. Burns GT, Tam N. Is it the shoes? A simple proposal for regulating footwear in road
26
27 11 running. *British Journal of Sports Medicine* 2020; 54: 439–441.
28
29
30 12 9. Frederick EC. No evidence of a performance advantage attributable to midsole
31
32 13 thickness. *Footwear Science* 2020; 12: 1–2.
33
34
35 14 10. Hoogkamer W. More isn't always better. *Footwear Science* 2020; 12: 75–77.
36
37
38 15 11. Nigg BM, Cigoja S, Nigg SR. Teeter-totter effect: A new mechanism to understand
39
40 16 shoe-related improvements in long-distance running. *British Journal of Sports*
41
42 17 *Medicine*. Epub ahead of print 7 August 2020. DOI: 10.1136/bjsports-2020-102550.
43
44
45 18 12. Roy J-PR, Stefanyshyn DJ. Shoe Midsole Longitudinal Bending Stiffness and Running
46
47 19 Economy, Joint Energy, and EMG. *Med Sci Sport Exerc* 2006; 38: 562–569.
48
49
50 20 13. Madden R, Sakaguchi M, Tomaras EK, et al. Forefoot bending stiffness, running
51
52 21 economy and kinematics during overground running. *Footwear Sci* 2016; 8: 91–98.
53
54
55 22 14. Flores N, Delattre N, Berton E, et al. Does an increase in energy return and/or
56
57 23 longitudinal bending stiffness shoe features reduce the energetic cost of running? *Eur J*
58
59
60

- 1
2
3 1 *Appl Physiol* 2019; 119: 429–439.
4
5
6 2 15. Healey L, Hoogkamer W. Longitudinal bending stiffness does not affect running
7
8 3 economy in Nike Vaporfly shoes. *SportRxiv*. Epub ahead of print 2021. DOI:
9
10 4 10.31236/OSF.IO/37UZR.
11
12
13 5 16. Beck ON, Golyski PR, Sawicki GS. Adding carbon fiber to shoe soles may not
14
15 6 improve running economy: a muscle-level explanation. *Sci Rep*; 10. Epub ahead of
16
17 7 print 1 December 2020. DOI: 10.1038/s41598-020-74097-7.
18
19
20
21 8 17. Cigoja S, Fletcher JR, Esposito M, et al. Increasing the midsole bending stiffness of
22
23 9 shoes alters gastrocnemius medialis muscle function during running. *Sci Rep* 2021; 11:
24
25 10 749.
26
27
28 11 18. Day E, Hahn M. Optimal footwear longitudinal bending stiffness to improve running
29
30 12 economy is speed dependent. *Footwear Sci* 2019; 1–11.
31
32
33 13 19. Ortega JA, Healey LA, Swinnen W, et al. Energetics and Biomechanics of Running
34
35 14 Footwear with Increased Longitudinal Bending Stiffness: A Narrative Review. *Sport*
36
37 15 *Med* 2021; 51: 873–894.
38
39
40
41 16 20. Oh K, Park S. The bending stiffness of shoes is beneficial to running energetics if it
42
43 17 does not disturb the natural MTP joint flexion. *J Biomech*. Epub ahead of print 18
44
45 18 January 2017. DOI: 10.1016/j.jbiomech.2017.01.014.
46
47
48 19 21. McLeod AR, Bruening D, Johnson AW, et al. Improving running economy through
49
50 20 altered shoe bending stiffness across speeds. *Footwear Sci*. Epub ahead of print 2020.
51
52 21 DOI: 10.1080/19424280.2020.1734870.
53
54
55
56 22 22. Flores N, Le Gendre E, Rao G, et al. Relationships between biomechanics,
57
58 23 anthropometrics, and running economy as a function of shoe mechanical feature
59
60

- 1
2
3 1 variations. *Footwear Sci* 2019; 11: S48–S49.
4
5
6 2 23. Cigoja S, Firminger CR, Asmussen MJ, et al. Does increased midsole bending stiffness
7
8 3 of sport shoes redistribute lower limb joint work during running? *J Sci Med Sport*
9
10 4 2019; 22: 1272–1277.
11
12
13 5 24. Čigoja S, Fletcher JR, Nigg BM. Can changes in midsole bending stiffness of shoes
14
15 6 affect the onset of joint work redistribution during a prolonged run? *J Sport Heal Sci*.
16
17 7 Epub ahead of print 9 January 2021. DOI: 10.1016/j.jshs.2020.12.007.
18
19
20
21 8 25. Sanno M, Willwacher S, Epro G, et al. Positive Work Contribution Shifts from Distal
22
23 9 to Proximal Joints during a Prolonged Run. *Med Sci Sport Exerc* 2018; 50: 2507–2517.
24
25
26 10 26. Hunter JG, Smith AMB, Sciarratta LM, et al. Standardized lab shoes do not decrease
27
28 11 loading rate variability in recreational runners. *J Appl Biomech* 2020; 36: 340–344.
29
30
31 12 27. Worobets J, Wannop JW. Influence of basketball shoe mass, outsole traction, and
32
33 13 forefoot bending stiffness on three athletic movements. *Sport Biomech* 2015; 14: 351–
34
35 14 360.
36
37
38
39 15 28. Mezzani A, Agostoni P, Cohen-Solal A, et al. Standards for the use of cardiopulmonary
40
41 16 exercise testing for the functional evaluation of cardiac patients: a report from the
42
43 17 Exercise Physiology Section of the European Association for Cardiovascular
44
45 18 Prevention and Rehabilitation. *Eur J Cardiovasc Prev Rehabil* 2009; 16: 249–267.
46
47
48
49 19 29. Adams D, Pozzi F, Carroll A, et al. Validity and Reliability of a Commercial Fitness
50
51 20 Watch for Measuring Running Dynamics. *J Orthop Sport Phys Ther* 2016; 46: 471–
52
53 21 476.
54
55
56 22 30. Péronnet F, Massicotte D. Table of nonprotein respiratory quotient: an update. *Can J*
57
58 23 *Sport Sci* 1991; 16: 23–29.
59
60

- 1
2
3 1 31. Enoka RM. *Neuromechanics of human movement*. 4th edn. Human Kinetics,
4
5 2 Champaign, 2008.
6
7
8 3 32. Morin J-B, Samozino P, Millet GY. Changes in Running Kinematics, Kinetics, and
9
10 4 Spring-Mass Behavior over a 24-h Run. *Med Sci Sport Exerc* 2011; 43: 829–836.
11
12
13 5 33. Pataky TC, Vanrenterghem J, Robinson MA. Zero- vs. one-dimensional, parametric vs.
14
15 6 non-parametric, and confidence interval vs. hypothesis testing procedures in one-
16
17 7 dimensional biomechanical trajectory analysis. *J Biomech* 2015; 48: 1277–1285.
18
19
20
21 8 34. Lakens D. Calculating and reporting effect sizes to facilitate cumulative science: A
22
23 9 practical primer for t-tests and ANOVAs. *Front Psychol*; 4. Epub ahead of print 2013.
24
25 10 DOI: 10.3389/fpsyg.2013.00863.
26
27
28 11 35. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. NY: Routle. New-
29
30 12 York, 1988.
31
32
33 13 36. Fletcher JR, Esau SP, MacIntosh BR. Economy of running: beyond the measurement of
34
35 14 oxygen uptake. *J Appl Physiol* 2009; 107: 1918–1922.
36
37
38 15 37. Barnes KR, Kilding AE. Running economy: measurement, norms, and determining
39
40 16 factors. *Sport Med - open* 2015; 1: 8.
41
42
43 17 38. Delattre N, Chambon N, Berton E, et al. Effect of time during a running session with
44
45 18 minimal footwear. *Comput Methods Biomech Biomed Engin* 2013; 16: 104–105.
46
47
48 19 39. Lindorfer J, Kröll J, Schwameder H. Familiarisation of novice and experienced
49
50 20 treadmill users during a running session: Group specific evidence, time and individual
51
52 21 patterns. *Hum Mov Sci* 2020; 69: 102530.
53
54
55 22 40. Mohr M, von Tschanner V, Nigg S, et al. Systematic reduction of leg muscle activity
56
57 23 throughout a standard assessment of running footwear. *J Sport Heal Sci*. Epub ahead of
58
59
60

- print 13 January 2021. DOI: 10.1016/j.jshs.2021.01.003.
41. Divert C, Baur H, Mornieux G, et al. Stiffness adaptations in shod running. *J Appl Biomech* 2005; 21: 311–21.
42. Kram R, Taylor CR. Energetics of running: a new perspective. *Nature* 1990; 346: 265–267.
43. Willwacher S, König M, Braunstein B, et al. The gearing function of running shoe longitudinal bending stiffness. *Gait Posture* 2014; 40: 386–390.
44. Lussiana T, Patoz A, Gindre C, et al. The implications of time on the ground on running economy: Less is not always better. *J Exp Biol*; 222. Epub ahead of print 15 March 2019. DOI: 10.1242/jeb.192047.
45. Ueno H, Suga T, Takao K, et al. Association between Forefoot Bone Length and Performance in Male Endurance Runners. *Int J Sports Med* 2018; 39: 275–281.
46. Carrier DR, Heglund NC, Earls KD. Variable gearing during locomotion in the human musculoskeletal system. *Science* 1994; 265: 651–3.
47. Fletcher JR, Groves EM, Pfister TR, et al. Can muscle shortening alone, explain the energy cost of muscle contraction in vivo? *Eur J Appl Physiol* 2013; 113: 2313–2322.
48. Dutto DJ, Smith GA. Changes in spring-mass characteristics during treadmill running to exhaustion. *Med Sci Sports Exerc* 2002; 34: 1324–31.
49. Hunter I, Smith GA. Preferred and optimal stride frequency, stiffness and economy: changes with fatigue during a 1-h high-intensity run. *Eur J Appl Physiol* 2007; 100: 653–661.
50. Nigg BM, Čigoja S, Nigg SR. Effects of running shoe construction on performance in long distance running. *Footwear Sci* 2020; 12: 133–138.

- 1
2
3 1 51. Franz JR, Wierzbinski CM, Kram R. Metabolic Cost of Running Barefoot versus Shod.
4
5 2 *Med Sci Sport Exerc* 2012; 44: 1519–1525.
6
7
8
9 3
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure captions

Figure 1. Time-course of the net metabolic energetic cost during the prolonged running duration with low (left) and high (right) LBS shoes. Participants with long dotted lines had lower energetic cost during the prolonged run with the low LBS shoes, those with full lines had lower energetic cost during the prolonged run with the high LBS shoes, and those with short dotted lines had randomly lower energetic cost during the prolonged run either with low or high LBS shoes (see Table 2 for corresponding data).

Figure 2. Time-course of the activations of TA, GM, and VL muscles during the prolonged running duration with low (left) and high (right) LBS shoes. Data of GM activations is missing for participants 3 and 10. * indicated a significant main effect of the running duration.

Figure 3. Time-courses of the stride cadence, the body vertical oscillation, the ground contact time, the leg stiffness, and the vertical stiffness during the prolonged running duration with low (left) and high (right) LBS shoes. Data is missing for participants 1 and 4.

Figure 4. Correlations between the individual variations of the net energetic cost induced by an increased LBS in participants' habitual shoes and the individual variations of the ground contact time (top left), the body vertical oscillation (top right), the TA activations (bottom left) induced by an increased LBS, and the participants' height (bottom right). Dark and grey points respectively indicated the participants benefiting from the high LBS and low LBS condition, while white points indicated unclear participants (based on Table 2). Data is missing for participants 1 and 4 for spatiotemporal variables.

Figure captions

Figure 1. Time-course of the net metabolic energetic cost during the prolonged running duration with low (left) and high (right) LBS shoes. Participants with long dotted lines had lower energetic cost during the prolonged run with the low LBS shoes, those with full lines had lower energetic cost during the prolonged run with the high LBS shoes, and those with short dotted lines had randomly lower energetic cost during the prolonged run either with low or high LBS shoes (see Table 2 for corresponding data).

Figure 2. Time-course of the activations of TA, GM, and VL muscles during the prolonged running duration with low (left) and high (right) LBS shoes. Data of GM activations is missing for participants 3 and 10. * indicated a significant main effect of the running duration.

Figure 3. Time-courses of the stride cadence, the body vertical oscillation, the ground contact time, the leg stiffness, and the vertical stiffness during the prolonged running duration with low (left) and high (right) LBS shoes. Data is missing for participants 1 and 4.

Figure 4. Correlations between the individual variations of the net energetic cost induced by an increased LBS in participants’ habitual shoes and the individual variations of the ground contact time (top left), the body vertical oscillation (top right), the TA activations (bottom left) induced by an increased LBS, and the participants’ height (bottom right). Dark and grey points respectively indicated the participants benefiting from the high LBS and low LBS condition, while white points indicated unclear participants (based on Table 2). Data is missing for participants 1 and 4 for spatiotemporal variables.

Table 1. Characteristics of the participants and participants' own running shoes (see supplementary material A for pictures of the shoes).

Participant	Height (cm)	Mass (kg)	Speed at 95% of VAT (m/s)	Shoe brand and version	Shoe size (EU size)	Shoe mass (g)	Low LBS (Nm/deg)	High LBS (Nm/deg)
1	172.1	63.1	3.19	Nike Lunar Launch	44	246	0.070	0.233 (+231%)
2	177.2	75.6	2.92	Mizuno Wave Inspire 9	44	312	0.114	0.306 (+168%)
3	176.0	74.2	3.19	Nike Air Max	45	325	0.043	0.156 (+260%)
4	182.1	82.3	3.33	Nike Free Run 2	44	343	0.027	0.148 (+454%)
5	175.3	67.8	3.19	Asics Gel Innovate	44	286	0.057	0.206 (+263%)
6	177.0	68.3	3.19	Adidas Tubular	43	268	0.038	0.124 (+225%)
7	170.2	61.0	2.92	Kalenji Active Trail	41	257	0.099	0.398 (+304%)
8	180.4	78.4	3.19	Adidas Boost Supernova 9	44	338	0.067	0.162 (+143%)
9	168.2	58.2	2.78	Brooks Aduro 4	42	278	0.083	0.287 (+246%)
10	176.0	86.8	2.92	Adidas Terrex	45	352	0.047	0.149 (+220%)
11	170.0	64.9	3.33	Mizuno Wave Elixir 7	42.5	257	0.041	0.196 (+377%)
Mean	175.0	71.0	3.11		43.5	297	0.062	0.215 (+263%)
SD	4.4	9.2	0.19		1.5	39	0.027	0.084 (89%)

Table 2. Individual (absolute and relative) variations of the net energetic cost of running induced by an increased LBS at each time interval of the prolonged running. Negative (dark grey cell) or positive (light grey cell) values respectively indicated that an increased LBS in the participant's habitual shoes induced decreased or increased net energetic cost of running compared to their lower habitual LBS. A beneficial shoe condition was unclear if negative and positive values were combined for a given participant.

	Δ net energetic cost (W/kg)							
Participants	From 9 to 10 min	From 14 to 15 min	From 19 to 20 min	From 24 to 25 min	From 29 to 30 min	From 34 to 35 min	From 39 to 40 min	Beneficial shoe condition
1	0.16 1.1%	-0.98 -6.5%	-0.80 -5.2%	-0.94 -6.1%	-0.45 -3.0%	-0.23 -1.5%	-0.15 -0.9%	Unclear
2	-1.94 -14.4%	-0.74 -5.8%	-1.17 -9.0%	-1.15 -8.7%	-1.07 -8.2%	-0.71 -5.4%	-0.69 -5.4%	High LBS
3	0.76 5.4%	0.24 1.7%	-0.43 -2.9%	0.19 1.3%	-0.24 -1.7%	0.25 1.7%	0.19 1.2%	Unclear
4	-0.60 -4.3%	-1.03 -7.4%	-1.62 -11.3%	-1.78 -12.2%	-3.18 -21.9%	-3.13 -20.9%	-3.95 -26.2%	High LBS
5	-0.28 -1.8%	0.19 1.2%	-0.47 -2.8%	-0.02 -0.2%	-0.26 -1.6%	0.56 3.6%	0.96 6.5%	Unclear
6	1.61 11.1%	1.29 8.8%	0.27 1.7%	-0.20 -1.3%	0.09 0.6%	-0.19 -1.3%	-0.11 -0.8%	Unclear
7	-1.31 -8.5%	-0.68 -4.4%	-0.25 -1.6%	-0.06 -0.4%	-0.47 -2.9%	0.50 3.5%	1.34 9.4%	Unclear
8	0.82 6.3%	1.06 8.4%	1.68 13.1%	0.49 3.8%	1.33 10.9%	0.38 3.0%	0.46 3.8%	Low LBS
9	3.11 30.3%	3.08 29.1%	3.11 28.9%	2.51 22.6%	1.85 16.2%	1.46 12.7%	1.25 10.3%	Low LBS
10	0.89 8.3%	1.17 10.8%	0.84 7.7%	1.18 10.8%	1.47 13.5%	1.54 14.1%	1.58 14.6%	Low LBS
11	1.02 7.1%	1.67 11.7%	1.74 12.5%	1.86 13.1%	1.99 13.9%	1.24 8.7%	1.74 12.1%	Low LBS
Number of participants benefiting from low LBS (Δ net energetic cost>0)	7	7	5	5	5	7	7	
Number of participants benefiting from high LBS (Δ net energetic cost<0)	4	4	6	6	6	4	4	

Supplementary material A

Figure A. Pictures of the participant’s habitual running shoes. The order of the shoes corresponds to the order of participants in each Table or Figure of the main document. The last picture is the stiff carbon plate used in this study.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supplementary material B

Procedure of the mechanical bending test

A shoe last was cut at the metatarsophalangeal joint level (70% of the shoe last length) to remove the anterior part. This shoe last was inserted inside the shoe. The shoe was then secure upside down on a horizontal plate from its posterior part with a clamp. The anterior ending of the shoe last inside the shoe (*B*) was identified and marked on the external part of the shoe (Figure B), and then marked on a graph paper perpendicularly located to the bending axis of the shoe forefoot. A horizontal straight line was drawn through *B* on the graph paper. A hook was then set on the shoe outsole at the anterior ending of the shoe. Either no mass ($m=0$ kg), or m masses of 1 kg, 2 kg, and 3 kg were hung along the hook. The location of the hook (*H*) was also marked on the graph paper (Figure B). A vertical straight line was drawn through *H* on the graph paper, which cut perpendicularly the horizontal straight line in a point *O* (Figure B). The length of $[BO]_m$, in meters, was the lever arm of the force applied by each m mass relative to the bending axis. The moment (M_m , in Nm, equation B1) applied to the anterior part of the shoe was thus computed as:

$$M_m = m \times g \times [BO]_m \text{ (B1)}$$

g being the gravity constant value equal to 9.81 m/s² and m being the mass equal to 0 kg, 1 kg, 2 kg, and 3 kg.

The length of $[OH]_m$, in meters, was the vertical displacement of the anterior part of the shoe due to the applied moment of force (for $m=0$ kg, the length $[OH]_0$ was the height of the curvature of the anterior part of the shoe, i.e. the toe spring). The bending angle ($\hat{B}_m$, in deg, equation B2) of the shoe was computed as:

$$\hat{B}_m = \tan^{-1} \left(\frac{[OH]_m}{[BO]_m} \right) \text{ (B2)}$$

Finally, the bending stiffness value (in Nm/deg) was obtained through the slope of a linear regression fitting the four force moments (M_m) and bending angles ($\hat{B}_m$) plotted on a moment-angle graph.

Figure B. Schema illustrating the identification of the point of the anterior ending of the shoe last inside the shoe (or bending axis, B), the location of the hook after a mass was hung (H), and the lever arm of the force applied by the mass relative to the bending axis (O) for the low (grey) and high (dark) longitudinal bending stiffness shoe conditions.

Supplementary material C

Figure C. Picture of the experimental set-up with a participant running on the loop with hard concrete surface (top), and illustration of the loop set-up with its dimensions and the direction of running (bottom). The total distance of the loop was about 230 metres. Each straight line measured 80 metres. The ground marks in both extremities of the loop were located at around 18 metres from the previous ones, thus allowing the participants to turn behind the ground mark in a secure way thanks to a slight deceleration.

Supplementary material D

Table D1. List of variables including in the correlation analyses. ^V indicates the variables presented in the main text for the correlations performed at the first and the last time intervals.

Variable	Description	Correlation analyses	
Δ Net energetic cost (W/kg) ^v	Variation of the net energetic cost of running by subtracting values induced by the high LBS to those induced by the low LBS from 9 to 10 min, from 14 to 15 min, from 19 to 20 min, from 24 to 25 min, from 29 to 30 min, from 34 to 35 min, and from 39 to 40 min.	These variables were used in the correlation analyses only for identical time intervals than those of the net energetic cost of running For example, Δ Cadence from 14 to 15 min was used in the correlation analyses as a potential explanatory variable for Δ Net energetic cost from 14 to 15 min.	
Δ TA (a.u.) ^v	Variation of the lower limb muscle activations by subtracting values induced by the high LBS to those induced by the low LBS from 9 to 10 min, from 14 to 15 min, from 19 to 20 min, from 24 to 25 min, from 29 to 30 min, from 34 to 35 min, and from 39 to 40 min.		
Δ GM (a.u.) ^v			
Δ VL (a.u.) ^v			
Δ Cadence (stride/min) ^v			
Δ Vertical oscillation (cm) ^v			
Δ Ground contact time (ms) ^v			
Δ Vertical stiffness (kN/m)	Variation of spatiotemporal parameters of running by subtracting values induced by the high LBS to those induced by the low LBS from 9 to 10 min, from 14 to 15 min, from 19 to 20 min, from 24 to 25 min, from 29 to 30 min, from 34 to 35 min, and from 39 to 40 min.		
Δ Leg stiffness (kN/m)			
Mass (kg) ^v	Mass of the participant	These variables were used in the correlation analyses for all time intervals of the variation of the net energetic cost of running For example, height was used in the correlation analyses as a potential explanatory variable for Δ Net energetic cost in each of the seven time intervals.	
Height (cm) ^v	Height of the participant		
Speed (m/s) ^v	Imposed speed to the participant equal to 95% of the VAT speed		
Shoe size (EU size)	Size of the participant's own shoe		
Low LBS (Nm/deg)	LBS of the participant's own shoe		
Δ LBS ^v	Variation of LBS when adding a stiff plate		
Shoe mass (g)	Mass of the participant's own shoe		
Habitual cadence (stride/min)	Spatiotemporal parameter of running measured at baseline (from 9 to 10 min) with the participant's own shoe (low LBS) considered as the runners' habitual running technique		
Habitual vertical oscillation (cm)			
Habitual ground contact time (ms)			
Habitual vertical stiffness (kN/m)			
Habitual leg stiffness (kN/m)			

Table D2. Explanatory variables correlated with the variation of the net energetic cost of running induced by an increased shoe LBS according to the time interval of the prolonged running. Based on criteria set in the statistical section, only the significant correlated variables were presented while NS indicated non-significant correlation.

Explanatory variables	Δ net energetic cost						
	From 9 to 10 min	From 14 to 15 min	From 19 to 20 min	From 24 to 25 min	From 29 to 30 min	From 34 to 35 min	From 39 to 40 min
Δ ground contact time from 9 to 10 min	$p=0.067$ $r=0.63$	NS	NS	NS	NS	NS	NS
Δ vertical oscillation from 9 to 10 min	$p=0.070$ $r=-0.63$	NS	NS	NS	NS	NS	NS
Δ TA activations from 9 to 10 min and from 14 to 15 min	$p=0.094$ $r=-0.53$	$p=0.082$ $r=-0.55$	NS	NS	NS	NS	NS
Δ VL activations from 24 to 25 min	NS	NS	NS	$p=0.049$ $r=-0.60$	NS	NS	NS
Height	NS	NS	NS	$p=0.043$ $r=-0.62$	$p=0.094$ $r=-0.53$	$p=0.026$ $r=-0.66$	$p=0.019$ $r=-0.69$
Other variables at any time intervals	NS	NS	NS	NS	NS	NS	NS

Supplementary material E

Figure E. Typical VL (yellow), GM (red), and TA (blue) raw activations during a running cycle. Vertical dotted lines represented estimations of instants of heel strike (HS) and toe-off (TO).