



HAL
open science

Detection of *Rickettsia raoultii* in *Dermacentor reticulatus* and *Haemaphysalis inermis* ticks in Slovakia

Basma Ouarti, Basma El Hamzaoui, Michal Stanko, Maureen Laroche, Oleg Mediannikov, Philippe Parola, Zuzana Sekeyová

► **To cite this version:**

Basma Ouarti, Basma El Hamzaoui, Michal Stanko, Maureen Laroche, Oleg Mediannikov, et al.. Detection of *Rickettsia raoultii* in *Dermacentor reticulatus* and *Haemaphysalis inermis* ticks in Slovakia. *Biologia*, 2022, 77 (6), pp.1611-1617. 10.1007/s11756-021-00789-8 . hal-03334365

HAL Id: hal-03334365

<https://hal-amu.archives-ouvertes.fr/hal-03334365>

Submitted on 6 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License



Detection of *Rickettsia raoultii* in *Dermacentor reticulatus* and *Haemaphysalis inermis* ticks in Slovakia

Basma Ouarti^{1,2} · Basma El Hamzaoui^{1,2} · Michal Stanko³ · Maureen Laroche^{1,2} · Oleg Mediannikov^{1,2} · Philippe Parola^{1,2} · Zuzana Sekeyová⁴

Received: 8 March 2021 / Accepted: 14 May 2021
© The Author(s) 2021

Abstract

Ticks are vector arthropods responsible for the transmission of several pathogenic agents that affect both human and animal health worldwide. In this study our objective was to analyse, using molecular tools, the bacterial community of *Dermacentor reticulatus* and *Haemaphysalis inermis* ticks collected in south-eastern Slovakia. Using real-time PCR, we identified the presence of *Rickettsia* spp. DNA at levels of 14/59 (23.72 %) and 29/173 (16.76 %) in *D. reticulatus* and *H. inermis*, respectively. In addition, using standard PCR and sequencing, we identified the presence of *Rickettsia raoultii* DNA in 13 ticks belonging to the two investigated species. *Rickettsia raoultii* blast results revealed an average identification percentage of 99.62 %. Following the results of this molecular study there is a possibility that *D. reticulatus* and *H. inermis* play a potential role in the transmission of *R. raoultii*. To prove the possibility of validity of this hypothesis, we suggest performing experimental models in future studies. Our results can serve as preliminary data for future transmission models.

Keywords Ticks · *Dermacentor reticulatus* · *Haemaphysalis inermis* · Slovakia

Introduction

Bacteria belonging to the genus *Rickettsia* are known as causative agents of various vector-borne zoonotic diseases. They are responsible for mild to severe diseases in humans (Parola et al. 2013; Eldin and Parola 2018).

The diversity of pathogens in ecosystems also depends on the species diversity of vectors and hosts and the diversity of

habitats (de la Fuente et al. 2008). Many ticks in Europe, including *Dermacentor reticulatus* Fabricius, 1794 are known to act as carriers, reservoirs and/or vectors of different pathogenic *Rickettsia* species e.g., *Rickettsia slovaca* (Sekeyová et al. 1998; Garcia-Vozmediano et al. 2020) and *Rickettsia raoultii* (Boldiš et al. 2008; Földvári et al. 2013).

Revising the role of another tick species, *Haemaphysalis inermis* Birula, 1895 in the spread of rickettsiae continues to be challenging as few data have been previously published (Portillo et al. 2008; Špitalská et al. 2018). In a study carried out in Hungary, the authors detected the presence of *R. helvetica* in *H. inermis* and suggested that this tick species could be a potential vector of this pathogen (Hornok et al. 2010).

Recent epidemiological studies of *Rickettsia* in Slovakia were completed in suburban, natural and rural habitats (Minichová et al. 2017), or in an urban temperate forest (Chvostáč et al. 2018) in the western part of the country. A dry forest-steppe biotope is rarely referenced in literature except, in the past century, by Řeháček et al. (1976a, b). The authors indicated that: “The biotopes of the spotted fever group rickettsiae occurring in east Slovakia—namely, cultivated and meadow steppes with sparse forests—share some of the characteristics of the biotopes of *Rickettsia sibirica*. Thus, besides the new species of spotted fever group rickettsiae,

Highlights

- *Dermacentor reticulatus* and *Haemaphysalis inermis* ticks collected in south-eastern Slovakia are infected by *Rickettsia raoultii*.
- These preliminary data could be used in a future transmission model.

✉ Philippe Parola
philippe.parola@univ-amu.fr

¹ Aix Marseille Univ, IRD, AP-HM, SSA, VITROME, Marseille, France

² IHU-Méditerranée Infection, Marseille, France

³ Institute of Parasitology, Slovak Academy of Sciences, Hlinkova 3, 040 01 Košice, Slovak Republic

⁴ Biomedical Research Centre, Institute of Virology, Slovak Academy of Sciences, Dúbravská cesta 9, 845 05 Bratislava, Slovak Republic

the circulation of *R. sibirica* and perhaps of *Rickettsia conorii* in east Slovakia should not be excluded". However, none of them were ever confirmed in south-eastern Slovakia afterwards (Řeháček et al. 1976a, b).

Our objective was to investigate the possibility of detecting rickettsiae in *D. reticulatus* and *H. inermis*, two relatively abundant tick species in the steppe habitats of south-eastern Slovakia.

Materials and methods

Tick collection and sampling site

The tick collection was carried out on October 23, 2015 by flagging grass and shrubs using a white cotton flag passed over the vegetation at the level of three microhabitats (forest, ecotone and meadow) along a line of 100 m in the vegetation of the protected zone of Slovak Karst National Park in south-eastern Slovakia (Central Europe) near the village of Hrhov (200–220 m above sea level, 48°34.899 N, 20°46.743 E) (Fig. 1).

DNA extraction and molecular detection of bacteria in ticks using real time PCR

Half of the tick body without legs was selected for DNA extraction using EZ1 DNA tissue kit (Qiagen, Hilden, Germany) following a protocol previously elaborated in our laboratory (Diarra et al. 2017).

After extraction, the DNA of each sample was screened with the CFX96 real-time PCR (Bio -Rad, Marnes-la-Coquette, France) and the LightCyclerR 480 Probes Master

Mix (Roche Diagnostics, Indianapolis, USA) for the presence of the following bacterial microorganisms (*Rickettsia* spp., *Bartonella* spp., *Borrelia* spp., *Coxiella burnetii* and *Anaplasma* spp.) using specific primers and probes listed in Diarra et al. (2020) and Ouarti et al. (2020).

The qPCR reaction mixture for each plate has been used according to the manufacturer's protocol. Negative controls were used in each qPCR and consisted of DNA extracted from uninfected *Rhipicephalus sanguineus* ticks from the laboratory colony of IHU - VITROME (Marseille, France). Positive controls included DNA extracted from a dilution of cultured strains of *Borrelia crocidurae*, *Bartonella henselae*, *Rickettsia montanensis*, *Ehrlichia canis* and *Coxiella burnetii* (Socolovschi et al. 2012b; Aouadi et al. 2017). Results were deemed positive if the cycle threshold (Ct) value obtained by CFX96 was lower than ≤ 35 (Ouarti et al. 2020).

Bacterial species identification using sequencing

All ticks tested positive for *Anaplasma* spp., *Borrelia* spp. and *Rickettsia* spp. in qPCR were subjected to amplification using standard PCR prior to sequencing for identification of the bacterial species (Dahmani et al. 2015b; Diarra et al. 2017). In standard PCR the primers used to amplify the *ompA* gene were (f-70 and r-701). For the sequencing we used the following primers (f-70, f-180 and r-701) which amplify (629 to 632 bp) of protein A (Table 1) (Socolovschi et al. 2012a).

Primers targeting the Anaplasmataceae 23S, *Borrelia* 16S rRNA and *Rickettsia ompA* genes (Table 1), were used as described previously (Socolovschi et al. 2012a; Dahmani et al. 2015a; Ouarti et al. 2020).

Fig. 1 Geographical position of the tick collection locality in the south-eastern region of Slovakia




Table 1 Standard PCR primers used for the detection of microorganisms in ticks

Bacterial species	Targeted sequences	Primers (5'–3')	Reference
Anaplasmataceae	23S rRNA	f_ATAAGCTGCGGGGAATTGTC	(Dahmani et al. 2015a)
<i>Rickettsia</i> spp.	<i>ompA</i>	r_TGCAAAAGGTACGCTGTAC f_70_ATGGCGAATATTCTCCAAA A r_701_GTTCCGTTAATGGC AGCATCT f_180_GCAGCGATAATGCT GAGTA	(Socolovschi et al. 2012a)
<i>Borrelia</i> spp.	16S rRNA	f_GCTGGCAGTGCCTTAAGC r_GCTTCGGGTATCCTCAACT	(Socolovschi et al. 2012a)

The purified DNA products were sequenced using a Big Dye Terminator kit and a genetic analyzer ABI PRISM 3130 (Applied BioSystems, Courtabouef, France). The sequences obtained were analyzed with the software ChromasPro, version 1.34 (Technelysium Pty, Ltd., Tewantin, Queensland, Australie) in order to compare them to the GenBank database using BLAST (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) (Diarra et al. 2020).

Results

A total of 232 adult ticks (63 *H. inermis* males and 110 females, 31 *D. reticulatus* males and 28 females) were collected from vegetation in the protected zone of the Slovak Karst National Park. The proportion of ticks examined are detailed in (Table 2).

Using qPCR, 14/59 (23.72 %) *D. reticulatus* ticks tested positive for *Rickettsia* spp., 2/173 (1.15 %) *H. inermis* ticks tested positive for Anaplasmataceae bacteria, 1/173 (0.57 %) for *Borrelia* spp., and 29/173 (16.76 %) for *Rickettsia* spp.

The 43 ticks positive for *Rickettsia* in qPCR were then subjected to standard PCR amplification.

A total of 13/43 (30.23 %) ticks were successfully sequenced. BLAST analysis of these 13 sequences obtained showed an identity percentage ranging from 97.98 to 100 % with the *R. raoultii* sequence. Details are given in Table 3. No *Borrelia* spp. or Anaplasmataceae sequences could be amplified.

Discussion

The investigated locality (Slovak karst near the village of Hrhov) is specific and unique due to the joint occurrence of seven tick species, *Ixodes ricinus* (Linnaeus, 1758), *I. trianguliceps* Birula, 1895, *I. frontalis* (Panzer, 1795), *Dermacentor marginatus* Sulzer, 1776, *D. reticulatus*, *Haemaphysalis concinna* C. L. Koch, 1844 and *H. inermis* (Černý 1972; Nosek 1972; Bona and Stanko 2013; Heglasová et al. 2020), which is an exceptional phenomenon in the conditions of Central Europe. *Haemaphysalis punctata* Canestrini & Fanzago, 1878 which was the dominant species in tick communities in the area of Slovak karst in the last century, has not been confirmed there in recent decades. The high concentration of potential vectors (seven tick species) and hosts (cattle, free-living ungulates, several species of rodents and shrews) in the studied locality makes the Slovak karst a unique model for studying the circulation of several species of pathogens in their natural habitat (Nosek 1971; Heglasová et al. 2020). Uniqueness of the site (hereby presented) is from the point of view of the common occurrence of several species of ticks (see above). In terms of uniqueness or differentiation of pathogens in ticks compared to other sites, confirmation of a variety of bacteria in *Haemaphysalis* spp. which live on a common site with both species the genus *Dermacentor* is exclusive. In Slovakia, the presence of *D. reticulatus* has been identified along the rivers in southwestern and south-eastern Slovakia (Nosek 1972). Ticks of the genus *Haemaphysalis* (*H. inermis*, *H. concinna*, *H. punctata*) are known to have a more focal distribution (Černý 1972; Nosek 1973).

Table 2 The proportion of ticks examined according to microhabitats (forest: ecotone: meadow)

Species of ticks	Forest	Ecotone	Meadow	Total
<i>Dermacentor reticulatus</i>	2 (3.4%)	20 (33.9%)	37 (62.7%)	59
<i>Haemaphysalis inermis</i>	27 (15.61%)	121(69.94%)	25 (14.45%)	173

Table 3 BLAST analysis of *Rickettsia* obtained from tested ticks

Ticks species	Query cover (%)	Percent identity (%)	Accession number	Molecular identification by BLAST
<i>Dermacentor reticulatus</i>	100	97.98	HM161789.1	<i>Rickettsia raoultii</i>
<i>Dermacentor reticulatus</i>	100	100	KX506737.1	<i>Rickettsia raoultii</i>
<i>Dermacentor reticulatus</i>	100	99.84	HM161789.1	<i>Rickettsia raoultii</i>
<i>Dermacentor reticulatus</i>	100	100	KX506737.1	<i>Rickettsia raoultii</i>
<i>Dermacentor reticulatus</i>	100	99.68	HM161792.1	<i>Rickettsia raoultii</i>
<i>Haemaphysalis inermis</i>	99	99.84	CP003426.1	<i>Rickettsia raoultii</i>
<i>Haemaphysalis inermis</i>	99	98.63	CP003426.1	<i>Rickettsia raoultii</i>
<i>Haemaphysalis inermis</i>	100	99.41	CP003426.1	<i>Rickettsia raoultii</i>
<i>Haemaphysalis inermis</i>	99	99.84	HM161789.1	<i>Rickettsia raoultii</i>
<i>Dermacentor reticulatus</i>	100	100	KX506737.1	<i>Rickettsia raoultii</i>
<i>Dermacentor reticulatus</i>	100	99.84	HM161789.1	<i>Rickettsia raoultii</i>
<i>Haemaphysalis inermis</i>	100	100	KX506737.1	<i>Rickettsia raoultii</i>
<i>Dermacentor reticulatus</i>	99	100	HM161792.1	<i>Rickettsia raoultii</i>

In previously published studies from the Slovak karst area, we confirmed several species of rickettsiae (*R. slovacica*, *R. helvetica*, *Rickettsia felis*, *Rickettsia* sp.) and *Borrelia miyamotoi* in small mammals (especially in rodents), from rodent-attached ticks, as well as from questing ticks sampled from vegetation (Radzijejskaja et al. 2015; Heglasová et al. 2018). There is a characteristically high concentration of deer in the area, as well as cattle grazing on the pastures and both vertebrate groups are hosts for adult ticks. These factors increase the significance of the study area from an epidemiological point of view (Černý 1972).

Here, we demonstrated the circulation of *R. raoultii* in the studied area. Although *R. raoultii* has been already detected in *D. reticulatus* and in patients bitten by *D. reticulatus* (Parola et al. 2009), the vector competence of *D. marginatus* and *H. inermis* to transmit *R. raoultii* cannot be confirmed, because the detection of this bacterium in the latter could be the consequence of a bacterial blood meal of these ticks. Therefore, additional epidemiological studies and experimental models will be needed.

Both *Dermacentor* and *Haemaphysalis* ticks from which confirmed positive cases of *Rickettsia* originate are typical species for forest-steppe zones (*D. marginatus*, *H. inermis*), alluvial forests and wet meadows, which are commonly seen biotopes in eastern Slovakia (*D. reticulatus*, *H. concinna*). The joint occurrence of steppe landscape and xerothermic tick species together with ticks that prefer humid habitats in the

studied area is probably caused by the presence of ponds and canals in the karst area as well as by the high density of hosts (birds, small mammals, wild ungulates, cattle) for all life stages of ticks. *Dermacentor reticulatus* was described as harbouring more bacteria than *D. marginatus* (Zhang et al. 2019). *Rickettsia raoultii* is usually associated with *Dermacentor* spp. (Špitalská et al. 2012; Švehlová et al. 2014). Sequencing partial *ompB* genes revealed the presence of *R. raoultii* in the larvae and nymphs of *D. reticulatus* ticks in Germany (Schmuck et al. 2020). In recent decades, the spread of *D. reticulatus* to new territories has been confirmed, having been reported in, for example, in Poland (Kiewra and Czulowska 2013; Mierzejewska et al. 2016), Germany (Dautel et al. 2006) as well as other European countries (Földvári et al. 2016; Kjær et al. 2019; Capligina et al. 2020). One study carried out in Latvia detected the presence of *R. raoultii* in *D. reticulatus*, their results corroborate with our study. Experimental models will therefore be necessary to understand the role of *D. reticulatus* in the appearance of *R. raoultii* (Capligina et al. 2020).

The indication of *H. inermis* harbouring rickettsiae was noted (Řeháček et al. 1976a, b). However, the role of *Haemaphysalis* spp. as vectors of *Rickettsia* spp. is unknown (Minichová et al. 2017). There is little data on bacterial associations with *H. inermis*. *Rickettsia aeschlimannii* was detected in La Rioja, Spain (Portillo et al. 2008). *Rickettsia slovacica* (Ibarra et al. 2006) and *R. raoultii* are responsible for tick-

borne lymphadenopathy/ Dermacentor-borne necrosis erythema and lymphadenopathy/scalp eschar and neck lymphadenopathy (TIBOLA-/DEBONEL/SENLAT) (Oteo et al. 2004; Selmi et al. 2008; Parola et al. 2013). This illness commonly occurs in Slovakia (Sekeyová et al. 2013) and was recently identified as the causative agent of tick-borne lymphadenopathy in Belgium (Lernout et al. 2019). *Rickettsia helvetica* is considered less pathogenic than the two aforementioned rickettsial pathogens (Sprong et al. 2009; Boulanger et al. 2019).

Conclusions

This molecular study supports the data that *R. raoultii* could be a source of lymphadenopathy in Slovak populations. Further studies, including experimental models, will be needed to assess the role of *D. reticulatus* and *H. inermis* in the transmission of *R. raoultii*. Nevertheless, based on the positive detection of *R. raoultii*, we assume that *Haemaphysalis* and *Dermacentor* spp. should be considered as potential reservoirs of *R. raoultii* in Slovakia.

Acknowledgements The authors thank Ladislav Mošanský for his help with tick collection.

Author Contributions ZS, PP and OM conceived and designed the research. BO, BE and ML conducted the experiments. MS, and ZS contributed analytical tools. BO and BE analysed the data. BO and ZS wrote the manuscript. All authors read and approved the manuscript.

Funding This study was supported by the Institut Hospitalo-Universitaire (IHU) Méditerranée Infection, the French National Research Agency under the “Investissements d’avenir” programme, reference ANR-10-IAHU-03, the Région Provence-Alpes-Côte d’Azur and the European funding ERDF PRIMI. Z.S. acknowledges financial support from SRDA-19-0066 and VEGA 2/0010/19, and 2/0021/21, from the Scientific Grant Agency of the Ministry of Education, Science, Research and Sport.

Declarations

Conflict of interest The authors have no conflicts of interest to disclose.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Aouadi A, Leulmi H, Boucheikhchoukh M, Benakhla A, Raoult D, Parola P (2017) Molecular evidence of tick-borne hemoprotozoan parasites (*Theileria ovis* and *Babesia ovis*) and bacteria in ticks and blood from small ruminants in Northern Algeria. *Comp Immunol Microbiol Infect Dis* 50:34–39. <https://doi.org/10.1016/j.cimid.2016.11.008>
- Boldiš V, Kocianová E, Štrus J, Tušek-Žnidarič M, Sparagano OA, Štefanidesová K, Špitalská E (2008) Rickettsial agents in Slovakian ticks (Acarina, Ixodidae) and their ability to grow in Vero and L929 cell lines. *Ann N Y Acad Sci* 1149:281–285. <https://doi.org/10.1196/annals.1428.090>
- Bona M, Stanko M (2013) First records of the tick *Ixodes frontalis* (Panzer, 1795) (Acari, Ixodidae) in Slovakia. *Ticks Tick Borne Dis* 4:478–481. <https://doi.org/10.1016/j.ttbdis.2013.06.002>
- Boulanger N, Boyer P, Talagrand-Reboul E, Hansmann Y (2019) Ticks and tick-borne diseases. *Med Mal Infect* 49:87–97. <https://doi.org/10.1016/j.medmal.2019.01.007>
- Caplignina V, Seleznova M, Akopjana S, Freimane L, Lazovska M, Krumins R, Kivrane A, Namina A, Aleinikova D, Kimsis J (2020) Large-scale countrywide screening for tick-borne pathogens in field-collected ticks in Latvia during 2017–2019. *Parasit Vectors* 13:1–12. <https://doi.org/10.1186/s13071-020-04219-7>
- Černý V (1972) The tick fauna of Czechoslovakia. *Folia Parasitol* 19:87–92
- Chvostáč M, Špitalská E, Václav R, Vaculová T, Minichová L, Derdánková M (2018) Seasonal patterns in the prevalence and diversity of tick-borne *Borrelia burgdorferi* sensu lato, *Anaplasma phagocytophilum* and *Rickettsia* spp. in an urban temperate forest in south western Slovakia. *Int J Environ Res Publ Health* 15:994. <https://doi.org/10.3390/ijerph15050994>
- Dahmani M, Davoust B, Benterki MS, Fenollar F, Raoult D, Mediannikov O (2015a) Development of a new PCR-based assay to detect Anaplasmataceae and the first report of *Anaplasma phagocytophilum* and *Anaplasma platys* in cattle from Algeria. *Comp Immunol Microbiol Infect Dis* 39:39–45. <https://doi.org/10.1016/j.cimid.2015.02.002>
- Dahmani M, Loudahi A, Mediannikov O, Fenollar F, Raoult D, Davoust B (2015b) Molecular detection of *Anaplasma platys* and *Ehrlichia canis* in dogs from Kabylie, Algeria. *Ticks Tick Borne Dis* 6:198–203. <https://doi.org/10.1016/j.ttbdis.2014.12.007>
- Dautel H, Dippel C, Oehme R, Hartelt K, Schettler E (2006) Evidence for an increased geographical distribution of *Dermacentor reticulatus* in Germany and detection of *Rickettsia* sp. RpA4. *Int J Med Microbiol* 296:149–156. <https://doi.org/10.1016/j.ijmm.2006.01.013>
- de la Fuente J, Estrada-Pena A, Venzal JM, Kocan KM, Sonenshine DE (2008) Overview: ticks as vectors of pathogens that cause disease in humans and animals. *Front Biosci* 13:6938–6946. <https://doi.org/10.2741/3200>
- Diarra AZ, Almeras L, Laroche M, Berenger J-M, Koné AK, Bocoum Z, Dabo O, Doumbo O, Raoult D, Parola P (2017) Molecular and MALDI-TOF identification of ticks and tick-associated bacteria in Mali. *PLoS Negl Trop Dis* 11:e0005762. <https://doi.org/10.1371/journal.pntd.0005762>
- Diarra AZ, Kone AK, Niare SD, Laroche M, Diatta G, Atteynine SA, Coulibaly M, Sangare AK, Kouriba B, Djimde A (2020) Molecular detection of microorganisms associated with small mammals and their ectoparasites in Mali. *Am J Trop Med Hyg* 103:2542–2551. <https://doi.org/10.4269/ajtmh.19-0727>
- Eldin C, Parola P (2018) Update on tick-borne bacterial diseases in travelers. *Curr Infect Dis Rep* 20:17. <https://doi.org/10.1007/s11908-018-0624-y>

- Földvári G, Rigó K, Lakos A (2013) Transmission of *Rickettsia slovaca* and *Rickettsia raoultii* by male *Dermacentor marginatus* and *Dermacentor reticulatus* ticks to humans. *Diagn Microbiol Infect Dis* 76:387–389. <https://doi.org/10.1016/j.diagmicrobio.2013.03.005>
- Földvári G, Široký P, Szekeres S, Majoros G, Sprong H (2016) *Dermacentor reticulatus*: a vector on the rise. *Parasit Vectors* 9:1–29. <https://doi.org/10.1186/s13071-016-1599-x>
- García-Vozmediano A, Giglio G, Ramassa E, Nobili F, Rossi L, Tomassone L (2020) *Dermacentor marginatus* and *Dermacentor reticulatus*, and their infection by SFG Rickettsiae and *Francisella-Like* Endosymbionts, in mountain and periurban habitats of Northwestern Italy. *Vet Sci* 7:157. <https://doi.org/10.3390/vetsci7040157>
- Heglasová I, Vichová B, Kraljik J, Mošanský L, Miklisová D, Stanko M (2018) Molecular evidence and diversity of the spotted-fever group *Rickettsia* spp. in small mammals from natural, suburban and urban areas of Eastern Slovakia. *Ticks Tick Borne Dis* 9:1400–1406. <https://doi.org/10.1016/j.ttbdis.2018.06.011>
- Heglasová I, Rudenko N, Golovchenko M, Zubriková D, Miklisová D, Stanko M (2020) Ticks, fleas and rodent-hosts analyzed for the presence of *Borrelia miyamotoi* in Slovakia: the first record of *Borrelia miyamotoi* in a *Haemaphysalis inermis* tick. *Ticks Tick Borne Dis* 11:101456. <https://doi.org/10.1016/j.ttbdis.2020.101456>
- Hornok S, Meli ML, Perreten A, Farkas R, Willi B, Beugnet F, Lutz H, Hofmann-Lehmann R (2010) Molecular investigation of hard ticks (Acari: Ixodidae) and fleas (Siphonaptera: Pulicidae) as potential vectors of rickettsial and mycoplasmal agents. *Vet Microbiol* 140:98–104. <https://doi.org/10.1016/j.vetmic.2009.07.013>
- Ibarra V, Oteo J, Portillo A, Santibañez S, Blanco J, Metola L, Eiros Bouza JM, Pérez Martínez L, Sanz M (2006) *Rickettsia slovaca* infection: DEBONEL/TIBOLA. *Ann N Y Acad Sci* 1078:206–214. <https://doi.org/10.1196/annals.1374.040>
- Kiewra D, Czulowska A (2013) Evidence for an increased distribution range of *Dermacentor reticulatus* in south-west Poland. *Exp Appl Acarol* 59:501–506. <https://doi.org/10.1007/s10493-012-9612-3>
- Kjær LJ, Soleng A, Edgar KS, Lindstedt HEH, Paulsen KM, Andreassen ÅK, Korslund L, Kjelland V, Slettan A, Stuen S (2019) A large-scale screening for the taiga tick, *Ixodes persulcatus*, and the meadow tick, *Dermacentor reticulatus*, in southern Scandinavia, 2016. *Parasit Vectors* 12:338. <https://doi.org/10.1186/s13071-019-3596-3>
- Lernout T, De Regge N, Tersago K, Fonville M, Suin V, Sprong H (2019) Prevalence of pathogens in ticks collected from humans through citizen science in Belgium. *Parasit Vectors* 12:550. <https://doi.org/10.1186/s13071-019-3806-z>
- Mierzejewska EJ, Estrada-Peña A, Alsarraf M, Kowalec M, Bajer A (2016) Mapping of *Dermacentor reticulatus* expansion in Poland in 2012–2014. *Ticks Tick Borne Dis* 7:94–106. <https://doi.org/10.1016/j.ttbdis.2015.09.003>
- Minichová L, Hamšíková Z, Mahriková L, Slovák M, Kocianová E, Kazimirová M, Škultéty L, Štefanidesová K, Špitalská E (2017) Molecular evidence of *Rickettsia* spp. in ixodid ticks and rodents in suburban, natural and rural habitats in Slovakia. *Parasit Vectors* 10:158. <https://doi.org/10.1186/s13071-017-2094-8>
- Nosek J (1971) The ecology, bionomics, and behaviour of *Haemaphysalis (Aboimialis) punctata* tick in central Europe. *Ztschr Parasitenk* 37:198–210. <https://doi.org/10.1007/BF00259499>
- Nosek J (1972) The ecology and public health importance of *Dermacentor marginatus* and *D. reticulatus* ticks in central Europe. *Folia Parasitol* 19:93–102
- Nosek J (1973) Some characteristic features of the life history, ecology and behaviour of the ticks *Haemaphysalis inermis*, *H. concinna* and *H. punctata*. In: Daniel M, Rosický B (eds) Proceedings of the 3rd International Congress of Acarology, Springer, Dordrecht, pp 479–482. https://doi.org/10.1007/978-94-010-2709-0_90
- Oteo J, Ibarra V, Blanco J, de Artola VM, Márquez F, Portillo A, Raoult D, Anda P (2004) *Dermacentor*-borne necrosis erythema and lymphadenopathy: clinical and epidemiological features of a new tick-borne disease. *Clin Microbiol Infect* 10:327–331. <https://doi.org/10.1111/j.1198-743X.2004.00782.x>
- Ouarti B, Mediannikov O, Righi S, Benakhla A, Raoult D, Parola P (2020) Molecular detection of microorganisms in lice collected from farm animals in Northeastern Algeria. *Comp Immunol Microbiol Infect Dis* 74:101569–101569. <https://doi.org/10.1016/j.cimid.2020.101569>
- Parola P, Roveery C, Rolain JM, Brouqui P, Davoust B, Raoult D (2009) *Rickettsia slovaca* and *R. raoultii* in tick-borne rickettsioses. *Emerg Infect Dis* 15:1105. <https://doi.org/10.3201/eid1507.081449>
- Parola P, Paddock CD, Socolovschi C, Labruna MB, Mediannikov O, Kernif T, Abdad MY, Stenos J, Bitam I, Fournier P-E (2013) Update on tick-borne rickettsioses around the world: a geographic approach. *Clin Microbiol Rev* 26:657–702. <https://doi.org/10.1128/CMR.00032-13>
- Portillo A, Santibañez P, Santibañez S, Pérez-Martínez L, Oteo JA (2008) Detection of *Rickettsia* spp. in *Haemaphysalis* ticks collected in La Rioja, Spain. *Vector Borne Zoonotic Dis* 8:653–658. <https://doi.org/10.1089/vbz.2007.0272>
- Radzijeuskaja J, Paulauskas A, Aleksandraviciene A, Jonauskaitė I, Stanko M, Karbowiak G, Petko B (2015) New records of spotted fever group rickettsiae in Baltic region. *Microbes Infect* 17:874–878. <https://doi.org/10.1016/j.micinf.2015.09.006>
- Řeháček J, Kováčová E, Kováč P (1976a) Rickettsiae belonging to the spotted fever group from ticks in the Tribeč mountains. *Folia Parasitol* 23:69–73
- Řeháček J, Župančičová M, Āč P, Brezina R, Úrvölgyi J, Kováčová E, Tarasevič I, Jablonskaja V, Pospíšil R, Baloghova D (1976b) Rickettsioses studies: 2. Natural foci of rickettsioses in east Slovakia. *Bull World Health Org* 53:31–38
- Schmuck HM, Chitimia-Dobler L, Król N, Kacza J, Pfeiffer M (2020) Collection of immature *Dermacentor reticulatus* (Fabricius, 1794) ticks from vegetation and detection of *Rickettsia raoultii* in them. *Ticks Tick Borne Dis* 11:101543. <https://doi.org/10.1016/j.ttbdis.2020.101543>
- Sekeyova Z, Roux V, Xu W, Řeháček J, Raoult D (1998) Note: *Rickettsia slovaca* sp. nov., a member of the spotted fever group rickettsiae. *Int J Syst Evol Microbiol* 48:1455–1462. <https://doi.org/10.1099/00207713-48-4-1455>
- Sekeyová Z, Socolovschi C, Špitalská E, Kocianová E, Boldiš V, Quevedo Diaz M, Berthová L, Boháčová M, Valáriková J, Fournier PE (2013) Update on rickettsioses in Slovakia. *Acta Virol* 57:180–199. https://doi.org/10.4149/av_2013_02_180
- Selmi M, Bertolotti L, Tomassone L, Mannelli A (2008) *Rickettsia slovaca* in *Dermacentor marginatus* and tick-borne lymphadenopathy Tuscany, Italy. *Emerg Infect Dis* 14:817. <https://doi.org/10.3201/eid1405.070976>
- Socolovschi C, Kernif T, Raoult D, Parola P (2012a) *Borrelia*, *Rickettsia*, and *Ehrlichia* species in bat ticks, France, 2010. *Emerg Infect Dis* 18:1966–1975. <https://doi.org/10.3201/eid1812.111237>
- Socolovschi C, Pagés F, Raoult D (2012b) *Rickettsia felis* in *Aedes albopictus* mosquitoes, Libreville, Gabon. *Emerg Infect Dis* 18:1687–1689. <https://doi.org/10.3201/eid1810.120178>
- Špitalská E, Štefanidesová K, Kocianová E, Boldiš V (2012) *Rickettsia slovaca* and *Rickettsia raoultii* in *Dermacentor marginatus* and *Dermacentor reticulatus* ticks from Slovak Republic. *Exp Appl Acarol* 57:189–197. <https://doi.org/10.1007/s10493-012-9539-8>
- Špitalská E, Sparagano O, Stanko M, Schwarzová K, Špitalský Z, Škultéty L, Fumačová Havlíková S (2018) Diversity of *Coxiella-like* and *Francisella-like* endosymbionts, and *Rickettsia* spp., *Coxiella burnetii* as pathogens in the tick populations of Slovakia. Central Europe. *Ticks Tick Borne Dis* 9:1207–1211. <https://doi.org/10.1016/j.ttbdis.2018.05.002>

- Sprong H, Wielinga PR, Fonville M, Reusken C, Brandenburg AH, Borgsteede F, Gaasenbeek C, van der Giessen JW (2009) *Ixodes ricinus* ticks are reservoir hosts for *Rickettsia helvetica* and potentially carry flea-borne *Rickettsia* species. *Parasit Vectors* 2:41. <https://doi.org/10.1186/1756-3305-2-41>
- Švehlová A, Berthová L, Sallay B, Boldiš V, Sparagano OA, Špitalská E (2014) Sympatric occurrence of *Ixodes ricinus*, *Dermacentor reticulatus* and *Haemaphysalis concinna* ticks and *Rickettsia* and *Babesia* species in Slovakia. *Ticks Tick Borne Dis* 5:600–605. <https://doi.org/10.1016/j.ttbdis.2014.04.010>
- Zhang YK, Yu ZJ, Wang D, Vichová B, Peťko B, Liu JZ (2019) The bacterial microbiome of field-collected *Dermacentor marginatus* and *Dermacentor reticulatus* from Slovakia. *Parasit Vectors* 12: 325. <https://doi.org/10.1186/s13071-019-3582-9>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.