



## GPCRs in invertebrate innate immunity

Jerome Reboul, Jonathan J Ewbank

### ► To cite this version:

Jerome Reboul, Jonathan J Ewbank. GPCRs in invertebrate innate immunity. *Biochemical Pharmacology*, 2016, 114, pp.82-87. 10.1016/j.bcp.2016.05.015 . hal-03540828

**HAL Id: hal-03540828**

**<https://amu.hal.science/hal-03540828>**

Submitted on 24 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

**GPCRs in invertebrate innate immunity**

Jerome Reboul and Jonathan J. Ewbank

Centre d'Immunologie de Marseille-Luminy, Aix Marseille Université UM2, Inserm,  
U1104, CNRS UMR7280, 13288 Marseille, France

Corresponding author: Jonathan Ewbank

Centre d'Immunologie de Marseille-Luminy,  
163 avenue de Luminy  
13009 Marseille  
FRANCE

ewbank@ciml.univ-mrs.fr

Tel: +33 491 269 472

## ABSTRACT

G-protein coupled receptors (GPCRs) represent a privileged point of contact between cells and their surrounding environment. They have been widely adopted in vertebrates as mediators of signals involved in both innate and adaptive immunity. Invertebrates rely on innate immune defences to resist infection. We review here evidence from a number of different species, principally the genetically tractable *Caenorhabditis elegans* and *Drosophila melanogaster* that points to an important role for GPCRs in modulating innate immunity in invertebrates too. In addition to examples of GPCRs involved in regulating the expression of defence genes, we discuss studies in *C. elegans* addressing the role of GPCR signalling in pathogen aversive behaviour. Despite the many lacunae in our current knowledge, it is clear that GPCR signalling contributes to host defence across the animal kingdom.

## 1. Introduction

Compared to the extensive literature linking G-protein coupled receptors (GPCRs) to immunity in vertebrates, the subject has been relatively poorly explored in invertebrate animals. In this review, we will cover early biochemical studies using the horseshoe crab, then more recent work with the genetically tractable models *Caenorhabditis elegans* and *Drosophila melanogaster*, and lastly a brief overview of research with diverse non-model systems. For the sake of restricting this review to a single phylogenetic group of animals, we will not include any studies on chordates, such as Ascidians.

Lacking adaptive immunity, invertebrates rely on their innate immune system to defend themselves against infection. The first step in triggering an immune response involves recognition of stranger/danger signals. Stranger signals are molecular hallmarks of a particular group of microbial species, hence their alternative name of microbe/pathogen-associated molecular patterns (MAMP/PAMP). The archetypal MAMP is lipopolysaccharide, an indispensable component of the outer membrane of Gram-negative bacteria. MAMPs are recognized by dedicated receptors, such as Toll-like receptor (TLR) 4 in vertebrates [1]. Danger signals, also known as damage-associated molecular patterns (DAMPs), on the other hand, can be endogenous molecules, like mitochondrial proteins or ATP, aberrantly released into the extracellular milieu [2]. The molecular architecture of GPCRs is particularly well suited to allow binding of diverse chemical structures, from small organic molecules to neuropeptides (reviewed in [3]). They are thus good candidates for mediating perception of diverse danger signals in host cells.

## 2. GPCRs in horseshoe crab immunity

### 2.1. LPS-triggered signalling

The first indications of a role for GPCRs in defence came from work with horseshoe crabs. These are large, evolutionary ancient, marine arthropods, with a predicted origin some 450 million years ago. Of the 4 known species, primarily *Limulus polyphemus* and *Tachypleus tridentatus* have been used in research. Their blood contains motile cells called hemocytes or amebocytes, which are thought to play a similar role in host defence to macrophages in vertebrates. More than half a century ago, it was shown that exposing *Limulus* blood to Gram-negative bacteria provoked a rapid coagulation [4]. This response requires exocytosis of the clotting factor coagulogen, and is triggered by recognition of bacterial lipopolysaccharide (LPS) [5]. Interestingly, coagulogen is structurally similar to the *Drosophila* protein spaetzle [6]. In both species these proteins are ligands for immune receptors of the Toll family required for activation of the key transcription factor NFκB. This has led to the suggestion of an evolutionary ancient origin for NFκB signalling in defence throughout animals [7]. LPS recognition itself relies on the zymogen factor C [8]. Binding of LPS triggers an auto-activation of factor C to give rise to an active serine protease. By analogy with the activation of vertebrate GPCRs of the protease-activated receptor family (PARs) by thrombin, it is believed that the factor C protease would cleave the N-terminus of an as yet unidentified GPCR leading to the activation of diverse cellular responses (Figure 1; reviewed in [9]). There are clear parallels between this system and the PAR-dependent activation of NFκB in vertebrates [10].

### 2.2. Potentiation by tachyplesin

Hemocytes can also be activated in the absence of LPS by host-derived peptides including tachyplesin [11], originally described as an antimicrobial peptide (AMP) [12]. Tachyplesin can interact *in vitro* with G<sub>o</sub> $\alpha$  and G<sub>i</sub> $\alpha$  proteins from bovine brain, and both U-73122, an inhibitor of phospholipase C, and pertussis toxin, a G protein inhibitor, strongly inhibit hemocyte exocytosis. Although tachyplesin has not been demonstrated to activate G proteins, these results provide further evidence for a role of GPCR signalling in horseshoe crab defence and suggest that a positive feedback mechanism for hemocyte secretion exists [11].

### 3. GPCRs in worm immunity

#### 3.1. Defence via CO<sub>2</sub>/O<sub>2</sub> detection and chemical sensing

Nematodes live in a microbe-rich environment and eat bacteria. They need to be able to distinguish innocuous bacteria from potential pathogens. Data from a broad range of species indicates that the Toll-like receptor (TLR) - NF $\kappa$ B signalling axis mentioned above has been lost from nematodes [13]. There are no obvious NF $\kappa$ B orthologs in worms and although *C. elegans* has one TLR, TOL-1 [14], it appears not to have a direct role in host defence (reviewed in [15]). It does, however, have the potential to influence the interaction between *C. elegans* and bacteria, including pathogens [14] since *tol-1* is required for the terminal differentiation of a class of neurons required for detection of CO<sub>2</sub> [16]. Like any metabolically active bacteria, pathogens will produce CO<sub>2</sub>, and locally depress O<sub>2</sub> levels. *C. elegans* is believed to measure gradients of O<sub>2</sub> and CO<sub>2</sub> to guide it to bacteria. It also integrates a variety of other sensory cues, both attractive and repulsive, via GPCR chemoreceptors [17], to distinguish good from bad food. *C. elegans* thus has the capacity to recognise

different foods and make choices based on the CO<sub>2</sub>/O<sub>2</sub> balance in combination with several other factors, including via chemosensation [18].

### 3.1.1. G<sub>i</sub>α-like protein ODR-3 and the GPCR kinase GRK-2 in *Serratia marcescens* avoidance

As an example, *C. elegans* detects the *Serratia marcescens*-derived surfactant serrawettin W2 via the AWB chemosensory neurons and is repelled by it, even in the presence of an attractive CO<sub>2</sub>/O<sub>2</sub> environment. Thus worms will avoid a bacterial lawn that contains serrawettin W2 [19]. Since *S. marcescens* is a pathogen of *C. elegans* [20], this clearly has the potential to contribute to host survival. The AWB neurons express a range of GPCRs that all require the G<sub>i</sub>α-like protein ODR-3 and the GPCR kinase GRK-2 for their function in chemosensation [21, 22]. Phosphorylation by GRKs allows GPCRs to bind the negative regulatory protein arrestin. This blocks further G protein-mediated signalling and targets GPCRs for internalization. Mutants in either *odr-3* or *grk-2* are defective in their avoidance of lawns containing serrawettin W2, suggesting a role for one or more GPCR in this type of pathogen avoidance behaviour [19].

### 3.1.2. GPCR signalling in *Pseudomonas aeruginosa* avoidance

Worms also avoid the pathogen *Pseudomonas aeruginosa*. This was shown recently to depend on detection of the bacterial secondary metabolites phenazine-1-carboxamide and pyochelin. These are recognised by an as yet undefined GPCR that acts via the Gα proteins GPA-2 and GPA-3. One consequence of this chemosensation is an increased neuronal secretion of the TGFβ orthologue DAF-7. This cytokine modifies how worms respond to ambient oxygen levels, so that they leave a lawn of *P. aeruginosa* in spite of an oxygen concentration that would normally be attractive [23].

1 There is plasticity in the behavioural response to pathogenic bacteria. *P. aeruginosa*  
2 induces aversive olfactory learning so that given a choice between *P. aeruginosa* and  
3 non-pathogenic *E. coli*, worms prefer *E. coli* [24]. For this, the AWB neurons act  
4 together with a second pair, called AWC, again requiring the function of the G<sub>i</sub>α-like  
5 protein ODR-3. Food preferences require the neuropeptide NLP-9 produced in AWB  
6 and its putative receptor, the GPCR NPR-18, as well as NLP-1 produced in AWC  
7 [25]. AWB and AWC act upstream of ADL sensory neurons to control the preference  
8 behaviour that can limit exposure to bacterial pathogens. In this context, recent work  
9 has suggested a role for several other neuropeptides, including NLP-10, for which the  
10 receptor is not known, and FLP-4 released from ADL, acting through its receptor, the  
11 GPCR NPR-4, in AIB interneurons. It also indicated a possible role in ADL for the  
12 orphan GPCR SRH-220. Loss of function *srh-220* mutants show a substantially  
13 reduced preference for *E. coli* when given a choice between it and pathogenic  
14 *P. aeruginosa*. This putative role for a GPCR is consistent with the similar deficit in  
15 odour preference observed in mutants for the unique arrestin orthologue ARR-1. This  
16 phenotype was rescued when ARR-1 was specifically expressed in the ADL sensory  
17 neurons [26]. A full understanding of these mechanisms will require the identification  
18 of the ligand for SRH-220.

### 3.1.3. The role of the neuropeptide receptor NPR-1 in immune responses

45 The GPCR neuropeptide receptor NPR-1 had been proposed to play a direct role in  
46 modulating innate immune gene expression upon *P. aeruginosa* infection. This has  
47 been challenged by other studies that suggested an indirect role for NPR-1 in defence  
48 (reviewed in [27, 28]). Indeed, *npr-1* mutants exhibit a broad range of behavioural  
49 phenotypes [29], including a change in CO<sub>2</sub> and O<sub>2</sub> sensation [30] that impact on its  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



capacity to avoid pathogenic bacteria. These could explain the observed alterations in the interaction between *npr-1* mutants and pathogens.

#### 3.1.4. A broad role for a GPCR-LRR protein FSHR-1 in defence

Another GPCR, the follicle stimulating hormone receptor homologue FSHR-1, is also required for the worm's capacity to recognise and avoid pathogenic bacteria. Moreover it is needed for the expression of defence genes in the gut of worms infected with *P. aeruginosa* [31]. FSHR-1 additionally acts in the intestine to regulate genes required to resist heavy metal and oxidative stress, thereby contributing to the maintenance or re-establishment of homeostasis following infection [31]. Finally, FSHR-1 antagonizes the capacity of *C. elegans* to resist low temperatures; mutants lacking *fshr-1* function survive cold stress better than wild-type worms [32]. FSHR-1 was originally studied because of its structure; it combines GPCR and leucine-rich repeat (LRR) domains [33]. LRR domains are frequently found in innate immune receptors, including TLRs and most NOD-like receptors (NLRs). In vertebrate TLRs, the LRRs mediate direct recognition with the appropriate MAMP. In invertebrates, the LRRs do not necessarily play such a role. Rather, as mentioned above, they bind endogenous cytokine-like proteins [34, 35]. Given its pleiotropic function, it is unlikely that FSHR-1 acts as a MAMP receptor in worms. Nevertheless, it will be very interesting to discover its physiological ligand.

#### 3.2. The DAMP receptor, DCAR-1

One GPCR that acts as a DAMP receptor has been identified in *C. elegans*. DCAR-1 was originally described as being expressed in chemosensory neurons and to be a putative receptor for the DOPA-derived small molecule dihydrocaffeic acid (DHCA) [36]. It is also expressed in the nematode epidermis where it can be activated by hydroxyphenyllactic acid (HPLA), a derivative of tyrosine that accumulates when

1 worms are wounded or infected by the fungus *Drechmeria coniospora* [37]. The  
2 signal transduction pathway downstream of DCAR-1 has been extensively studied  
3  
4 through genetic and biochemical approaches (Figure 2). DCAR-1 acts upstream of the  
5  
6  $G_{12}\alpha$  protein GPA-12 [37] that in turn is upstream of 2 phospholipase C $\beta$  (PLC $\beta$ )  
7  
8 enzymes (PLC-3 and EGL-8). These hydrolyse phosphatidylinositol 4,5-bisphosphate  
9  
10 (PIP2) to produce inositol trisphosphate (IP3) and diacylglycerol (DAG). DAG then  
11  
12 activates a protein kinase C (TPA-1; [38]) that switches on a conserved p38 MAPK  
13  
14 cascade [39, 40], ultimately leading to the activation of a STAT-like transcription  
15  
16 factor and the expression of antimicrobial peptide genes in the epidermis [41].  
17  
18  
19  
20

### 21 3.3. Multiple roles for the $G_q\alpha$ EGL-30 in host defence

22  
23 A second  $G\alpha$  protein (the  $G_q\alpha$  EGL-30) functions in a cell autonomous manner within  
24  
25 the intestine via the PLC $\beta$  EGL-8 to regulate the activity of the p38 MAPK pathway  
26  
27 and thereby the expression of intestinal immune effectors upon infection with  
28  
29 *P. aeruginosa*. EGL-30 and EGL-8 additionally act in a cell non-autonomous manner,  
30  
31 by modulating the level of insulin/IGF1 signalling from neurons [42]. Secretion of  
32  
33 insulin has a wide-ranging impact on the physiology of the worm, influencing lifespan  
34  
35 as well as stress and pathogen resistance (reviewed in [28]).  
36  
37  
38  
39  
40

41 The G proteins GPA-12 and EGL-30 are also involved in the interaction between  
42  
43 *C. elegans* and *Microbacterium nematophilum*, a Gram-positive bacterium [43] that  
44  
45 worms avoid. EGL-30 intervenes at two different levels, influencing both the aversive  
46  
47 behavioural response to *M. nematophilum* mediated by neuronal activity, and an  
48  
49 epithelial defence mechanism. Thus GPA-12 and EGL-30 are both positive regulators  
50  
51 of neurotransmitter release in cholinergic motor neurons, required for the changes in  
52  
53 locomotion behaviour that underlie pathogen avoidance [44]. At the same time, EGL-  
54  
55 30 acts in the rectal epithelium to modulate the immune response, involving changes  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 in gene expression and alteration of cellular morphology. The GPCR that acts  
2 upstream of EGL-30 has recently been identified. Like DCAR-1, it appears to  
3 recognise an amino acid derivative (R. McMullan, personal communication).  
4  
5

6  
7 The EGL-30-dependent pathway functions in cooperation with a Ras signalling  
8 pathway [45]. Serotonin, released from ADF chemosensory neurons, acting directly  
9 or indirectly via serotonin receptors, the GPCRs SER-1 and SER-7, and possibly  
10 others, activates the G<sub>o</sub>α protein GOA-1 that in turn triggers a downstream protective  
11 signal transduction cascade in the rectal epithelium. The system is complex, since  
12 GOA-1 acts antagonistically to EGL-30 both in neurons and the rectal epithelium  
13 [46]. Further, the level of serotonin production in ADF depends on the expression of  
14 *tph-1*, corresponding to a rate-limiting tryptophan hydroxylase. It is modulated  
15 indirectly by a GPCR pathway in olfactory sensory neurons involving the G<sub>q</sub>α protein  
16 EGL-30 [47]. The identity of the putative GPCR that acts upstream of EGL-30 in the  
17 AWB and AWC sensory neurons for the detection of pathogens is not yet known. But  
18 it is noteworthy that the arrestin ARR-1 plays a broad role in immune regulation [48].  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

### 36 3.3. Other GPCRs in *C. elegans* defence

37 DCAR-1 was identified through a genome-wide RNAi screen for genes required for  
38 antimicrobial peptide gene expression [49]. Several other GPCRs also emerged as  
39 candidates in this screen, but their precise role in innate immunity is currently unclear  
40 [37]. In different species, GPCR families have undergone lineage-specific expansions.  
41 In the case of *C. elegans* there are more than 1500 predicted GPCRs, many involved  
42 in chemosensation [50-52]. Before any functional studies had been undertaken, *in*  
43 *silico* analyses of sequence evolution and polymorphisms in natural isolates suggested  
44 a role for certain GPCR families in mediating interactions with microbes. Among  
45 them, the SRH family stands out because of its very high functional polymorphism  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

[53, 54]. For example, there is at least one naturally occurring deletion allele of *srh-220*, a gene involved in pathogen discrimination as described above, present in the Germany isolate MY1 (see [http://www.wormbase.org/species/c\\_elegans/strain/MY1](http://www.wormbase.org/species/c_elegans/strain/MY1)). Thus GPCRs are central to diverse aspects of host-pathogen relationships in the worm, but whether SRH-220 or other GPCRs are *bona fide* MAMP receptors remains to be determined.

#### 4. GPCRs in fly immunity

The best-characterised aspect of innate immune defences in *Drosophila* is the regulation by NF $\kappa$ B of antimicrobial gene expression in the fat body following activation of Toll (by the cytokine-like protein Spaetzle) or via the peptidoglycan-triggered IMD pathway [35]. Flies additionally have the capacity to produce reactive oxygen species in the intestine through the activity of DUOX enzymes. These contribute to protect flies from gut infection. In contrast to peptidoglycan-induced IMD activation, acute DUOX activation requires membrane-localised PLC $\beta$  and was therefore proposed to be principally via GPCR signalling [55]. Subsequent work revealed uracil, excreted by non-commensal bacteria in the gut, to be a trigger for increasing DUOX activity [56]. The identity of the putative uracil-binding GPCR has yet to be reported (Figure 3).

The other evidence for a role of GPCR signalling in flies is also indirect. One significant hit in a genome-wide cell-based RNAi screen for genes required for Toll and IMD pathway corresponded to the *Drosophila* G protein-coupled receptor kinase Gprk2 [57]. The immune function for this GRK was suggested to be potentially related to its established role in regulating hedgehog signalling [58, 59], perhaps

1 through an effect on cell surface receptor recycling [60]. Interestingly, uracil-  
2 dependent DUOX activation in the fly gut also requires hedgehog, which is needed  
3  
4 for the formation of a subset of endosomes that act as signalling platforms [61].  
5  
6

7 There is circumstantial evidence for a link between GPCR signalling and defence  
8  
9 against viral infection in *Drosophila*. Thus, the G $\beta$ -like protein RACK1 (Receptor for  
10 Activated C Kinase 1) that acts downstream of the GPCR DCAR-1 in *C. elegans* [37,  
11 38] is required for the translation of Cricket Paralysis Virus (CrPV) proteins in fly  
12 cells. It should be noted, however, that RACK1 is also a ribosomal protein and its role  
13 in viral replication, which is linked to IRES-dependent translation, may be totally  
14 independent of GPCRs [62]. As a further example, *Drosophila* can be parasitized by  
15 wasps that deposit their eggs in larvae. Wasp secretions, transferred into the host with  
16 the egg, can contain symbiotic viruses that suppress the normal host defences that  
17 lead to encapsulation and killing of wasp eggs [63-65]. One viral effector, CrV2, has  
18 been demonstrated to bind directly to a host G $\alpha$  protein *in vitro* [66], raising the  
19 possibility of a wasp block of GPCR signalling important in host defence.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

## 39 **5. GPCRs in immunity in other invertebrate species**

40  
41  
42

43 If the literature on GPCRs in defence in the 2 model organisms *C. elegans* and  
44 *D. melanogaster* is sparse, in other invertebrates it is even more patchy. There are  
45 occasional reports of a role for GPCRs in immunity for a diverse mix of species.  
46 Thus, for example, in crayfish, exposure to dead Gram-negative bacteria provokes up-  
47 regulation of the GPCR-encoding gene HP1R. Knocking down HP1R expression  
48 renders animals more susceptible to infection, supporting a role for GPCR signalling  
49 in defence [67]. The response to LPS involves production of astakines, the crayfish  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

equivalent of prokineticins (PROKs). In vertebrates, these cytokines signal through GPCRs, and this is probably the case in crayfish too [68]. In shrimp, arrestins are required for defence against Gram-positive bacteria, but this has been suggested to be via a direct effect on the Cactus-Dorsal complex in the Toll signalling pathway rather than a modulation of GPCR signalling [69]. More examples are likely to be found in the future.

## 6. Concluding remarks

Evidence has accumulated for an important role for GPCRs in invertebrate innate immunity. Despite the many examples of functions for downstream components of GPCR signalling pathways, perhaps because of functional redundancy, there are extremely few cases where a single GPCR has been assigned an unambiguous and specific role in defence. Identifying innate immune GPCRs that acts as DAMP or MAMP receptors therefore remains a major challenge for the future.

## Footnote

Work in our laboratory is funded by institutional grants from AMU, INSERM and CNRS, the ANR (ANR-12-BSV3-0001-01, ANR-11-LABX-0054 (Investissements d'Avenir–Labex INFORM) and ANR-11-IDEX-0001-02 (Investissements d'Avenir–A\*MIDEX)). Images were generously provided by Sheri Amsel ([www.exploringnature.org](http://www.exploringnature.org)), Christopher Crocker and David Hall ([www.wormatlas.org](http://www.wormatlas.org)) and Andrew Leach (<http://andrewleachprojects.com>). We thank Rachel McMullan for sharing unpublished results.

## Figure Legends

Figure 1. A model for activation of effector secretion by LPS in horseshoe crab. Binding of LPS to factor C activates its proteolytic domain leading to cleavage of the putative GPCR and triggering of downstream signaling. In addition to IP<sub>3</sub>-dependent exocytosis, there is the potential for DAG-dependent activation of p38 signaling via a PKC. Secretion of the host defense molecule tachyplesin has been proposed to potentiate heterotrimeric G protein signaling, providing a positive feedback mechanism. Figure adapted from [9].

Figure 2. Simplified model of the regulation of p38 signaling by the GPCR DCAR-1 in the epidermis of *C. elegans*. Fungal infection or physical injury leads to the production of hydrophenyllactic acid (HPLA) that activates DCAR-1 and downstream elements. The name of each nematode protein and its vertebrate orthologue is given. Figure adapted from [28].

Figure 3. Model of DUOX activation in *Drosophila*. Uracil produced by pathogenic bacteria activates an as yet unidentified GPCR. This triggers calcium release from the endoplasmic reticulum via a PLC, and subsequent ROS production by Duox. Infection also drives expression of Duox via PLC-dependent activation of p38 MAPK.

## References

- [1] T. Kawai, S. Akira, The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors, *Nat Immunol* 11(5) (2010) 373-84.
- [2] M. Heil, W.G. Land, Danger signals - damaged-self recognition across the tree of life, *Front Plant Sci* 5 (2014) 578.
- [3] A.J. Kooistra, R. Leurs, I.J. de Esch, C. de Graaf, From three-dimensional GPCR structure to rational ligand discovery, *Adv Exp Med Biol* 796 (2014) 129-57.
- [4] F.B. Bang, A bacterial disease of *Limulus polyphemus*, *Bull Johns Hopkins Hosp* 98(5) (1956) 325-51.
- [5] T. Osaki, S. Kawabata, Structure and function of coagulogen, a clottable protein in horseshoe crabs, *Cell Mol Life Sci* 61(11) (2004) 1257-65.
- [6] A. Bergner, V. Oganessyan, T. Muta, S. Iwanaga, D. Typke, R. Huber, W. Bode, Crystal structure of a coagulogen, the clotting protein from horseshoe crab: a structural homologue of nerve growth factor, *Embo J* 15(24) (1996) 6789-97.
- [7] X.W. Wang, N.S. Tan, B. Ho, J.L. Ding, Evidence for the ancient origin of the NF-kappaB/IkappaB cascade: its archaic role in pathogen infection and immunity, *Proc Natl Acad Sci U S A* 103(11) (2006) 4204-9.
- [8] Y. Kobayashi, T. Shiga, T. Shibata, M. Sako, K. Maenaka, T. Koshiba, H. Mizumura, T. Oda, S. Kawabata, The N-terminal Arg residue is essential for autocatalytic activation of a lipopolysaccharide-responsive protease zymogen, *J Biol Chem* 289(37) (2014) 25987-95.
- [9] S. Kurata, S. Ariki, S. Kawabata, Recognition of pathogens and activation of immune responses in *Drosophila* and horseshoe crab innate immunity, *Immunobiology* 211(4) (2006) 237-49.



- [10] C. Cunha, A. Carvalho, A. Esposito, F. Bistoni, L. Romani, DAMP signaling in fungal infections and diseases, *Front Immunol* 3 (2012) 286.
- [11] A. Ozaki, S. Ariki, S. Kawabata, An antimicrobial peptide tachyplesin acts as a secondary secretagogue and amplifies lipopolysaccharide-induced hemocyte exocytosis, *FEBS J* 272(15) (2005) 3863-71.
- [12] T. Nakamura, H. Furunaka, T. Miyata, F. Tokunaga, T. Muta, S. Iwanaga, M. Niwa, T. Takao, Y. Shimonishi, Tachyplesin, a class of antimicrobial peptide from the hemocytes of the horseshoe crab (*Tachyplesus tridentatus*). Isolation and chemical structure, *J Biol Chem* 263(32) (1988) 16709-13.
- [13] J.C. Sullivan, F.S. Wolenski, A.M. Reitzel, C.E. French, N. Traylor-Knowles, T.D. Gilmore, J.R. Finnerty, Two alleles of NF-kappaB in the sea anemone *Nematostella vectensis* are widely dispersed in nature and encode proteins with distinct activities, *PLoS One* 4(10) (2009) e7311.
- [14] N. Pujol, E.M. Link, L.X. Liu, C.L. Kurz, G. Alloing, M.W. Tan, K.P. Ray, R. Solari, C.D. Johnson, J.J. Ewbank, A reverse genetic analysis of components of the Toll signalling pathway in *Caenorhabditis elegans*, *Curr Biol* 11(11) (2001) 809-21.
- [15] J.J. Ewbank, N. Pujol, Local and long-range activation of innate immunity by infection and damage in *C. elegans*, *Curr Opin Immunol* 38 (2016) 1-7.
- [16] J.P. Brandt, N. Ringstad, Toll-like Receptor Signaling Promotes Development and Function of Sensory Neurons Required for a *C. elegans* Pathogen-Avoidance Behavior, *Curr Biol* 25(17) (2015) 2228-37.
- [17] Q. Li, S.D. Liberles, Aversion and attraction through olfaction, *Curr Biol* 25(3) (2015) R120-9.
- [18] B.B. Shtonda, L. Avery, Dietary choice behavior in *Caenorhabditis elegans*, *J Exp Biol* 209(Pt 1) (2006) 89-102.

- [19] E. Pradel, Y. Zhang, N. Pujol, T. Matsuyama, C.I. Bargmann, J.J. Ewbank, Detection and avoidance of a natural product from the pathogenic bacterium *Serratia marcescens* by *Caenorhabditis elegans*, Proc Natl Acad Sci U S A 104(7) (2007) 2295-300.
- [20] C.L. Kurz, S. Chauvet, E. Andres, M. Aurouze, I. Vallet, G.P. Michel, M. Uh, J. Celli, A. Filloux, S. De Bentzmann, I. Steinmetz, J.A. Hoffmann, B.B. Finlay, J.P. Gorvel, D. Ferrandon, J.J. Ewbank, Virulence factors of the human opportunistic pathogen *Serratia marcescens* identified by *in vivo* screening, Embo J 22(7) (2003) 1451-1460.
- [21] E.R. Troemel, B.E. Kimmel, C.I. Bargmann, Reprogramming chemotaxis responses: sensory neurons define olfactory preferences in *C. elegans*, Cell 91(2) (1997) 161-9.
- [22] H.S. Fukuto, D.M. Ferkey, A.J. Apicella, H. Lans, T. Sharmeen, W. Chen, R.J. Lefkowitz, G. Jansen, W.R. Schafer, A.C. Hart, G protein-coupled receptor kinase function is essential for chemosensation in *C. elegans*, Neuron 42(4) (2004) 581-93.
- [23] J.D. Meisel, O. Panda, P. Mahanti, F.C. Schroeder, D.H. Kim, Chemosensation of bacterial secondary metabolites modulates neuroendocrine signaling and behavior of *C. elegans*, Cell 159(2) (2014) 267-80.
- [24] Y. Zhang, H. Lu, C.I. Bargmann, Pathogenic bacteria induce aversive olfactory learning in *Caenorhabditis elegans*, Nature 438(7065) (2005) 179-84.
- [25] G. Harris, Y. Shen, H. Ha, A. Donato, S. Wallis, X. Zhang, Y. Zhang, Dissecting the signaling mechanisms underlying recognition and preference of food odors, J Neurosci 34(28) (2014) 9389-403.

- [26] Y. Yu, L. Zhi, X. Guan, D. Wang, D. Wang, FLP-4 neuropeptide and its receptor in a neuronal circuit regulate preference choice through functions of ASH-2 trithorax complex in *Caenorhabditis elegans*, *Sci Rep* 6 (2016) 21485.
- [27] T. Kawli, F. He, M.W. Tan, It takes nerves to fight infections: insights on neuro-immune interactions from *C. elegans*, *Dis Model Mech* 3(11-12) (2010) 721-31.
- [28] D.H. Kim, J.J. Ewbank, Signaling in the innate immune response, in: *T.C.e.R.C.* ed (Ed.) *WormBook*, <http://www.wormbook.org>, 2015, pp. 1-51.
- [29] E.C. Andersen, J.S. Bloom, J.P. Gerke, L. Kruglyak, A variant in the neuropeptide receptor *npr-1* is a major determinant of *Caenorhabditis elegans* growth and physiology, *PLoS Genet* 10(2) (2014) e1004156.
- [30] M.A. Carrillo, M.L. Guillermin, S. Rengarajan, R.P. Okubo, E.A. Hallem, O<sub>2</sub>-sensing neurons control CO<sub>2</sub> response in *C. elegans*, *J Neurosci* 33(23) (2013) 9675-83.
- [31] E.V. Miller, L.N. Grandi, J.A. Giannini, J.D. Robinson, J.R. Powell, The Conserved G-Protein Coupled Receptor FSHR-1 Regulates Protective Host Responses to Infection and Oxidative Stress, *PLoS One* 10(9) (2015) e0137403.
- [32] J.D. Robinson, J.R. Powell, Long-term recovery from acute cold shock in *Caenorhabditis elegans*, *BMC Cell Biol* 17(1) (2016) 2.
- [33] J.R. Powell, D.H. Kim, F.M. Ausubel, The G protein-coupled receptor FSHR-1 is required for the *Caenorhabditis elegans* innate immune response, *Proc Natl Acad Sci U S A* 106(8) (2009) 2782-7.
- [34] K. Inamori, S. Aiki, S. Kawabata, A Toll-like receptor in horseshoe crabs, *Immunol Rev* 198 (2004) 106-15.
- [35] B. Lemaitre, J. Hoffmann, The Host Defense of *Drosophila melanogaster*, *Annu Rev Immunol* (2007).

- [36] R. Aoki, T. Yagami, H. Sasakura, K. Ogura, Y. Kajihara, M. Ibi, T. Miyamae, F. Nakamura, T. Asakura, Y. Kanai, Y. Misu, Y. Iino, M. Ezcurra, W.R. Schafer, I. Mori, Y. Goshima, A seven-transmembrane receptor that mediates avoidance response to dihydrocaffeic acid, a water-soluble repellent in *Caenorhabditis elegans*, J Neurosci 31(46) (2011) 16603-10.
- [37] O. Zugasti, N. Bose, B. Squiban, J. Belougne, C.L. Kurz, F.C. Schroeder, N. Pujol, J.J. Ewbank, Activation of a G protein-coupled receptor by its endogenous ligand triggers the innate immune response of *Caenorhabditis elegans*, Nat Immunol 15(9) (2014) 833-8.
- [38] K. Ziegler, C.L. Kurz, S. Cypowyj, C. Couillault, M. Pophillat, N. Pujol, J.J. Ewbank, Antifungal innate immunity in *C. elegans*: PKCdelta links G protein signaling and a conserved p38 MAPK cascade, Cell Host Microbe 5(4) (2009) 341-52.
- [39] N. Pujol, O. Zugasti, D. Wong, C. Couillault, C.L. Kurz, H. Schulenburg, J.J. Ewbank, Anti-fungal innate immunity in *C. elegans* is enhanced by evolutionary diversification of antimicrobial peptides, PLoS Pathog 4(7) (2008) e1000105.
- [40] N. Pujol, S. Cypowyj, K. Ziegler, A. Millet, A. Astrain, A. Goncharov, Y. Jin, A.D. Chisholm, J.J. Ewbank, Distinct innate immune responses to infection and wounding in the *C. elegans* epidermis, Curr Biol 18(7) (2008) 481-9.
- [41] K. Dierking, J. Polanowska, S. Omi, I. Engelmann, M. Gut, F. Lembo, J.J. Ewbank, N. Pujol, Unusual regulation of a STAT protein by an SLC6 family transporter in *C. elegans* epidermal innate immunity, Cell Host Microbe 9(5) (2011) 425-35.

- [42] T. Kawli, C. Wu, M.W. Tan, Systemic and cell intrinsic roles of Gqalpha signaling in the regulation of innate immunity, oxidative stress, and longevity in *Caenorhabditis elegans*, *P Natl Acad Sci USA* 107(31) (2010) 13788-93.
- [43] J. Hodgkin, P.E. Kuwabara, B. Corneliussen, A novel bacterial pathogen, *Microbacterium nematophilum*, induces morphological change in the nematode *C. elegans*, *Curr Biol* 10(24) (2000) 1615-1618.
- [44] A. Anderson, R. McMullan, G-proteins: Fighting infection on two fronts, *Worm* 1(4) (2012) 196-201.
- [45] R. McMullan, A. Anderson, S. Nurrish, Behavioral and Immune Responses to Infection Require Galphaq- RhoA Signaling in *C. elegans*, *PLoS Pathog* 8(2) (2012) e1002530.
- [46] A. Anderson, H. Laurenson-Schafer, F.A. Partridge, J. Hodgkin, R. McMullan, Serotonergic chemosensory neurons modify the *C. elegans* immune response by regulating G-protein signaling in epithelial cells, *PLoS Pathog* 9(12) (2013) e1003787.
- [47] Y. Qin, X. Zhang, Y. Zhang, A neuronal signaling pathway of CaMKII and Gqalpha regulates experience-dependent transcription of *tph-1*, *J Neurosci* 33(3) (2013) 925-35.
- [48] V. Singh, A. Aballay, Endoplasmic reticulum stress pathway required for immune homeostasis is neurally controlled by arrestin-1, *J Biol Chem* 287(40) (2012) 33191-7.
- [49] B. Squiban, J. Belougne, J. Ewbank, O. Zugasti, Quantitative and automated high-throughput genome-wide RNAi screens in *C. elegans*, *J Vis Exp* 60 (2012) e3448.

- [50] C. Bergamasco, P. Bazzicalupo, Chemical sensitivity in *Caenorhabditis elegans*, Cell Mol Life Sci 63(13) (2006) 1510-22.
- [51] C.I. Bargmann, Chemosensation in *C. elegans*, WormBook, The *C. elegans* Research Community ed, <http://www.wormbook.org>, 2006, pp. 1-29.
- [52] H.M. Robertson, J.H. Thomas, The putative chemoreceptor families of *C. elegans*, in: T.C.e.R. Community (Ed.) WormBook, WormBook, 2006.
- [53] J.H. Thomas, J.L. Kelley, H.M. Robertson, K. Ly, W.J. Swanson, Adaptive evolution in the SRZ chemoreceptor families of *Caenorhabditis elegans* and *Caenorhabditis briggsae*, Proc Natl Acad Sci U S A 102(12) (2005) 4476-81.
- [54] M.K. Stewart, N.L. Clark, G. Merrihew, E.M. Galloway, J.H. Thomas, High genetic diversity in the chemoreceptor superfamily of *Caenorhabditis elegans*, Genetics 169(4) (2005) 1985-96.
- [55] E.M. Ha, K.A. Lee, Y.Y. Seo, S.H. Kim, J.H. Lim, B.H. Oh, J. Kim, W.J. Lee, Coordination of multiple dual oxidase-regulatory pathways in responses to commensal and infectious microbes in drosophila gut, Nat Immunol 10(9) (2009) 949-57.
- [56] K.A. Lee, S.H. Kim, E.K. Kim, E.M. Ha, H. You, B. Kim, M.J. Kim, Y. Kwon, J.H. Ryu, W.J. Lee, Bacterial-derived uracil as a modulator of mucosal immunity and gut-microbe homeostasis in *Drosophila*, Cell 153(4) (2013) 797-811.
- [57] S. Valanne, H. Myllymaki, J. Kallio, M.R. Schmid, A. Kleino, A. Murumagi, L. Airaksinen, T. Kotipelto, M. Kaustio, J. Ulvila, S.S. Esfahani, Y. Engstrom, O. Silvennoinen, D. Hultmark, M. Parikka, M. Ramet, Genome-wide RNA interference in *Drosophila* cells identifies G protein-coupled receptor kinase 2 as a conserved regulator of NF-kappaB signaling, J Immunol 184(11) (2010) 6188-98.

- [58] C. Molnar, H. Holguin, F. Mayor, Jr., A. Ruiz-Gomez, J.F. de Celis, The G protein-coupled receptor regulatory kinase GPRK2 participates in Hedgehog signaling in *Drosophila*, *Proc Natl Acad Sci U S A* 104(19) (2007) 7963-8.
- [59] C. Molnar, A. Ruiz-Gomez, M. Martin, S. Rojo-Berciano, F. Mayor, J.F. de Celis, Role of the *Drosophila* non-visual ss-arrestin kurtz in hedgehog signalling, *PLoS Genet* 7(3) (2011) e1001335.
- [60] K. Pal, S.H. Hwang, B. Somatilaka, H. Badgandi, P.K. Jackson, K. DeFea, S. Mukhopadhyay, Smoothed determines beta-arrestin-mediated removal of the G protein-coupled receptor Gpr161 from the primary cilium, *J Cell Biol* 212(7) (2016) 861-75.
- [61] K.A. Lee, B. Kim, J. Bhin, H. Kim do, H. You, E.K. Kim, S.H. Kim, J.H. Ryu, D. Hwang, W.J. Lee, Bacterial uracil modulates *Drosophila* DUOX-dependent gut immunity via Hedgehog-induced signaling endosomes, *Cell Host Microbe* 17(2) (2015) 191-204.
- [62] K. Majzoub, M.L. Hafirassou, C. Meignin, A. Goto, S. Marzi, A. Fedorova, Y. Verdier, J. Vinh, J.A. Hoffmann, F. Martin, T.F. Baumert, C. Schuster, J.L. Immler, RACK1 controls IRES-mediated translation of viruses, *Cell* 159(5) (2014) 1086-95.
- [63] N.E. Beckage, D.B. Gelman, Wasp parasitoid disruption of host development: implications for new biologically based strategies for insect control, *Annu Rev Entomol* 49 (2004) 299-330.
- [64] K. Bitra, R.J. Suderman, M.R. Strand, Polydnavirus Ank proteins bind NF-kappaB homodimers and inhibit processing of Relish, *PLoS Pathog* 8(5) (2012) e1002722.
- [65] G. Gueguen, M.E. Kalamarz, J. Ramroop, J. Uribe, S. Govind, Polydnaviral ankyrin proteins aid parasitic wasp survival by coordinate and selective inhibition of

hematopoietic and immune NF-kappa B signaling in insect hosts, PLoS Pathog 9(8)  
(2013) e1003580.

[66] T.H. Cooper, K. Bailey-Hill, W.R. Leifert, E.J. McMurchie, S. Asgari, R.V. Glatz, Identification of an in vitro interaction between an insect immune suppressor protein (CrV2) and G alpha proteins, J Biol Chem 286(12) (2011) 10466-75.


[67] C. Dong, P. Zhang, A putative G protein-coupled receptor involved in innate immune defense of *Procambarus clarkii* against bacterial infection, Comp Biochem Physiol A Mol Integr Physiol 161(2) (2012) 95-101.

[68] X. Lin, I. Soderhall, Crustacean hematopoiesis and the astakine cytokines, Blood 117(24) (2011) 6417-24.

[69] J.J. Sun, J.F. Lan, X.Z. Shi, M.C. Yang, G.J. Niu, D. Ding, X.F. Zhao, X.Q. Yu, J.X. Wang, Beta-Arrestins negatively regulate the Toll pathway in shrimp by preventing Dorsal translocation and inhibiting Dorsal transcriptional activity, J Biol Chem (2016).



Figure 1



**Figure 2**  
[Click here to download high resolution image](#)

**Figure 2**






Figure 3





From Wikimedia Commons




Illustration by WormAtlas




Illustration by Andrew Leach