



HAL
open science

Neural bases of the bodily self as revealed by electrical brain stimulation: A systematic review

Zoé Dary, Bigna Lenggenhager, Stanislas Lagarde, Samuel Medina Villalon,
Fabrice Bartolomei, Christophe Lopez

► To cite this version:

Zoé Dary, Bigna Lenggenhager, Stanislas Lagarde, Samuel Medina Villalon, Fabrice Bartolomei, et al.. Neural bases of the bodily self as revealed by electrical brain stimulation: A systematic review. Human Brain Mapping, 2023, 1 (24), 10.1002/hbm.26253 . hal-04008492

HAL Id: hal-04008492

<https://amu.hal.science/hal-04008492>

Submitted on 10 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

REVIEW ARTICLE

Neural bases of the bodily self as revealed by electrical brain stimulation: A systematic review

Zoé Dary¹ | Bigna Lenggenhager^{2,3} | Stanislas Lagarde^{4,5} | Samuel Medina Villalon^{4,5} | Fabrice Bartolomei^{4,5} | Christophe Lopez¹

¹Aix Marseille University, CNRS, LNC, FR3C, Marseille, France

²Department of Psychology, University of Zurich, Zurich, Switzerland

³Department of Psychology, University of Konstanz, Konstanz, Germany

⁴Aix Marseille University, APHM, Inserm, INS, Timone Hospital, Marseille, France

⁵APHM, Timone Hospital, Epileptology and Cerebral Rhythmology, Marseille, France

Correspondence

Christophe Lopez, Laboratoire de Neurosciences Cognitives – UMR 7291, Aix Marseille University & Centre National de la Recherche Scientifique (CNRS), Centre Saint-Charles, Fédération de Recherche 3C – Case B. 3, Place Victor Hugo. 13331 Marseille Cedex 03, France.
Email: christophe.lopez@univ-amu.fr

Funding information

Agence Nationale de la Recherche, Grant/Award Number: ANR-19-CE37-0027

Abstract

An increasing amount of recent research has focused on the multisensory and neural bases of the bodily self. This pre-reflective form of self is considered as multifaceted, incorporating phenomenal components, such as self location, body ownership, first-person perspective, agency, and the perceptual body image. Direct electrical brain stimulation (EBS) during presurgical evaluation of epilepsy and brain tumor resection is a unique method to causally relate specific brain areas to the various phenomenal components of the bodily self. We conducted a systematic review of the literature describing altered phenomenal experience of the bodily self evoked by EBS. We included 42 articles and analyzed self reports from 221 patients. Three-dimensional density maps of EBS revealed that stimulation in the middle cingulum, inferior parietal lobule, supplementary motor area, posterior insula, hippocampal complex/amygdala, and precuneus most consistently altered one or several components of the bodily self. In addition, we found that only EBS in the parietal cortex induced disturbances of all five components of the bodily self considered in this review article. These findings inform current neuroscientific models of the bodily self.

KEYWORDS

self consciousness, bodily self, electrical brain stimulation, epilepsy, parietal cortex

1 | INTRODUCTION

The functionality, morphology, and state of our body have shown to play a major role in many cognitive, affective, and social processes and crucially define our self consciousness. Such bodily self is considered to result from a constant multisensory integration and fine-tuning of bottom-up signals (visual, somatosensory, interoceptive, and vestibular signals) and top-down expectancy (Blanke, 2012; Lenggenhager & Lopez, 2015; Park & Blanke, 2019; Tsakiris, 2010). Various phenomenal components of the bodily self have been described in the literature, of which five main components will be

differentiated in the present review article: self location, body ownership, the first-person perspective, the sense of agency, and the perceptual body image (Table 1). Self location refers to the experience of occupying a volume of space, typically localized within one's own physical body boundaries (Blanke & Metzinger, 2009). Body ownership is the experience of owning a physical body (Tsakiris et al., 2007). The first-person perspective refers to “the experience from where I perceive the world” (Blanke, 2012). The first-person perspective is usually centered on the body and corresponds to an egocentric perspective/viewpoint (Vogeley & Fink, 2003). Agency is the sense of being in control of one's own actions (Jeannerod, 2006). The

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Human Brain Mapping* published by Wiley Periodicals LLC.

TABLE 1 Five core phenomenal experiences underlying the bodily self

Bodily experiences	Definition	Exemplary experimental paradigm in healthy participants	Exemplary cases in neurology and psychiatry
Self location	A volume of space, normally localized within the physical boundaries of one's body	Rubber-hand illusion (Botvinick & Cohen, 1998), full-body illusion (Lenggenhager et al., 2007)	Out-of-body experience, heautoscopy
Body ownership	The feeling that our body and its parts belong to us	Rubber hand illusion, full-body illusion (Petkova et al., 2011), real hand illusion (Kannape et al., 2019)	Somatoparaphrenia, hemiasomatognosia, body integrity dysphoria
First-person perspective	The experience from where I perceive the world	Mental imagery perspective-taking tasks (Vogeley et al., 2004), full-body illusion (Ehrsson, 2007)	Heautoscopy, out-of-body experience
Agency	The feeling of being at the origin of our actions	Rubber hand illusion, full body illusion, arm agency illusions (Farrer et al., 2003), full-body agency illusions (Kannape & Blanke, 2013)	Schizophrenia, alien hand syndrome
Perceptual body image	Own body perceptions that can be verbalized: size, weight, body shape, etc.	Body shape distortions in the full-body illusion (Piryankova et al., 2014), rubber hand illusion (Linkenauger et al., 2013), Pinocchio illusion (Lackner, 1988)	Macro/microsomatognosia, supernumerary phantom limb, depersonalization, body integrity dysphoria, anorexia, Alice in Wonderland syndrome

Note: Definition of the bodily experiences, examples of experimental paradigms used to alter and study the respective component in healthy participants, and examples of neurological and psychiatric conditions leading to alterations in these bodily experiences.

perceptual body image refers to the perceptual experience of one's own body, such as the perceived size and shape of one's own body (Gallagher, 2005).

An increasing amount of research has investigated the neural mechanisms underlying the bodily self and its different phenomenal aspects (Blanke, 2012; Blanke et al., 2015). Such research includes investigations in neurological and psychiatric patients with specific alterations in the bodily self, as well as functional and structural neuroimaging in healthy participants (reviewed in Berlucchi & Aglioti, 2010; Blanke, 2012; Park & Blanke, 2019; see Table 1 for a summary). Results point to a widely distributed cerebral network, encompassing particularly the temporo-parietal, premotor, posterior parietal, and extrastriate cortices, underlying the stable sense of a bodily self. In neurological patients, for example, it has been proposed that lesion to the posterior insula and parietal operculum can lead to a sense of loss of ownership for one's own hand or arm (e.g., somatoparaphrenia; Gandola et al., 2012). Disembodied self location, on the other hand, has been linked to lesion or seizure involving the temporo-parietal junction (Blanke et al., 2004), or the posterior insula (Heydrich & Blanke, 2013). Lesions to the posterior parietal and the prefrontal cortex have been shown to distort own-body representations, for example the sense that the body is split in two (Heydrich et al., 2010), or the experience of having an additional limb (Hari et al., 1998).

Most studies in healthy participants combined functional neuroimaging with experimental manipulations of the bodily self through synchronous, but mismatching, multisensory stimulation, altering specific components of the bodily self (reviewed in Blanke, 2012; Dieguez & Lopez, 2017). Illusory ownership of a fake hand

(Botvinick & Cohen, 1998; Ehrsson et al., 2004), self identification with a virtual body and altered self location (Ionta et al., 2011; Lenggenhager et al., 2007; Nakul, Orlando-Dessaints, et al., 2020), altered sense of agency (Farrer et al., 2003), or distortions in the structure and size of the felt body (de Vignemont et al., 2005; Ehrsson, Kito, Sadato, Passingham & Naito, 2005) have been induced using such techniques in healthy participants (see Table 1 for a summary). Functional neuroimaging studies revealed an involvement of the premotor cortex, primary somatosensory cortex, extrastriate body area, insula, the putamen, and intraparietal sulcus in illusory self identification with a rubber hand or with a virtual body (Chancel et al., 2022; Ehrsson et al., 2004; Ehrsson, Holmes, & Passingham, 2005; Gentile et al., 2015; Guterstam, Björnsdotter, Bergouignan, et al., 2015; Ionta et al., 2011; Limanowski et al., 2014; Limanowski & Blankenburg, 2015; Petkova et al., 2011). Other studies found that the temporo-parietal junction, primary somatosensory cortex, premotor cortex, cingulate cortex, and the posterior superior temporal gyrus are related to illusory changes in the self location (Guterstam, Björnsdotter, Gentile, & Ehrsson, 2015; Ionta et al., 2011). The sense of agency on the other hand, has been related to the insula and inferior parietal cortex (Chambon et al., 2013; Farrer et al., 2003; Farrer, Frey, et al., 2008; Farrer, Bouchereau, et al., 2008; Farrer & Frith, 2002), whereas experimentally-induced changes in the perceptual body image (perceived shape and size or posture of the body) have been related to the postcentral sulcus, intraparietal sulcus, insula, and inferior parietal lobule (Ehrsson et al., 2005; Kavounoudias et al., 2008; Naito et al., 2017). While the neural networks revealed by these neuroimaging studies are typically based on a correlative approach, a more causal link has been suggested by non-invasive transcranial

direct current stimulation (de Boer et al., 2020; Lira et al., 2018; van Elk et al., 2017), or even more directly by invasive, intracranial electrical brain stimulation (EBS) (Arzy et al., 2006; Blanke et al., 2002; Bos et al., 2016; Desmurget et al., 2009; Schaller et al., 2021).

EBS is typically used during presurgical evaluation of focal intractable epilepsy to define as accurately as possible the seizure onset zone and perform functional brain mapping (Trebuchon et al., 2020; Trébuchon & Chauvel, 2016), or in awake patients during brain tumor resection (Duffau, 2015). During EBS an electrical current of several mA is directly delivered to the brain through intracerebrally implanted electrodes (stereoelectroencephalography, SEEG) or through subdural grids and strips of electrodes placed at the surface of the cerebral cortex (electrocorticography, ECoG; Grande et al., 2020; Isnard et al., 2018; Lesser et al., 2010). SEEG electrodes are implanted either unilaterally or bilaterally, and each electrode has multiple contacts to record activity from several brain regions and/or to apply EBS (Isnard et al., 2018). The electrodes allow to stimulate and record activity in deep brain regions such as the insular, cingulate, and orbitofrontal cortex, or the amygdala. During EBS, clinicians record the patients' phenomenal experience (including perceived disturbances of the bodily self), behavioral responses (e.g., muscle contraction), and electroencephalographic activity (i.e., SEEG and ECoG). During brain tumor resection, electrical current is applied through electrodes directly on the cortex or on subcortical fibers. Under local anesthesia, patients perform various cognitive and motor tasks to identify the resection boundaries, while largely preserving functionally significant areas and pathways. Since the pioneering work from neurosurgeon Wilder Penfield (e.g., Penfield & Rasmussen, 1950), clinicians have increasingly used EBS, providing functional maps (Duffau et al., 2003; Salanova et al., 1995a, 1995b) and connectivity maps (Duffau, 2015) of the human cerebral cortex. Given these advantages, EBS during awake surgery might be a particularly interesting perspective to assess the sense of self (Schaller et al., 2021).

EBS offers some advantages for mapping brain functions (reviewed in Mercier et al., 2022). EBS enables three-dimensional recordings of epileptic discharges, changes in cortical excitability and in brain connectivity evoked by EBS with a high temporal resolution (Isnard et al., 2018). Furthermore, EBS is usually considered to provide *causal* rather than just *correlative* evidence for a link between neural structures and phenomenal experience, such as somatosensory, visual, auditory, vestibular, emotional, and autonomic perceptions (reviewed in Selimbeyoglu & Parvizi, 2010), or more complex sensations, like reminiscence of past experience, *déjà-vu*, dreamy state and memory illusions (Curot et al., 2017; Penfield, 1955).

While in his seminal investigations Penfield did not investigate specifically disturbances of the bodily self, he reviewed 190 cases of surgery for focal epilepsy in awake patients under local anesthesia carried out during a nine-year period and described “psychical responses,” during which the patient is conscious and usually capable of introspection (Penfield, 1947, 1955). Psychical responses include experiential responses, defined as psychical hallucinations of past experience (e.g., flash back, dream), and interpretive responses, defined as psychical illusion about the present experience (e.g., *déjà-*

vu, fear, disembodiment). Interestingly, both experiential and interpretive responses encompass distortions of the bodily self as defined in the present study, as they refer to illusions about the experienced state of the body (posture, weight, size, shape) and self (location, connection to the body). Penfield (1955) identified 10 relevant cases and concluded that psychical responses “result from the stimulation principally of the lateral and superior surfaces of either temporal lobe. The superior surface is that portion of the temporal lobe that is hidden within the Sylvian fissure and in the circular sulcus that surrounds the insula beneath it” (see areas highlighted in pink in Figure 1). In line with earlier work by Penfield, more recent EBS studies confirmed the role of the superior temporal cortex and temporo-parietal junction in self location and embodiment. For example, an especially spectacular phenomenon reported during EBS in the temporo-parietal junction is an out-of-body experience (OBE), during which the self is perceived as located outside of the physical body, resulting in alterations of several components of the bodily self, especially self location and first-person perspective (Blanke et al., 2002; Bos et al., 2016; De Ridder et al., 2007; reviewed in Nakul & Lopez, 2017). However, full-blown OBEs seem very rare in the context of EBS, and investigations of the neural bases of other components of the bodily self have been poorly described in earlier EBS studies, have attracted less attention, and have not been systematically reviewed.




FIGURE 1 Brain areas responsible for psychical responses identified by Penfield. In this anatomical description of the left cerebral hemisphere, Penfield has represented the temporal lobe cut and turned down in order to show the superior and mesial surface of the temporal lobe. Modified from Penfield (1955) “The Twenty-Ninth Maudsley Lecture: The Role of the Temporal Cortex in Certain Psychical Phenomena” with the permission from Cambridge University Press

Here, we provide a systematic review of the literature on the neural bases of the bodily self as revealed by EBS in neurological patients, following the preferred reporting items for systematic reviews and meta-analyses (PRISMA) guidelines (Page et al., 2021). We focus on disturbances in the five core phenomenal components of the bodily self described above—self location, body ownership, first-person perspective, agency and perceptual body image—and aim at identifying the underlying brain areas.

2 | MATERIALS AND METHODS

2.1 | Search strategy

This systematic literature review is reported in accordance with the PRISMA guidelines (Page et al., 2021; Tables S1 and S2). We searched articles published in English in PubMed up to March 23, 2022 without date restriction using keywords in all available fields (title, abstract, keywords). The full search strategy is detailed in Table S3. Keywords searched included several synonyms for the stimulation and recording methods: “SEEG,” “stereo-EEG,” “electrical cortical stimulation,” “stereoencephalography,” “electrical stimulation,” “direct electrical stimulation,” “direct stimulation,” “stereo EEG,” “intracranial electroencephalography,” “stereo-electro-encephalography,” “epilepsy,” “awake,” “craniotomy,” “glioma.” For the bodily self disturbances, keywords included various disorders of self and own body perception: “embodiment,” “disembodiment,” “doppelgänger,” “feeling of a presence,” “heautoscopy,” “depersonalisation,” “self location,” “derealisation,” “autoscopy,” “bodily self,” “out of body experience,” “self consciousness,” “out-of-body illusion,” “body schema,” “body image,” “ownership,” “agency,” “vestibular,” “posture,” “proprioception.”

2.2 | Study selection

We included only primary articles (original research articles, case reports, prospective and retrospective case series) but not review articles. Inclusion criteria were studies reporting patients who received EBS with implanted depth electrodes (SEEG), subdural grids and strips of electrodes (EcoG), or during awake brain surgery. Studies were included in our review only if they reported a disturbance of one or several phenomenal components of the bodily self that could be related to the direct effect of EBS. Inclusion criteria were defined as follows:

- *Disturbances of self location* include (1) *OBE*, that is, full-blown illusory perception of being disembodied; (2) *vestibular sensations*, that is, illusory self motion without disembodiment, including illusory translations of the entire body, sensation of falling, floating in the air, rocking, and being tilted; and (3) *proprioceptive sensations*, that is, illusory motion of a body part without disembodiment, such as illusory elevation of an arm.

- *Disturbances of body ownership* correspond to the inability to recognize a body part as one's own, or to sensation of disownership, estrangement or disembodiment of a body part or of the whole body.
- *Disturbances of first-person perspective* correspond to a change in the multisensory perspective, or viewpoint on the world, which may be allocentric instead of egocentric, distanced and elevated.
- *Disturbances of agency* correspond to the sensations of not being at the origin of the bodily actions, of not controlling and generating them, or when one feels the urge to execute actions, or when one experiences a resistance to planned actions.
- *Disturbances of the perceptual body image* correspond to own body perceptions that can be verbalized, including sensations of lightness or heaviness of a body part (without illusory or real motion of this body part), sensations that a body part is absent or lost, or a perceived body size distortion.
- *Other disturbances of the bodily self*, which did not clearly fit into the above categories were also considered. This category encompasses the sense of depersonalization, that is, a dissociative experience combining sensations of being detached from the body, of losing control over the body, actions, or thoughts (Simeon & Abugel, 2006). This category also includes the feeling of a presence, the vivid sensation that somebody is present nearby (Fénelon et al., 2011), as it has been proposed that a sensed presence is a sensorimotor double of the patient's own body (Arzy et al., 2006; Bernasconi et al., 2021).

2.3 | Data extraction

The characteristics of the publications were extracted (authors, year of publication, region in the world), as well as information about the clinical population (sample size, number of patients reported, disease), methods of the applied EBS (brain areas explored, stimulation parameters). For each EBS-evoked bodily self disturbance, we identified the category of bodily self disturbance and the location of the stimulation (cerebral hemisphere, lobe, brain area, and coordinates of electrode location). Only four publications reported coordinates in a standardized atlas (14 cases from 2 studies in Talairach coordinates; 14 cases from 2 studies in MNI coordinates). Authors from recent studies were contacted by email to require coordinates of activation if they were not reported in their original publication. Six authors provided additional electrode coordinates (71 cases from 6 studies in MNI coordinates). All data are summarized in Table S4.

2.4 | Analysis of the neuroanatomical localization of EBS

The localizations of the EBS evoking different types of bodily self disturbances were summarized on a 3D rendering of the right cerebral hemisphere on the MNI brain template from FreeSurfer (Fischl et al., 2004). We used 3Dviewer tool for the visual




FIGURE 2 Flow diagram for the preferred reporting items for systematic reviews and meta-analyses (PRISMA)

representation (Medina Villalon et al., 2018) available at <https://meg.univ-amu.fr/wiki/3DViewer>. We converted the coordinates from Talairach to MNI coordinates when necessary, using custom-based scripts in MATLAB 2020 (The MathWorks, Inc., USA). When possible, we reported the other EBS localizations with the best approximation possible on the MNI template, considering the published MRI or implantation schema, and considering the original description of EBS localizations in terms of gyrus, sulcus, and Brodmann area. Sites of EBS were plotted on a single right cerebral hemisphere surface, by projecting coordinates on its closest mesial or lateral parts. Each point was projected on the surface, except for the amygdala, hippocampus, and insula, for which we created a mesh on which points were projected.

2.5 | Statistical analysis

To identify the core areas underpinning the bodily self, we calculated 3D spatial density of electrode contacts evoking disturbances of the bodily self using custom-based scripts in MATLAB 2020 and projected the spatial density maps on a 3D rendering of the right cerebral hemisphere on the MNI template. We also analyzed the proportion of evoked responses per brain region, as well as the hemispheric dominance of EBS for each component of the bodily self, using the Fisher's Exact test and z-tests (SPSS 26, IBM, USA).

3 | RESULTS

3.1 | Literature review

We identified 393 articles in the database, of which 20 were eligible to be included in the systematic review. We also searched for relevant studies in the references provided in each of these publications and included 22 other articles on the effects of EBS on the bodily self. In total, 42 articles were included in the systematic review. Figure 2 presents the PRISMA flow diagram of the study selection.

3.2 | Characteristics of the studies

The 42 articles included in the systematic review were published between 1937 and 2022. More than half of the studies were from Europe (52.3%; $n = 22$), 33.3% ($n = 14$) were conducted in North America, and 14.3% ($n = 6$) in Asia. Twenty-three articles were retrospective studies (sample size: 33–1132 patients; mean \pm SD: 196 ± 231 patients), 7 articles were case series (13–47 patients; 21 ± 16 patients), and 12 studies were single case reports. Table 2 summarizes the study characteristics, as well as clinical population, stimulation parameters, location of electrodes, and phenomenal experience.

TABLE 2 Summary of the publications included in the systematic review reporting bodily self disturbances evoked by electrical brain stimulation

Publication	Region	Clinical population and EBS procedure	Type of study: Sample size	Number of patients included in the review	Prevalence of bodily self disturbances (%)	Category of bodily self disturbance reported	Brain areas explored by EBS
Penfield and Boldrey (1937)	N. America	Epilepsy (ABS)	RS: 163	3	1.8	SL (vest), Ag.	Frontal, parietal, temporal
Penfield (1947)	N. America	Epilepsy (ABS)	RS: 190	1	0.5	SL (vest), Other	Temporal
Penfield (1955)	N. America	Epilepsy (ABS)	RS: 190	1	0.5	SL (OBE)	Temporal
Penfield (1957)	N. America	Brain tumor and epilepsy (ABS)	RS: 108	1	0.9	SL (vest)	Temporal, central
Mullan and Penfield (1959)	N. America	Epilepsy (ABS)	RS: 217	4	1.8	Other	Temporal
Penfield and Perot (1963)	N. America	Epilepsy (ABS)	RS: 1132	1	0.1	SL (vest)	Whole brain
Halgren et al. (1978)	N. America	Epilepsy (ABS)	CS: 36	15	–	SL (prop), Ag., SL (vest)	Temporal
Fried et al. (1991)	N. America	Epilepsy (subdural grid)	CS: 13	7	–	Ag., SL (prop)	Fronto-parietal
Richer et al. (1993)	N. America	Epilepsy (SEEG)	RS: 40	3	7.5	SL (prop), SL (vest)	Rolandic, parietal
Salanova et al. (1995a)	N. America	Epilepsy (ABS)	RS: 82	6	7.3	SL (vest), BI, Other	Parietal
Salanova et al. (1995b)	N. America	Epilepsy and brain tumor (ABS, subdural grid)	RS: 34	1	2.9	BI	Parietal
Blanke et al. (2000)	Europe	Epilepsy (subdural grid)	CR: 1	1	–	SL (vest)	Frontal, temporal, parietal
Kremer et al. (2001)	Europe	Epilepsy (SEEG)	CR: 1	1	–	Ag.	Cingulate
Blanke et al. (2002)	Europe	Epilepsy (subdural grid)	CR: 1	1	–	SL (vest), BI, 1PP, SL (OBE)	Frontal, temporal, parietal
Ostrowsky et al. (2002)	Europe	Epilepsy (SEEG)	CS: 30	2	–	BI	Temporal
Kahane et al. (2003)	Europe	Epilepsy (SEEG)	RS: 260	13	5	SL (vest)	Whole brain
Wiest et al. (2004)	Europe	Epilepsy (subdural grid)	CR: 1	1	–	SL (vest)	Parietal
So and Schaüble (2004)	N. America	Epilepsy (subdural grid)	CR: 1	1	–	Own.	Fronto-central
Isnard et al. (2004)	Europe	Epilepsy (SEEG)	RS: 50	12	24	SL (vest), Other	Temporal, insular
Arzy et al. (2006)	Europe	Epilepsy (subdural grid)	CR: 1	1	–	Other	Frontal, temporal, parietal
Vignal et al. (2007)	Europe	Epilepsy (SEEG)	RS: 180	3	1.7	Other, SL (OBE)	Temporal
De Ridder et al. (2007)	Europe	Tinnitus (paddle electrode)	CR: 1	1	–	SL (OBE)	Temporo-parietal
Mulak et al. (2008)	Europe	Epilepsy (SEEG)	RS: 339	5	1.5	Other	Whole brain
Desmurget et al. (2009)	Europe	Brain tumor (ABS)	CS: 7	3	–	Ag., SL (prop)	Parietal, premotor

TABLE 2 (Continued)

Publication	Region	Clinical population and EBS procedure	Type of study: Sample size	Number of patients included in the review	Prevalence of bodily self disturbances (%)	Category of bodily self disturbance reported	Brain areas explored by EBS
Mazzola et al. (2014)	Europe	Epilepsy (SEEG)	RS: 219	13	5.9	SL (vest)	Insula
Blanke et al. (2014)	Europe	Epilepsy, brain tumor (subdural grid)	CS: 5	3	–	Other	Insular, temporal, fronto-parietal
Balestrini et al. (2015)	Europe	Epilepsy (SEEG)	RS: 274	16	5.8	BI, Other, SL (vest)	Parietal
Bos et al. (2016)	Europe	Brain tumor (ABS)	CR: 1	1	–	SL (prop), SL (OBE), 1PP	Parietal, temporal, Sylvian fissure
Caruana et al. (2018)	Europe	Epilepsy (SEEG)	RS: 329	25	7.6	SL (vest)	Cingulate
Yu, Liu, et al. (2018)	Asia	Epilepsy (SEEG)	CR: 1	1	–	SL (OBE), 1PP	Frontal, temporal, parietal
Yu, Yu, et al. (2018)	Asia	Epilepsy (SEEG)	RS: 43	3	7	BI, SL (vest)	Operculo-insular
Popa et al. (2019)	Europe	Epilepsy (SEEG)	RS: 110	11	10	BI, SL (vest), SL (prop), 1PP	Cingulate
Mandonnet et al. (2020)	Europe	Brain tumor (ABS)	CR: 1	1	–	BI	Superior parietal
Fornia et al. (2020)	Europe	Brain tumor (ABS)	CS: 12	8	–	Ag.	Premotor and S1
Andelman-Gur et al. (2020)	Asia	Epilepsy (subdural grid)	RS: 62	23*	–	Ag., SL (vest), SL (OBE)	Frontal, parietal, occipital, temporal
Oane et al. (2020)	Europe	Epilepsy (SEEG)	CS: 47	6	–	SL (vest)	Whole brain
Fox et al. (2020)	N. America	Epilepsy (SEEG, subdural grid, strip of electrodes)	RS: 67	7	10.4	SL (vest)	Frontal, temporal, parietal, cingulate
Sun, Zhang, Ren, et al. (2021)	Asia	Epilepsy (SEEG)	RS: 20	1	–	BI	Parietal (S1)
Sun, Zhang, Yu, et al. (2021)	Asia	Epilepsy (SEEG)	RS: 33	9	27.3	Ag.	Parietal (S1)
Parvizi et al. (2021)	N. America	Epilepsy (SEEG)	CR: 1	1	–	Other	Frontal, temporal, parietal
Bratu et al. (2021)	Europe	Epilepsy (SEEG)	CR: 1	1	–	SL (OBE)	Whole brain
Hao et al. (2022)	Asia	Epilepsy (SEEG)	RS: 376	3	0.8	BI	Whole brain

Note: Publications are sorted by year of publication. Region of the sample: North America (N. America). Procedures for electrical brain stimulation (EBS): stimulation during awake brain surgery (ABS) for resection of brain tumor or epileptic zone, and stimulation during presurgical evaluation of epilepsy (stereoelectroencephalography [SEEG] or subdural electrodes), or implantation to treat tinnitus. Type of study: retrospective study (RS), case report (CR), and case series (CS). To avoid bias, the prevalence of bodily self disturbance (in % of the patients sample) is reported only when sample size is above 30. Categories of bodily self disturbance: SL (vest): self location (vestibular); SL (OBE): self location (out-of-body experience); SL (prop): self location (proprioceptive); Ag: agency; BI: body image; 1PP: first-person perspective; Own: ownership; Other (feeling of a presence, depersonalization, derealization). S1: primary somatosensory cortex. *Individual EBS evoking bodily self disturbances were reported as different cases as the number of patients was not indicated.

3.3 | Characteristics of clinical populations

Together, the studies included in this systematic review analyzed the EBS responses of a total of 4680 patients, of which 51.7% ($n = 2421$) received EBS during the presurgical evaluation of intractable epilepsy, 48.2% ($n = 2258$) received EBS during awake brain surgery for epilepsy or brain tumor, and 0.02% ($n = 1$) received EBS for intractable tinnitus. Out of these, we identified 221 patients (4.7% of all reported patients) who experienced a disturbance of one or several components of the bodily self during EBS. Eighty percent of the patients who reported a disturbance of their bodily self ($n = 176$) received EBS through SEEG, ECoG or strips of electrodes, whereas 20% ($n = 45$) received EBS during awake brain surgery.

3.4 | Prevalence of bodily self disturbances during EBS

When considering only the retrospective studies with a sample above 30 patients in order to avoid bias related to small samples (for similar procedures, see Dai et al., 2022), we observed that the prevalence of bodily self disturbances ranged from 0.1% to 27.3% (mean \pm SD: 6.2% \pm 7.3%; Table 2).

3.5 | Categories of bodily self disturbances evoked by EBS

Figure 3 compares the number of EBS-evoked disturbances of the different components of the bodily self and Table S4 summarizes the individual phenomenal reports.




FIGURE 3 Distribution of the EBS-evoked disturbances in the different components of the bodily self. Disturbances of self location are plotted separately for vestibular illusions, proprioceptive illusions, and out-of-body experiences (OBE). Note that the number of responses reported exceeds the number of patients: when patients reported disturbances in several components of the bodily self for a given EBS, all components were counted separately

The by far most commonly reported disturbance of the bodily self was a change in the perceived self location (163 responses representing 55.8% of all responses). Disturbances in the perceived self location were evoked significantly more often than any other component of the bodily self (all $z > 9.05$, and all p -values $< .0001$). The most common responses within this category included a perceived change in self location with vestibular sensations ($n = 131$; 44.9% of all responses; all $z > 6.55$ and $p < .0001$) characterized by illusory translation of the entire body (reported by 43 patients, e.g., Case 43 in Table S4: “Sliding towards the lower end of the bed”; Blanke et al., 2000), sensation of falling (in 39 patients, e.g., Case 45: “The patient reported that she was ‘sinking into the bed’ or ‘falling from a height’.”; Blanke et al., 2002), or feeling of floating in the air (in 31 patients, e.g., Case 34: “Feeling their whole body floating just above the bed”; Richer et al., 1993), without the sense of disembodiment.

The second most frequent EBS-evoked disturbance of the bodily self affected the sense of agency ($n = 57$; 19.5% of all responses). Disturbance in the sense of agency was also significantly more frequently evoked than other components of the bodily self, excluding self location (all $z > 2.13$ and all p -values $< .05$). Disorders of agency included the sensation an external agent moving (parts of) the body (reported by 17 patients, e.g., Cases 11–19: “The patients perceived all the above movements as being induced by an agent outside themselves”; Halgren et al., 1978), an “urge to move” for 8 patients (Andelman-Gur et al., 2020; Sun, Zhang, Yu, et al., 2021), an intention to move (5 patients, e.g., Case 85: “A pure intention, that is, a felt desire to move without any overt movement being produced”; Desmurget et al., 2009), or a resistance to perform an action (10 patients, e.g., Cases 171–180: “I feel resistance to anything I am told to do [...] the possibility of continuing in any action is blocked”; Andelman-Gur et al., 2020).

Disturbances of the perceptual body image represented 13% ($n = 38$) of the bodily self related responses. The cases included sensations that a body part was larger (in 15 patients, e.g., Cases 104–117: “He feels his right hand larger and swollen”; Balestrini et al., 2015), sensations that a body part was smaller (only in Case 45: “She reported seeing her legs ‘becoming shorter’.”; Blanke et al., 2002), or heavier (in 6 patients, e.g., Case 151: “Sensation that the upper right limb is heavier”; Popa et al., 2019), that a body part was missing (in 11 patients, e.g., Case 146: “The patient reported that he could not feel the existence of his right hand”; Yu, Yu, et al., 2018; Case 40: “Feeling that the leg is absent” (Salanova et al., 1995b), or the sensation that a body part was distorted (in 4 patients, e.g., Case 42: “My hand feels as if it is going around like a screwdriver”; Salanova et al., 1995a). There was no reported case of a supernumerary limb.

Other disturbances represented 9.9% ($n = 29$) of all bodily self related responses after EBS, a proportion of responses that did not differ from that of the perceptual body image ($z = 1.17$, $p = .24$). They included the sense of depersonalization (in 22 patients, e.g., Case 217: “This feeling of being disconnected from something. [...] It's like being weightless in your own mind as a personality”; Parvizi et al., 2021) and the feeling of a presence (in 5 patients, e.g., Case 75: “He is behind me, almost at my body, but I do not feel it”; Arzy et al., 2006).

There were 22 cases of disturbances of self location with proprioceptive illusion, which represented 7.5% of all responses (e.g., Case 45: “The patient felt her right leg being drawn towards the opposite wall of the operating theatre”; Blanke et al., 2002); Case 159: “Sensation that the upper part of the body moves upwards” (Popa et al., 2019).

We found only very rare cases of full-blown OBEs, which include changes in the first-person perspective and body ownership during EBS. We identified 10 cases of illusory self location during OBEs, representing 3.4% of all responses. OBE included the perception that the self was disembodied with autoscopia (e.g., Case 120: “She felt as if she floated just below the ceiling and saw her own body lying on the operating table”; Bos et al., 2016) or disembodiment without autoscopia (e.g., Case 5: “Oh God! I am leaving my body, an altered relationship to his own person as though he were outside of his body”; Penfield, 1955). We identified three responses of changes in the first-person perspective, associated with OBE, which represented 1% of all responses (e.g., Case 45: “I see myself lying in bed, from above, but I only see my legs and lower trunk”; Blanke et al., 2002). However, we found only one case of altered sense of body ownership during EBS in a patient with epilepsy (Case 62: “Sudden estrangement of the left lower extremity from the rest of his body”; So & Schaüble, 2004).

3.6 | Localization of EBS evoking disturbances of the different components of the bodily self

3.6.1 | Effect of EBS in the different lobes

Figure 4a summarizes the localization of EBS evoking disturbances of the bodily self and Figure 4b quantifies the different categories of

bodily self disturbances separately for each lobe and the cingulum. Disturbances of the bodily self were evoked by EBS in the parietal lobe ($n = 90$ responses evoked by EBS, representing 32.1% of all evoked responses in all brain regions stimulated), cingulum ($n = 60$, 21.4%), temporal lobe ($n = 54$, 19.3%), frontal lobe ($n = 39$, 13.9%) and insular lobe ($n = 32$, 11.4%). EBS in the occipital lobe only rarely evoked disturbances of the bodily self ($n = 5$, 1.8%). Statistical analyses indicated a significantly higher proportion of responses in the parietal lobe (32.1%) than in all other regions stimulated (all $z > 2.86$ and all p -values $< .01$). The proportion of responses did not differ between the cingulate and temporal cortex ($z = 0.63$, $p = .53$), but was higher in the cingulate than in the frontal cortex ($z = 2.32$, $p < .05$).

Figure 4b shows that only EBS in the parietal lobe evoked disturbances of all five phenomenal components underlying the ordinary and healthy sense of the bodily self that were considered in the present systematic review (including all three subcategories of self location, agency, perceptual body image, first-person perspective, body ownership).

3.6.2 | Localization of EBS changing the perceived self location

Vestibular sensations were evoked by EBS in the cingulate (36% of self location responses), parietal (27%), insular (16%) and temporal (15%) cortex (Figure 4b). Although the proportion of vestibular sensations did not differ significantly between the cingulate and parietal cortex ($z = 1.51$, $p = .12$), the proportion of vestibular responses in each of these areas was significantly higher than in the insular cortex (all $z = 2.01$ and $p < .05$) and temporal cortex (all $z = 2.36$ and $p < .05$). Stimulation in the anterior, middle and posterior cingulate cortex induced an illusory displacement of entire body. Similar sensations were evoked by insula stimulation, especially during EBS in the posterior insula. Illusory whole-body displacements were also evoked by temporal cortex stimulation, mostly in the hippocampal/amygdala complex (e.g., Cases 20–25), superior temporal gyrus, as well as less frequently by EBS in the middle and inferior temporal gyri. In the parietal cortex, EBS in the parietal operculum, angular gyrus or precuneus predominately evoked feelings of body elevation and illusions that the body moved toward one side (e.g., Case 54).

Proprioceptive sensations were evoked by EBS in the parietal (36% of the proprioceptive responses), temporal (32%), and cingulate (18%) cortex (statistical tests on proportions were not conducted due to the low number of responses). Responses including illusory translation of a body part (e.g., Case 159) occurred after stimulation of the right angular and supramarginal gyri, subcortical white matter in the temporo-parietal junction or middle cingulate cortex.

Ten OBEs were evoked during EBS in the temporal (50% of all OBE responses), parietal (20%), occipital (10%), frontal (10%) and insular (10%) cortex (statistical tests on proportions were not conducted due to the low number of responses). More precisely, OBE was evoked by EBS in the angular gyrus, the posterior part of the superior




FIGURE 4 Localization of EBS evoking bodily self disturbances. (a) EBS sites are summarized on 3D views of the right cerebral hemisphere, with a color code indicating the category of bodily self disturbances. The lower part shows EBS in the insula and mesio-temporal region. (b) Number of disturbances in the different components of the bodily self according to the brain region stimulated. 292 EBS could be located based on the anatomical data available in the articles considered in the review. Changes in the experience of self location are plotted separately for vestibular illusions, proprioceptive illusions and out-of-body experiences (OBE). Note that patients may report disturbances of several components of the bodily self for a given stimulation, which was coded in several categories

temporal gyrus at the junction with the angular gyrus, the subcortical white matter in the temporo-parietal junction, the anterior insula and in middle frontal gyrus and middle occipital gyrus.

3.6.3 | Localization of EBS changing the sense of agency

Fifty-six percent of the sensations to urge movements or sensations of a resistance to move were reported after EBS in the frontal cortex, whereas 26% of the responses were reported after parietal lobe stimulation, such as in the angular gyrus or somatosensory cortex. There was a significantly higher proportion of responses in the frontal cortex than in all other regions stimulated (all $z > 3.23$ all p -values $< .01$). Disturbances in the sense of agency were especially evoked by EBS in

the superior frontal gyrus (mesial and lateral parts) and precentral gyrus, including the supplementary motor area and the premotor cortex (e.g., Cases 171 and 172; “feeling a strong resistance to performing the task”). Stimulation in the mesial temporal lobe (hippocampus; 16% of the response) induced, for example, the sensation that the patient’s movements were “induced by an agent outside themselves” (e.g., Cases 11–19).

3.6.4 | Localization of EBS changing the perceptual body image

Disturbance of the perceptual body image was reported after EBS in the parietal (63% of all body image responses), cingulate (24%) and insular (8%) cortex. There was a significantly higher proportion of

responses in the parietal cortex than in all other regions stimulated (all $z > 3.47$ and p -values $< .001$). Sensation that a body part becomes heavier or lighter was induced, for example, after EBS in the angular gyrus or middle cingulate cortex (e.g., case 151). Illusory distortions of the body size (e.g., Case 45) occurred after EBS in the precuneus, postcentral gyrus, posterior cingulum, and superior parietal lobule. Finally, sensations that a body part is absent (e.g., Case 161) was reported when the superior part of the anterior insular long gyrus, middle cingulate cortex, superior parietal lobule and postcentral gyrus were stimulated.

3.6.5 | Localization of EBS changing the first-person perspective

Patients reported observing their environment from a viewpoint outside their body (together with a feeling of disembodiment, OBE) after stimulation in the parietal (25% of the responses), temporal (25%), cingulate (25%), and insular (25%) cortex (e.g., Cases 45, 120, 149, 153; statistical tests on proportions were not conducted due to the low number of responses). More precisely, changes in the first-person perspective were found after EBS in the angular gyrus, subcortical white matter in the temporo-parietal junction, anterior insula, and middle cingulate cortex. EBS at similar location evoked an OBE and a change in the first-person perspective, but OBE could also be evoked without a change in the first-person perspective.

3.6.6 | Localization of EBS changing the sense of body ownership

One case was identified during which EBS in the superior parietal lobule evoked an estrangement of the left lower extremity from the rest of the patient's body (Case 61).

3.6.7 | Localization of EBS evoking other disorders of the bodily self

Cases of depersonalization-derealization were reported during EBS in the superior temporal gyrus (12%), posterior cingulate cortex (4%), parahippocampal gyrus (19%), hippocampus (8%), insula (27%), and amygdala (4%); statistical tests on proportions were not conducted due to the low number of responses). The feeling of a presence was for example evoked by EBS in the left temporo-parietal junction (Case 75: "The patient had the impression that somebody was behind her").

3.7 | Identification of the main areas underpinning the bodily self

Two quantifications helped determine the main areas underpinning the bodily self. First, we plotted the histograms showing the number

of bodily self disturbances as a function of the main gyri and areas reported in the articles considered in our review. 280 EBS could be localized using this method (Figure 5). We found that disturbances of the bodily self were consistently evoked by EBS in the middle cingulum ($n = 50$, 18% of all responses evoked in all brain regions), inferior parietal lobule (angular and supramarginal gyrus, $n = 34$, 12%), mesial and lateral parts of the superior frontal gyrus (supplementary motor area and premotor cortex, $n = 23$, 8%), posterior insula ($n = 21$, 7.5%) and hippocampus ($n = 19$, 6.8%). EBS in these regions induced disturbances in one or more phenomenal components of the bodily self.

Second, we calculated 3D density maps of EBS for which coordinates were available, irrespective of the category of bodily self disturbance. Figure 6 revealed that six brain areas seem crucially involved in the bodily self as their stimulation most consistently altered the bodily self across the different studies. This approach revealed the predominant implication of the inferior parietal lobule, middle cingulum, supplementary motor area, posterior insula, hippocampal complex/amygdala, and the precuneus.

3.8 | Hemispheric laterality of EBS and category of bodily self disturbances

As not all information about the lateralization was available, we only listed the EBS where the lateralization was clearly indicated (203 responses; Figure 7). The statistical analysis of the effect of the stimulation side could only be carried out for self location, agency, body image and other sensations, for which there were enough cases reported. When considering these four experiences, we found a significant association between the hemisphere stimulated and the evoked responses ($\chi^2(3) = 55.84$, $p < .001$). The Bayes factor strongly supports the hypothesis for this association ($BF_3 = 1.23 \times 10^{11}$). Results of the z -tests indicated that for self location, the proportion of response after EBS in the right hemisphere was significantly higher than the proportion of response after EBS in the left hemisphere (77.6% of all responses evoked by EBS of the right hemisphere vs. 33.3% of all responses evoked by EBS of the left hemisphere, respectively). By contrast, for agency we found that the proportion of response in the left hemisphere was significantly higher than the proportion of response in the right hemisphere (32.1% vs. 0.8%, respectively). There was no statistically significant difference for the body image and other sensations.

4 | DISCUSSION

This systematic review of the literature presents the largest series of EBS in awake patients reporting disturbances of the bodily self. We describe below the nature of the phenomenal experience reported by the patients and the role of the stimulated brain areas in generating the sense of a bodily self.




FIGURE 5 Number of disturbances of the different components of the bodily self as a function of EBS location. 280 EBS could be localized according to the main gyri for each lobe and the cingulum

4.1 | Nature and frequency of distorted components of the bodily self during EBS

The present systematic review of the literature shows that not all components of the bodily self were equally altered by EBS in awake patients.

The high prevalence of illusory self location in space (vestibular illusions of floating, levitation or translation and proprioceptive illusions such as illusory movement of a body part) that do not involve the sense of disembodiment is likely related to the relatively simple and unisensory nature of these illusions. They are also commonly experienced during hypnagogic and hypnopompic states in




FIGURE 6 Color-coded density maps showing the number of EBS evoking disturbances of the bodily self. This analysis, conducted irrespective of the category of bodily self disturbances, revealed 6 main areas underlying the bodily self




FIGURE 7 Percentage of all reported responses according to the stimulation side. * Statistically significant differences ($p < .05$) for the z-tests comparing the number of responses evoked by the stimulation of the right vs. left cerebral hemisphere

neurotypical populations (reviewed in Windt, 2015). A high prevalence of illusory self location can also be accounted by the widespread cortical vestibular network, encompassing a large number of multisensory areas in the parietal, temporal, insular, and cingulate cortex (Dieterich et al., 2003; Kahane et al., 2003; Lopez, Blanke, & Mast, 2012; Zu Eulenburg et al., 2012), which are strongly and reciprocally interconnected (Raiser et al., 2020). Similarly, illusory self location with proprioceptive illusions are in line with a large frontoparietal network processing signals from muscle spindle afferents (Cignetti et al., 2014; Kavounoudias et al., 2008; Naito et al., 2017). In addition, there is evidence that the intrinsic cortical network architecture can predict the frequency of responses elicited by EBS in patients with epilepsy (Fox et al., 2020; McGonigal et al., 2021). EBS in “unimodal brain networks at the base of the cortical hierarchy elicited frequent and simple effects,” such as EBS in somatomotor and visual networks (Fox et al., 2020).

Abnormal sense of agency was the second most frequent type of EBS-evoked disturbances of the bodily self, whereas distortions of the perceptual body image and other disturbances were the third most frequent disturbances of the bodily self. One should note that this is in part related to the high prevalence of the reported “urge” to perform an action (Fried et al., 1991), which we included as disturbance of agency, defined in a non-restrictive way. Full-blown disturbance of agency, such as the feeling that someone else is controlling the action, an experience sometimes reported by patients with schizophrenia or depersonalization disorders, was however rarely found. Disturbances of the body image were mostly experiences that the body shape was distorted, usually involving the sense of a larger body part or a loss of a body part. Although we report of few cases of asomatognosia (missing limb, estrangement of body part) during EBS, the feeling of an additional body part seems to be rarely evoked by EBS. We found only one case of supernumerary phantom limb due to EBS of the motor cortex for treating central pain (Canavero et al., 1999), which did not follow our inclusion criteria. The patient reported once every 1–2 weeks the presence of a painful supernumerary left arm. However, this illusion was reported about 6 weeks after the stimulator implantation, and could not be causally related to the direct effect of EBS, when compared to the immediate effects of EBS in patients with epilepsy or during awake brain surgery. Consistent with our observations on the effects of EBS, supernumerary phantom limbs are also rare during seizures (Hari et al., 1998) or stroke (Khateb et al., 2009; Miyazawa et al., 2004). Other types of disturbances of the bodily self reported here were mainly related to depersonalization, which combines various sensations of being detached from the body, or losing control over the body, actions, or thoughts, or a feeling of a presence.

It is interesting to note that in our analysis OBE including changes in the first-person perspective and body ownership were very rarely found. While the rare cases have revealed important insights on the fundamentals of the bodily self (Blanke & Metzinger, 2009) and have been widely discussed in the scientific and general literature, there is no clear answer as to why these phenomena are so rare during EBS.

The low prevalence of OBE, changes in the first-person perspective and body ownership does not seem to be related to different excitability of the underlying intrinsic brain network (Fox et al., 2020), as these illusions were mostly triggered by EBS in the network where the most frequent disorders of the bodily self were also triggered. All these three disturbances of the bodily self are much more complex illusions than simple illusory elevation of the body, or illusory flexion of a body part, as they involve interpreting the experience from several senses together, and/or require a breakdown of a larger brain network. At a first glance, this low prevalence seems in contrast with the fact that 5%–10% of individuals from the general population have had at least an OBE in their lifetime, sometimes including changes in the visuospatial perspective (Blanke & Dieguez, 2009; Lopez & Elzière, 2018). On the other hand, experimentally-induced conflicts between senses in healthy participants have to date never evoked full-blown OBEs, but rather illusory self location toward an avatar and self identification with an avatar, without overt disembodiment (Ionta et al., 2011; Lenggenhager et al., 2007; Nakul, Orlando-Dessaints, et al., 2020). Alternatively, EBS may be effective only in populations prone to these illusory changes in embodiment, first-person perspective, and body ownership. Indeed, disturbances of self location and the perceptual body image, together with mental imagery abilities about the own body and spatial perspective, have been related to several factors and personality traits, including anxiety, migraine, depersonalization (Braithwaite et al., 2017; Lopez & Elzière, 2018), and schizotypy (Arzy et al., 2007; Mohr et al., 2006), or to different reliance on interoceptive and visual signals (Nakul, Dabard, et al., 2020; Pfeiffer et al., 2013; Tsakiris et al., 2011). It is therefore speculated that illusory disembodiment and changes in ownership were evoked mostly or only in these subpopulations of participants. Finally, it is also possible that the conditions to evoke such illusions require a particular involvement of the synchrony of the underlying neural systems (Hsu et al., 2022), as it is the case for complex phenomena such as memory reminiscence (Barbeau et al., 2005; Bartolomei et al., 2017).

4.2 | Prevalence of bodily self disturbances among experiential phenomena during EBS

Despite the increasing use of SEEG in clinical practice for presurgical evaluation of epilepsy (Parvizi & Kastner, 2018), there are only few retrospective studies in large samples of patients, and no prospective studies available, to correctly estimate the prevalence of EBS-evoked bodily self disturbances. A previous review article summarized the effects of EBS reported in the literature up 2010 (Selimbeyoglu & Parvizi, 2010). Based on their data, we calculated that out the 566 responses evoked after EBS, 9% were disturbances of the bodily self as defined in our present review article.

In the sample of patients included in our systematic review of the literature, we estimated that the prevalence of bodily self disturbances averaged 6.2% (of the patients receiving EBS) based on retrospective studies with a sample size above 30 patients. However, there was a very large variability in the prevalence across

studies, ranging from 0.1% to 27.3%, depending on the stimulated brain regions. The highest prevalence was found in the parietal cortex (primary somatosensory cortex: 27.3%; Sun, Zhang, Yu, et al., 2021) and temporo-insular cortex (24%; Isnard et al., 2004), whereas the lowest prevalence was found for early whole-brain or temporal cortex mapping by Penfield and colleagues (0.1%–1.8%; Penfield & Perot, 1963; Penfield, 1947, 1955, 1957).

The generally comparatively low occurrence of bodily self responses during EBS might have several causes. It has been argued that EBS is not ideal to investigate complex cognitive functions and perceptions due to the nature of the brain networks involved and because of methodological limitations pertaining to EBS procedures during SEEG and awake brain surgery (Herbet, 2021; Mandonnet, 2021). While low-level sensorimotor perceptions rely on rather local and unimodal neural networks, with limited spatial variability between brains, high-level cognitive and perceptual functions are based on more distributed networks of multisensory cortical and subcortical brain areas, which present with larger variability between individuals. There is recent evidence to suggest that EBS in network that are heteromodal and higher in the cortical hierarchy evoke rare and more variable perceptions (such as stimulation of the limbic and default mode network; Fox et al., 2020), which may render particularly difficult the study of higher cognitive functions. This may also account for the intra-individual and inter-individual variability of the responses during EBS in a same network (e.g., Halgren et al., 1978), sometimes given as a limitation of EBS procedures (Borchers et al., 2011). In addition to the nature of the networks involved, EBS procedures may limit the investigation of high-level cognitive functions and perceptions, such as distortions of the bodily self. Indeed, EBS is often applied during cognitive tasks such as reading, picture naming, memory tasks or motor tasks. Accordingly, the self reports and clinical responses collected are oriented toward the performance of these tasks, rather than toward detailed introspective analysis from the patients about their sense of self.

4.3 | Neural network underpinning the bodily self

The disturbances in the bodily self reported in the present review of the literature fall into the category of interpretive responses and psychical illusions defined by Penfield (1955), as they are illusions about the current state of the body and self. In contrast to Penfield's description of a core area for psychical responses in the temporal cortex (Figure 1), we identified a much larger network involved in the sense of self, encompassing mainly the inferior parietal lobule, middle cingulum, supplementary motor area, posterior insula, precuneus, and hippocampus (Figure 6). As reviewed in details elsewhere, all these areas belong to brain networks related to the sense of self in functional neuroimaging in healthy participants or in brain lesion studies (Berlucchi & Aglioti, 2010; Blanke, 2012; Blanke & Metzinger, 2009; Lenggenhager & Lopez, 2015; Park & Blanke, 2019; Serino et al., 2013; Seth, 2013; Tsakiris, 2010; Vogeley & Fink, 2003). These areas have also been related to phantom perceptions in general

(De Ridder et al., 2011). This might explain why EBS within areas belonging to this network can trigger various illusory contents about the self and body, either in relation to the perceptual and emotional state of the body (through cingulate, prefrontal, parietal, and precuneus activation), or to memories through activation of the hippocampus, parahippocampal area, and amygdala (De Ridder et al., 2011). Interestingly, the areas reported here as underpinning the bodily self overlap to a great degree with the posterior brain networks underlying conscious experience in general (Koch, 2018; Raccach et al., 2021). However, the bodily self, as a minimal and immediate form of self (Blanke & Metzinger, 2009; Gallagher, 2000), does not seem to involve the prefrontal cortex, which is more implicated in higher cognitive functions and reflective aspects of consciousness (Boly et al., 2017; Dehaene & Changeux, 2011; Odegaard et al., 2017) and is in general less likely to evoke responses and perceptions during EBS (Fox et al., 2020).

4.3.1 | Inferior parietal lobule

The present review revealed that only stimulation of the parietal cortex induced a modification of all components of bodily self and that the proportion of responses evoked by EBS in the parietal cortex was higher than in any other brain area stimulated. Except for disturbance of body ownership, which was only evoked once by EBS in the superior parietal lobule, EBS applied in the inferior parietal lobule resulted in a change in all other core components of the bodily self. Moreover, the analysis of EBS spatial density indicates a strong overlap of stimulation sites in the inferior parietal lobule (angular gyrus and supramarginal gyrus). This result from EBS studies showing causal implication of the inferior parietal lobule in the bodily self is in line with results from previous functional neuroimaging studies in healthy participants and brain lesion studies. Studies found that all areas within the inferior parietal lobule were strongly connected to the inferior frontal, insular and posterior temporal cortex, as well as to a broad network of other posterior brain regions, such as the somatosensory cortex and superior parietal areas, as well as with the auditory and visual cortex (Caspers et al., 2011). Results from diffusion tensor imaging and tractography studies (reviewed in Seghier, 2013) showed that the angular gyrus is especially strongly connected to areas that we identified as underpinning the bodily self, such as the precuneus, supramarginal gyrus, the hippocampus and parahippocampal gyrus, the middle temporal gyrus, the superior frontal gyrus, and the inferior frontal gyrus (Catani et al., 2005; Makris et al., 2007; Rushworth et al., 2006; Uddin et al., 2010). Because of this extensive pattern of connection, the inferior parietal lobule belongs to and interact with fronto-temporal systems, or the default mode network.

4.3.2 | Middle cingulate cortex

We found that EBS in the middle cingulate cortex evoked mostly sensations of illusory self location (vestibular illusions) and distortions of

the body image. The middle cingulate cortex has been involved in a very large range of functions, including “feedback processing, pain, salience, action-reward association, premotor functions, and conflict monitoring” in humans and in non-human primates (reviewed in Procyk et al., 2016; p. 467). More posterior stimulation during awake brain surgery have been related to altered states of consciousness (Herbet et al., 2016). A large retrospective analysis of the effects of EBS in the cingulate cortex indicates that the majority of vestibular responses—in relation to illusory self location in the present review article—were evoked by stimulation of the caudal part of the middle cingulate cortex (Caruana et al., 2018). In general, EBS in the posterior regions of the cingulate cortex (caudal part of the middle cingulate cortex and posterior cingulate cortex) were characterized by sensory illusions concerning the vestibular, interoceptive, somatosensory, and visual systems (Caruana et al., 2018). The role of the middle and posterior cingulate cortex in self location is also confirmed by their activation in fMRI studies that have used optokinetic stimulation or galvanic vestibular stimulation, evoking illusory self motion perception (Cardin & Smith, 2010; Smith et al., 2011), or experimentally-induced illusions of self location and body ownership (Guterstam, Björnsdotter, Gentile, & Ehrsson, 2015). Given the importance of these senses for the experienced bodily self, the posterior regions of the cingulate cortex may play an important role in the bodily self.

4.3.3 | Supplementary motor area

The supplementary motor area was mostly involved in changes in the sense of agency, as broadly defined in the present review article, although a few EBS also evoked illusory self location. The medial frontal cortex has consistently been involved in action monitoring in various fMRI studies (Yomogida et al., 2010), studies using transcranial magnetic stimulation and transcranial direct current stimulation (Cavazzana et al., 2015; Moore et al., 2010) and using SEEG recordings (Bonini et al., 2014). Here, we provide further causal evidence of the role of the mesial frontal cortex in the sense of agency.

4.3.4 | Insula

The insula has attracted a lot of attention recently in neuroscience. Functional MRI and PET studies have linked the insula to a large range of sensory, emotional and cognitive functions (Craig, 2002, 2009; Kurth, Zilles, et al., 2010; Kurth, Eickhoff, et al., 2010). The posterior insula has more specifically been involved in the processing of somatosensory, thermosensory, nociceptive, and vestibular information (Bottini et al., 1994, 2001; Dieterich et al., 2003; Mazzola et al., 2019; Ostrowsky et al., 2002). Recent meta-analyses of functional neuroimaging data revealed that the posterior insula is more particularly involved in sensorimotor processing, whereas the anterior insula is more involved in cognitive and socio-emotional functions (Kurth, Zilles, et al., 2010). Of note, the posterior insula contains

neurons responding to somatosensory stimuli to various body parts, and also to the entire body, making it a crucial area for whole-body integration and perception (Coq et al., 2004; Evrard, 2019; Schneider et al., 1993). The posterior insula also overlaps with the human equivalent of the monkey parieto-insular vestibular cortex, as revealed by functional neuroimaging and electrophysiological data (Bense et al., 2001; Bottini et al., 2001; Dieterich et al., 2003; Frank et al., 2014; Frank & Greenlee, 2018; Guldin & Grüsser, 1998; Lopez, Blanke, & Mast, 2012; Nakul et al., 2021). This area, which is considered the core of the vestibular cortical network, is involved in processing signals about self motion, and should therefore be important for the bodily self, given the recognized role of vestibular information for the neural underpinning of the bodily self (Lenggenhager & Lopez, 2015; Lopez, 2013, 2016; Lopez & Elzière, 2018; Lopez, Schreyer, et al., 2012). The fact that EBS evoked mostly changes in the bodily self during stimulation of the posterior rather than the anterior insula, is in line with the sensorimotor foundations of the bodily self, a minimal and immediate form of self (Blanke & Metzinger, 2009; Gallagher, 2000).

4.3.5 | Hippocampal complex

Stimulation of the hippocampus and parahippocampal gyrus evoked illusory self location (proprioceptive and vestibular), depersonalization, and distorted sense of agency. There is a large body of neurophysiological and neuroimaging evidence demonstrating that the hippocampus is crucially involved in self location and spatial memory, as it contains place cells coding the specific location of an animal (Barry & Burgess, 2014; Burgess & O'Keefe, 2003; O'Keefe & Conway, 1978; Poucet et al., 2003; Wiener et al., 2002) or a human (Ekstrom et al., 2003; Miller et al., 2013) within a real or virtual space. A seminal intracranial EEG study in patients with epilepsy identified place selectivity in the hippocampus in patients immersed in a virtual environment (Ekstrom et al., 2003). In addition, an MRI study during experimentally-induced OBE-like illusion teleporting the participants in another location of the room showed that activity in the hippocampus predicted the perceived location of the bodily self in the room (Guterstam, Björnsdotter, Gentile, & Ehrsson, 2015). Accordingly, EBS in the hippocampus may disturb the neural underpinnings of self location and/or disturb the encoding of sensory afferents to the hippocampus. There is indeed evidence that the hippocampus processes vestibular signals, which are crucial for self location (Hitier et al., 2021; Horii et al., 2004; O'Mara et al., 1994; Smith, 1997; Suzuki et al., 2001; Vitte et al., 1996). Moreover, the hippocampus is well-known for its role in episodic memory. EBS in the hippocampus has been shown to evoke various psychological experiences (Halgren et al., 1978; Penfield, 1958; Penfield & Perot, 1963), including episodic memories, personal semantics, familiarity and reminiscences of a dream (reviewed in Curot et al., 2017). This could explain the feelings of unreality reported by some patients during stimulation of the hippocampus.

4.3.6 | Precuneus

We found that stimulation of the precuneus also evoked disturbances of the body image, self location, and depersonalization. There is a large body of evidence indicating that the precuneus is involved in self processing, self awareness and consciousness (Cavanna & Trimble, 2006). Functional MRI and PET studies showed precuneus activations during numerous self related tasks, such as during visuo-spatial tasks conducted from a first- versus a third-person perspective, when participants attribute seen actions to self versus others, or judgment of personality traits pertaining to self versus others (David et al., 2006; Farrer & Frith, 2002; Kircher et al., 2000; Lambrey et al., 2012; Ruby & Decety, 2001; Vogeley et al., 2004). Interestingly, a PET study in a patient receiving EBS at the temporo-parietal junction, which triggered an OBE, revealed that the disembodied experience was related to activation of a brain network encompassing the right precuneus (De Ridder et al., 2007). The precuneus also receives vestibular information (Dieterich et al., 2003), which may explain the illusory self location reported during EBS. Tracer studies in animals indicate that the precuneus is interconnected with most of the brain areas that we found involved in the bodily self, including the supplementary motor area, the cingulate cortex, the inferior parietal lobule, and more generally the temporo-parieto-occipital cortex (Cavanna & Trimble, 2006). Accordingly, EBS in the precuneus may activate a large parieto-temporo-frontal network underlying self experience.

4.4 | Hemispheric dominance for the neural bases of the bodily self

Here, we found a right hemisphere dominance for self location and a left hemisphere dominance for the sense of agency, and no significant dominance for the perceptual body image and the other types of bodily self disturbances. The right dominance for self location is congruent with the dominance in the right cerebral hemisphere of vestibular information processing, which is crucial for the sense of self location, and represents the majority of the distortions of the bodily self collected in our systematic review. This right hemispheric dominance of the vestibular cortex has consistently been shown in right-handed individuals by a series of functional neuroimaging, anatomical and clinical studies (Dieterich et al., 2003; Dieterich et al., 2017; Janzen et al., 2008; Kirsch et al., 2016, 2018), as well as by a meta-analysis of functional neuroimaging data (Lopez, Blanke, & Mast, 2012). The right hemispheric dominance for self location is also in line with the right dominance of the proprioceptive networks (e.g., Naito et al., 2017). Overall, self awareness in a broad sense has long been associated with a right hemispheric dominance (Devinsky, 2000; Feinberg & Keenan, 2005; Keenan et al., 2005). Previous studies showed that the right hemisphere is specialized in own-body recognition and motor awareness (Antonello & Gottesman, 2017; Martinaud et al., 2017), as well as in first-person perspective (Vogeley et al., 2004) and the experience of an embodied self location (Ionta et al., 2011).

We found that the sense of agency was mostly disturbed when EBS was applied on the left side. While this result is in line with the large number of feeling of the “urge” to move and execute an action and the dominance of the motor areas in the left cerebral hemisphere of right-handed participants, it is not congruent with the implication of the mostly right insula and angular gyrus in the conscious sense of agency. However, these areas are more likely involved in the conscious sense of agency for ongoing actions (i.e., “monitoring intentional fluency” and “the subjective sense of control”; Haggard & Chambon, 2012) manipulated experimentally in fMRI studies, rather than in the feeling of urge to move evoked by EBS in otherwise immobile individuals.

4.5 | Limitations of EBS studies for neuroscientific investigations of the bodily self

Despite the recognized clinical advantages of EBS (George et al., 2020; Grande et al., 2020; Mandonnet et al., 2010), functional brain mapping using EBS has several limitations related to procedures and clinical constraints, which have been discussed in details elsewhere (Borchers et al., 2011; Parvizi & Kastner, 2018; Ritaccio et al., 2018).

First, EBS is limited by the unequal implantation of electrodes in the different cortices, which could be a bias for the results reported here. Most of focal epilepsies originate from the temporal and frontal lobes, which explains that the majority of electrodes was implanted in the temporal and frontal lobes (Parvizi & Kastner, 2018). In addition, there is an overrepresentation of some regions which have intensively been mapped and reported during the last years in retrospective studies, such as the effects of EBS in the insula (Mazzola et al., 2009, 2014, 2019; Ostrowsky et al., 2002; Yu, Yu, et al., 2018), cingulate cortex (Balestrini et al., 2015; Herbet et al., 2016; Oane et al., 2020; Popa et al., 2019), and frontal cortex (Fornia et al., 2020; Fox et al., 2018). We also note that because subcortical structures are only rarely implanted (with the exception of the hippocampus and amygdala) when compared to the cortex, we currently lack descriptions of the cerebellar and subcortical (brainstem, basal ganglia, thalamus) contributions to the bodily self, a spatial sampling bias which has been referred to as “corticocentric myopia” (Parvizi, 2009).

Second, EBS allows to investigate functions in various areas of the brain and create a causal link between neural activity and behavior (Borchers et al., 2011; Parvizi & Kastner, 2018) and is considered “the gold standard” for brain mapping (Mandonnet et al., 2010). However, distant and undirect effects of EBS have also been described and the causal effect of EBS is deemed controversial (Borchers et al., 2011; Desmurget et al., 2013). The effects of EBS are the result of inhibition and excitation of population of neurons at local and/or distant sites (Borchers et al., 2011; Ritaccio et al., 2018). Accordingly, locations of electrodes reported here may represent the effect of EBS on a larger brain network rather than the action on the neural populations close to the stimulation electrode only. There is indeed evidence that local stimulation, evoking for example ecstatic sensations, autobiographical

memories and own-body perceptions, were associated with increased or decreased pattern of functional connectivity within larger cerebral networks (Bartolomei et al., 2012, 2017, 2019; Popa et al., 2019).

Third, EBS is by definition applied to the brain of patients who have abnormal brain tissue functioning and/or tumors, which may hamper the generalization of findings to non-neurological individuals. Patients with long-lasting epilepsy exhibit changes in functional connectivity in epileptogenic networks (Besson et al., 2017; Bettus et al., 2008, 2009; Lagarde et al., 2022; Wirsich et al., 2016). We note that recent EBS studies report only the effects of stimulation in non-epileptic tissue and/or the effects of stimulation not followed by epileptic discharges. In the case of low-grade glioma, brain reorganization before EBS may have modified the structure and function of the brain networks investigated (Duffau, 2015). Of note, only 20% of the patients reported in the present review underwent awake brain surgery. The advantage of a systematic review like ours is to identify the brain areas and phenomenal responses that are consistent across studies and patients, despite differences in the stimulation methods and paradigms (frequency, duration, intensity...).

With respect to the study of the sense of self, another limitation is related to the way clinical assessments are done during EBS. Neuropsychological and sensorimotor assessments performed during EBS in patients with epilepsy or during awake brain surgery typically focus on language, memory and visuo-spatial abilities, whereas self and own-body perceptions may not be spontaneously reported by the patients and not recorded. Patients may also have difficulty expressing what they are feeling or be afraid to say what is happening to them, resulting in a reporting bias. Regarding spontaneous self reports, the lack of structured questionnaires or interview, rarely compatible with clinical routine during EBS, limits the detailed description that would be needed to fully describe the often complex phenomenal experience associated with changes in the sense of self. For example, structured questionnaires and scales have shown to be helpful for screening emotional responses during EBS in the amygdala (Lanteaume et al., 2007).

Finally, a limitation of systematic reviews of the literature is the efficiency of online search and citation searching in published articles to identify all published cases related to the sense of self. We note that our online search was not able to identify articles published before 1993, including the work by Penfield and colleagues, probably due to the lack of keywords and referenced abstract. The large number of keywords used to define the various facets of the bodily self in our review should have allowed a comprehensive identification of the cases of bodily self disturbances published until 2022.

5 | CONCLUSIONS AND PERSPECTIVES

This systematic review of the literature reveals that EBS can evoke a large variety of phenomenal content related to the bodily self, including a disturbed sense of self location (regarding the whole-body, body parts, or disembodiment) and agency, change in the first-person perspective, altered body ownership, distorted body image, or symptoms

of depersonalization. While these different phenomenal components of the bodily self are relatively rare and were not equally altered by EBS, they were all evoked by the stimulation of the parietal cortex. We also identified a network of six main areas that were most consistently involved in disturbances of the bodily self, including the inferior parietal lobule, middle cingulum, supplementary motor area, posterior insula, precuneus, and hippocampus. This seems in contrast with earlier work by Penfield (1955, 1947), who found that “psychical responses” (i.e., experiential and interpretive responses) were mostly evoked by EBS in the lateral and superior surfaces of the temporal lobes. Future electrophysiological studies should endeavor to determine how changes in neural network electrophysiology, such as dynamics of functional connectivity between distant brain areas, underpin the bodily self. The possibility to use sensory and cognitive tasks during EBS and SEEG recordings open interesting avenues, such as the simultaneous experimental manipulation of the bodily self (e.g., using the rubber hand illusion or the full-body illusion) and recordings of intracranial EEG signals (Guterstam et al., 2019).

ACKNOWLEDGMENTS

The authors are very thankful to the Drs. M. Andelman-Gur, S. Arzy, S. Balestrini, O. Blanke, E. Bos, F. Caruana, D. De Ridder, M. Desmurget, L. Fonia, K. Fox, F. Garbarini, L. Heydrich, L. Mazzola, I. Oane, J. Parvizi, S. Sunaert, G. Wiest, for kindly providing coordinates of electrodes or complementary information about their study. This work was supported by the ANR VESTISELF project, grant ANR-19-CE37-0027 of the French Agence Nationale de la Recherche to Christophe Lopez, Fabrice Bartolomei and Bigna Lenggenhager.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

Data used in this article are available in Table S4.

ORCID

Zoé Dary  <https://orcid.org/0000-0003-3243-3460>

Bigna Lenggenhager  <https://orcid.org/0000-0003-0418-9931>

Stanislas Lagarde  <https://orcid.org/0000-0003-2916-1302>

Fabrice Bartolomei  <https://orcid.org/0000-0002-1678-0297>

Christophe Lopez  <https://orcid.org/0000-0001-9298-2969>

REFERENCES

- Andelman-Gur, M. M., Gazit, T., Andelman, F., Kipervasser, S., Kramer, U., Neufeld, M. Y., Fried, I., & Fahoum, F. (2020). Spatial distribution and hemispheric asymmetry of electrically evoked experiential phenomena in the human brain. *Journal of Neurosurgery*, 133, 54–62. <https://doi.org/10.3171/2019.3.JNS183429>
- Antoniello, D., & Gottesman, R. (2017). Limb Misidentification: A Clinical-Anatomical Prospective Study. *The Journal of Neuropsychiatry and Clinical Neurosciences*, 29, 284–288. <https://doi.org/10.1176/appi.neuropsych.16090169>
- Arzy, S., Mohr, C., Michel, C. M., & Blanke, O. (2007). Duration and not strength of activation in temporo-parietal cortex positively correlates with schizotypy. *NeuroImage*, 35, 326–333.
- Arzy, S., Seeck, M., Ortigue, S., Spinelli, L., & Blanke, O. (2006). Induction of an illusory shadow person. *Nature*, 443, 287.
- Balestrini, S., Francione, S., Mai, R., Castana, L., Casaceli, G., Marino, D., Provinciali, L., Cardinale, F., & Tassi, L. (2015). Multimodal responses induced by cortical stimulation of the parietal lobe: A stereo-electroencephalography study. *Brain*, 138, 2596–2607. <https://doi.org/10.1093/brain/aww187>
- Barbeau, E., Wendling, F., Régis, J., Duncan, R., Poncet, M., Chauvel, P., & Bartolomei, F. (2005). Recollection of vivid memories after perirhinal region stimulations: synchronization in the theta range of spatially distributed brain areas. *Neuropsychologia*, 43, 1329–1337. <https://doi.org/10.1016/j.neuropsychologia.2004.11.025>
- Barry, C., & Burgess, N. (2014). Neural mechanisms of self-location. *Current Biology: CB*, 24, R330–R339. <https://doi.org/10.1016/j.cub.2014.02.049>
- Bartolomei, F., Barbeau, E. J., Nguyen, T., McGonigal, A., Régis, J., Chauvel, P., & Wendling, F. (2012). Rhinal-hippocampal interactions during déjà vu. *Clinical Neurophysiology*, 123, 489–495. <https://doi.org/10.1016/j.clinph.2011.08.012>
- Bartolomei, F., Lagarde, S., Médina Villalon, S., McGonigal, A., & Benar, C. G. (2017). The “Proust phenomenon”: Odor-evoked autobiographical memories triggered by direct amygdala stimulation in human. *Cortex*, 90, 173–175. <https://doi.org/10.1016/j.cortex.2016.12.005>
- Bartolomei, F., Lagarde, S., Scavarda, D., Carron, R., Bénar, C. G., & Picard, F. (2019). The role of the dorsal anterior insula in ecstatic sensation revealed by direct electrical brain stimulation. *Brain Stimulation*, 12, 1121–1126. <https://doi.org/10.1016/j.brs.2019.06.005>
- Bense, S., Stephan, T., Yousry, T. A., Brandt, T., & Dieterich, M. (2001). Multisensory cortical signal increases and decreases during vestibular galvanic stimulation (fMRI). *Journal of Neurophysiology*, 85, 886–899.
- Berlucchi, G., & Aglioti, S. M. (2010). The body in the brain revisited. *Experimental Brain Research*, 200, 25–35. <https://doi.org/10.1007/s00221-009-1970-7>
- Bernasconi, F., Blondiaux, E., Potheegadoo, J., Stripeikyte, G., Pagonabarraga, J., Bejr-Kasem, H., Bassolino, M., Akselrod, M., Martinez-Horta, S., Sampedro, F., Hara, M., Horvath, J., Franza, M., Konik, S., Bereau, M., Ghika, J.-A., Burkhard, P. R., Van De Ville, D., Faivre, N., ... Blanke, O. (2021). Robot-induced hallucinations in Parkinson's disease depend on altered sensorimotor processing in fronto-temporal network. *Science Translational Medicine*, 13, eabc8362. <https://doi.org/10.1126/scitranslmed.abc8362>
- Besson, P., Bandt, S. K., Proix, T., Lagarde, S., Jirsa, V. K., Ranjeva, J.-P., Bartolomei, F., & Guye, M. (2017). Anatomic consistencies across epilepsies: a stereotactic-EEG informed high-resolution structural connectivity study. *Brain*, 140, 2639–2652. <https://doi.org/10.1093/brain/awx181>
- Bettus, G., Guedj, E., Joyeux, F., Confort-Gouny, S., Soulier, E., Laguitton, V., Cozzone, P. J., Chauvel, P., Ranjeva, J.-P., Bartolomei, F., & Guye, M. (2009). Decreased basal fMRI functional connectivity in epileptogenic networks and contralateral compensatory mechanisms. *Human Brain Mapping*, 30, 1580–1591. <https://doi.org/10.1002/hbm.20625>
- Bettus, G., Wendling, F., Guye, M., Valton, L., Régis, J., Chauvel, P., & Bartolomei, F. (2008). Enhanced EEG functional connectivity in mesial temporal lobe epilepsy. *Epilepsy Research*, 81, 58–68. <https://doi.org/10.1016/j.eplepsyres.2008.04.020>
- Blanke, O. (2012). Multisensory brain mechanisms of bodily self-consciousness. *Nature Reviews. Neuroscience*, 13, 556–571. <https://doi.org/10.1038/nrn3292>
- Blanke, O., & Dieguez, S. (2009). Leaving body and life behind: Out-of-body and near-death experience. In S. Laureys & G. Tononi (Eds.), *The Neurology of Consciousness: Cognitive Neuroscience and Neuropathology* (pp. 303–325). Elsevier.
- Blanke, O., Landis, T., Spinelli, L., & Seeck, M. (2004). Out-of-body experience and autoscopia of neurological origin. *Brain*, 127, 243–258.

- Blanke, O., & Metzinger, T. (2009). Full-body illusions and minimal phenomenal selfhood. *Trends in Cognitive Sciences*, 13, 7–13.
- Blanke, O., Ortigue, S., Landis, T., & Seeck, M. (2002). Stimulating illusory own-body perceptions. *Nature*, 419, 269–270.
- Blanke, O., Perrig, S., Thut, G., Landis, T., & Seeck, M. (2000). Simple and complex vestibular responses induced by electrical cortical stimulation of the parietal cortex in humans. *Journal of Neurology, Neurosurgery, and Psychiatry*, 69, 553–556.
- Blanke, O., Pozeg, P., Hara, M., Heydrich, L., Serino, A., Yamamoto, A., Higuchi, T., Salomon, R., Seeck, M., Landis, T., Arzy, S., Herbelin, B., Bleuler, H., & Rognini, G. (2014). Neurological and robot-controlled induction of an apparition. *Current Biology*, 24, 2681–2686. <https://doi.org/10.1016/j.cub.2014.09.049>
- Blanke, O., Slater, M., & Serino, A. (2015). Behavioral, neural, and computational principles of bodily self-consciousness. *Neuron*, 88, 145–166. <https://doi.org/10.1016/j.neuron.2015.09.029>
- Boly, M., Massimini, M., Tsuchiya, N., Postle, B. R., Koch, C., & Tononi, G. (2017). Are the neural correlates of consciousness in the front or in the back of the cerebral cortex? Clinical and neuroimaging evidence. *The Journal of Neuroscience*, 37, 9603–9613. <https://doi.org/10.1523/JNEUROSCI.3218-16.2017>
- Bonini, F., Burle, B., Liégeois-Chauvel, C., Régis, J., Chauvel, P., & Vidal, F. (2014). Action monitoring and medial frontal cortex: leading role of supplementary motor area. *Science*, 343, 888–891. <https://doi.org/10.1126/science.1247412>
- Borchers, S., Himmelbach, M., Logothetis, N., & Karnath, H.-O. (2011). Direct electrical stimulation of human cortex—The gold standard for mapping brain functions? *Nature Reviews. Neuroscience*, 13, 63–70. <https://doi.org/10.1038/nrn3140>
- Bos, E. M., Spoor, J. K. H., Smits, M., Schouten, J. W., & Vincent, A. J. P. E. (2016). Out-of-body experience during awake craniotomy. *World Neurosurgery*, 92, 586.e9–586.e13. <https://doi.org/10.1016/j.wneu.2016.05.002>
- Bottini, G., Karnath, H. O., Vallar, G., Sterzi, R., Frith, C. D., Frackowiak, R. S., & Paulesu, E. (2001). Cerebral representations for egocentric space: Functional-anatomical evidence from caloric vestibular stimulation and neck vibration. *Brain*, 124, 1182–1196.
- Bottini, G., Sterzi, R., Paulesu, E., Vallar, G., Cappa, S. F., Erminio, F., Passingham, R. E., Frith, C. D., & Frackowiak, R. S. (1994). Identification of the central vestibular projections in man: a positron emission tomography activation study. *Experimental Brain Research*, 99, 164–169.
- Botvinick, M., & Cohen, J. (1998). Rubber hands “feel” touch that eyes see. *Nature*, 391, 756.
- Braithwaite, J. J., Watson, D. G., & Dewe, H. (2017). Predisposition to out-of-body experience (OBE) is associated with aberrations in multisensory integration: Psychophysiological support from a “rubber hand illusion” study. *Journal of Experimental Psychology. Human Perception and Performance*, 43, 1125–1143. <https://doi.org/10.1037/xhp0000406>
- Bratu, F.-I., Oane, I., Barborica, A., Donos, C., Pistol, C., Daneasa, A., Lentoiu, C., & Mindruta, I. (2021). Network of autoscopic hallucinations elicited by intracerebral stimulations of periventricular nodular heterotopia: An SEEG study. *Cortex*, 145, 285–294. <https://doi.org/10.1016/j.cortex.2021.08.018>
- Burgess, N., & O’Keefe, J. (2003). Neural representations in human spatial memory. *Trends in Cognitive Sciences*, 7, 517–519.
- Canavero, S., Bonicalzi, V., Castellano, G., Perozzo, P., & Massa-Micon, B. (1999). Painful supernumerary phantom arm following motor cortex stimulation for central poststroke pain: Case report. *Journal of Neurosurgery*, 91, 121–123. <https://doi.org/10.3171/jns.1999.91.1.0121>
- Cardin, V., & Smith, A. T. (2010). Sensitivity of human visual and vestibular cortical regions to egomotion-compatible visual stimulation. *Cerebral Cortex*, 20, 1964–1973. <https://doi.org/10.1093/cercor/bhp268>
- Caruana, F., Gerbella, M., Avanzini, P., Gozzo, F., Pelliccia, V., Mai, R., Abdollahi, R. O., Cardinale, F., Sartori, I., Lo Russo, G., & Rizzolatti, G. (2018). Motor and emotional behaviours elicited by electrical stimulation of the human cingulate cortex. *Brain*, 141(10), 3035–3051. <https://doi.org/10.1093/brain/awy219>
- Caspers, S., Eickhoff, S. B., Rick, T., von Kapri, A., Kuhlen, T., Huang, R., Shah, N. J., & Zilles, K. (2011). Probabilistic fibre tract analysis of cytoarchitecturally defined human inferior parietal lobule areas reveals similarities to macaques. *NeuroImage*, 58, 362–380. <https://doi.org/10.1016/j.neuroimage.2011.06.027>
- Catani, M., Jones, D. K., & Ffytche, D. H. (2005). Perisylvian language networks of the human brain. *Annals of Neurology*, 57, 8–16. <https://doi.org/10.1002/ana.20319>
- Cavanna, A. E., & Trimble, M. R. (2006). The precuneus: a review of its functional anatomy and behavioural correlates. *Brain*, 129, 564–583. <https://doi.org/10.1093/brain/awl004>
- Cavazzana, A., Penolazzi, B., Begliomini, C., & Bisiacchi, P. S. (2015). Neural underpinnings of the “agent brain”: new evidence from transcranial direct current stimulation. *The European Journal of Neuroscience*, 42, 1889–1894. <https://doi.org/10.1111/ejn.12937>
- Chambon, V., Wenke, D., Fleming, S. M., Prinz, W., & Haggard, P. (2013). An online neural substrate for sense of agency. *Cerebral Cortex*, 23, 1031–1037. <https://doi.org/10.1093/cercor/bhs059>
- Chancel, M., Iriye, H., & Ehrsson, H. H. (2022). Causal inference of body ownership in the posterior parietal cortex. *The Journal of Neuroscience*, 42, 7131–7143. <https://doi.org/10.1523/JNEUROSCI.0656-22.2022>
- Cignetti, F., Vaugoyeau, M., Nazarian, B., Roth, M., Anton, J.-L., & Assaiante, C. (2014). Boosted activation of right inferior frontoparietal network: a basis for illusory movement awareness. *Human Brain Mapping*, 35, 5166–5178. <https://doi.org/10.1002/hbm.22541>
- Coq, J.-O., Qi, H., Collins, C. E., & Kaas, J. H. (2004). Anatomical and functional organization of somatosensory areas of the lateral fissure of the New World titi monkey (*Callicebus moloch*). *The Journal of Comparative Neurology*, 476, 363–387. <https://doi.org/10.1002/cne.20237>
- Craig, A. D. (2002). How do you feel? Interoception: the sense of the physiological condition of the body. *Nature Reviews. Neuroscience*, 3, 655–666.
- Craig, A. D. (2009). How do you feel—now? The anterior insula and human awareness. *Nature Reviews. Neuroscience*, 10, 59–70. <https://doi.org/10.1038/nrn2555>
- Curot, J., Busigny, T., Valton, L., Denuelle, M., Vignal, J.-P., Maillard, L., Chauvel, P., Pariente, J., Trebuchon, A., Bartolomei, F., & Barbeau, E. J. (2017). Memory scrutinized through electrical brain stimulation: A review of 80 years of experiential phenomena. *Neuroscience and Biobehavioral Reviews*, 78, 161–177. <https://doi.org/10.1016/j.neubiorev.2017.04.018>
- Dai, S., Lemaire, C., Piscicelli, C., & Pérennou, D. (2022). Lateropulsion prevalence after stroke: a systematic review and meta-analysis. *Neurology*, 98, e1574–e1584. <https://doi.org/10.1212/WNL.000000000000200010>
- David, N., Bewernick, B. H., Cohen, M. X., Newen, A., Lux, S., Fink, G. R., Shah, N. J., & Vogeley, K. (2006). Neural representations of self versus other: visual-spatial perspective taking and agency in a virtual ball-tossing game. *Journal of Cognitive Neuroscience*, 18, 898–910.
- de Boer, D. M. L., Johnston, P. J., Kerr, G., Meinzer, M., & Cleeremans, A. (2020). A causal role for the right angular gyrus in self-location mediated perspective taking. *Scientific Reports*, 10, 19229. <https://doi.org/10.1038/s41598-020-76235-7>
- De Ridder, D., Elgoyhen, A. B., Romo, R., & Langguth, B. (2011). Phantom percepts: tinnitus and pain as persisting aversive memory networks. *Proceedings of the National Academy of Sciences of the United States of America*, 108, 8075–8080. <https://doi.org/10.1073/pnas.1018466108>
- De Ridder, D., Van Laere, K., Dupont, P., Menovsky, T., & Van de Heyning, P. (2007). Visualizing out-of-body experience in the brain. *The New England Journal of Medicine*, 357, 1829–1833. <https://doi.org/10.1056/NEJMoa070010>

- de Vignemont, F., Ehrsson, H. H., & Haggard, P. (2005). Bodily illusions modulate tactile perception. *Current Biology*, 15, 1286–1290. <https://doi.org/10.1016/j.cub.2005.06.067>
- Dehaene, S., & Changeux, J.-P. (2011). Experimental and theoretical approaches to conscious processing. *Neuron*, 70, 200–227. <https://doi.org/10.1016/j.neuron.2011.03.018>
- Desmurget, M., Reilly, K. T., Richard, N., Szathmari, A., Mottolese, C., & Sirigu, A. (2009). Movement intention after parietal cortex stimulation in humans. *Science*, 324, 811–813. <https://doi.org/10.1126/science.1169896>
- Desmurget, M., Song, Z., Mottolese, C., & Sirigu, A. (2013). Re-establishing the merits of electrical brain stimulation. *Trends in Cognitive Sciences*, 17, 442–449. <https://doi.org/10.1016/j.tics.2013.07.002>
- Devinsky, O. (2000). Right cerebral hemisphere dominance for a sense of corporeal and emotional self. *Epilepsy & Behavior*, 1, 60–73. <https://doi.org/10.1006/ebep.2000.0025>
- Dieguez, S., & Lopez, C. (2017). The bodily self: Insights from clinical and experimental research. *Annals of Physical and Rehabilitation Medicine*, 60, 198–207. <https://doi.org/10.1016/j.rehab.2016.04.007>
- Dieterich, M., Bense, S., Lutz, S., Drzegza, A., Stephan, T., Bartenstein, P., & Brandt, T. (2003). Dominance for vestibular cortical function in the non-dominant hemisphere. *Cerebral Cortex*, 13, 994–1007.
- Dieterich, M., Kirsch, V., & Brandt, T. (2017). Right-sided dominance of the bilateral vestibular system in the upper brainstem and thalamus. *Journal of Neurology*, 264, 55–62. <https://doi.org/10.1007/s00415-017-8453-8>
- Duffau, H. (2015). Stimulation mapping of white matter tracts to study brain functional connectivity. *Nature Reviews. Neurology*, 11, 255–265. <https://doi.org/10.1038/nrneuro.2015.51>
- Duffau, H., Capelle, L., Denvil, D., Sichez, N., Gatignol, P., Taillandier, L., Lopes, M., Mitchell, M.-C., Roche, S., Muller, J.-C., Bitar, A., Sichez, J.-P., & van Effenterre, R. (2003). Usefulness of intraoperative electrical subcortical mapping during surgery for low-grade gliomas located within eloquent brain regions: functional results in a consecutive series of 103 patients. *Journal of Neurosurgery*, 98, 764–778. <https://doi.org/10.3171/jns.2003.98.4.0764>
- Ehrsson, H. H. (2007). The experimental induction of out-of-body experiences. *Science*, 317, 1048.
- Ehrsson, H. H., Holmes, N. P., & Passingham, R. E. (2005). Touching a rubber hand: feeling of body ownership is associated with activity in multisensory brain areas. *The Journal of Neuroscience*, 25, 10564–10573.
- Ehrsson, H. H., Kito, T., Sadato, N., Passingham, R. E., & Naito, E. (2005). Neural substrate of body size: illusory feeling of shrinking of the waist. *PLoS Biology*, 3, e412. <https://doi.org/10.1371/journal.pbio.0030412>
- Ehrsson, H. H., Spence, C., & Passingham, R. E. (2004). That's my hand! Activity in premotor cortex reflects feeling of ownership of a limb. *Science*, 305, 875–877.
- Ekstrom, A. D., Kahana, M. J., Caplan, J. B., Fields, T. A., Isham, E. A., Newman, E. L., & Fried, I. (2003). Cellular networks underlying human spatial navigation. *Nature*, 425, 184–188. <https://doi.org/10.1038/nature01964>
- Evard, H. C. (2019). The organization of the primate insular cortex. *Frontiers in Neuroanatomy*, 13, 43. <https://doi.org/10.3389/fnana.2019.00043>
- Farrer, C., Bouchereau, M., Jeannerod, M., & Franck, N. (2008). Effect of distorted visual feedback on the sense of agency. *Behavioural Neurology*, 19, 53–57.
- Farrer, C., Franck, N., Georgieff, N., Frith, C. D., Decety, J., & Jeannerod, M. (2003). Modulating the experience of agency: A positron emission tomography study. *NeuroImage*, 18, 324–333.
- Farrer, C., Frey, S. H., Van Horn, J. D., Tunik, E., Turk, D., Inati, S., & Grafton, S. T. (2008). The angular gyrus computes action awareness representations. *Cerebral Cortex*, 18, 254–261.
- Farrer, C., & Frith, C. D. (2002). Experiencing oneself vs another person as being the cause of an action: the neural correlates of the experience of agency. *NeuroImage*, 15, 596–603.
- Feinberg, T. E., & Keenan, J. P. (2005). Where in the brain is the self? *Consciousness and Cognition*, 14, 661–678. <https://doi.org/10.1016/j.concog.2005.01.002>
- Fénelon, G., Soulas, T., De Langavant, L. C., Trinkler, I., & Bachoud-Lévi, A.-C. (2011). Feeling of presence in Parkinson's disease. *Journal of Neurology, Neurosurgery, and Psychiatry*, 82, 1219–1224. <https://doi.org/10.1136/jnnp.2010.234799>
- Fischl, B., van der Kouwe, A., Destrieux, C., Halgren, E., Ségonne, F., Salat, D. H., Busa, E., Seidman, L. J., Goldstein, J., Kennedy, D., Caviness, V., Makris, N., Rosen, B., & Dale, A. M. (2004). Automatically parcellating the human cerebral cortex. *Cerebral Cortex*, 14, 11–22. <https://doi.org/10.1093/cercor/bhg087>
- Fornia, L., Puglisi, G., Leonetti, A., Bello, L., Berti, A., Cerri, G., & Garbarini, F. (2020). Direct electrical stimulation of the premotor cortex shuts down awareness of voluntary actions. *Nature Communications*, 11, 705. <https://doi.org/10.1038/s41467-020-14517-4>
- Fox, K. C. R., Shi, L., Baek, S., Raccach, O., Foster, B. L., Saha, S., Margulies, D. S., Kucyi, A., & Parvizi, J. (2020). Intrinsic network architecture predicts the effects elicited by intracranial electrical stimulation of the human brain. *Nature Human Behaviour*, 4, 1039–1052. <https://doi.org/10.1038/s41562-020-0910-1>
- Fox, K. C. R., Yih, J., Raccach, O., Pendekanti, S. L., Limbach, L. E., Maydan, D. D., & Parvizi, J. (2018). Changes in subjective experience elicited by direct stimulation of the human orbitofrontal cortex. *Neurology*, 91, e1519–e1527. <https://doi.org/10.1212/WNL.0000000000006358>
- Frank, S. M., Baumann, O., Mattingley, J. B., & Greenlee, M. W. (2014). Vestibular and visual responses in human posterior insular cortex. *Journal of Neurophysiology*, 112, 2481–2491. <https://doi.org/10.1152/jn.00078.2014>
- Frank, S. M., & Greenlee, M. W. (2018). The parieto-insular vestibular cortex in humans: More than a single area? *Journal of Neurophysiology*, 120, 1438–1450. <https://doi.org/10.1152/jn.00907.2017>
- Fried, I., Katz, A., McCarthy, G., Sass, K., Williamson, P., Spencer, S., & Spencer, D. (1991). Functional organization of human supplementary motor cortex studied by electrical stimulation. *The Journal of Neuroscience*, 11, 3656–3666. <https://doi.org/10.1523/JNEUROSCI.11-11-03656.1991>
- Gallagher, S. (2000). Philosophical conceptions of the self: implications for cognitive science. *Trends in Cognitive Sciences*, 4, 14–21.
- Gallagher, S. (2005). *How the body shapes the mind*. Oxford University Press, New York.
- Gandola, M., Invernizzi, P., Sedda, A., Ferrè, E. R., Sterzi, R., Sberna, M., Paulesu, E., & Bottini, G. (2012). An anatomical account of somatoparaphrenia. *Cortex*, 48, 1165–1178. <https://doi.org/10.1016/j.cortex.2011.06.012>
- Gentile, G., Björnsdotter, M., Petkova, V. I., Abdulkarim, Z., & Ehrsson, H. H. (2015). Patterns of neural activity in the human ventral premotor cortex reflect a whole-body multisensory percept. *NeuroImage*, 109, 328–340. <https://doi.org/10.1016/j.neuroimage.2015.01.008>
- George, D. D., Ojemann, S. G., Drees, C., & Thompson, J. A. (2020). Stimulation mapping using stereoelectroencephalography: Current and future directions. *Frontiers in Neurology*, 11, 320. <https://doi.org/10.3389/fneur.2020.00320>
- Grande, K. M., Ihnen, S. K. Z., & Arya, R. (2020). Electrical stimulation mapping of brain function: A comparison of subdural electrodes and stereo-EEG. *Frontiers in Human Neuroscience*, 14. <https://doi.org/10.3389/fnhum.2020.611291>
- Guldin, W. O., & Grüsser, O. J. (1998). Is there a vestibular cortex? *Trends in Neurosciences*, 21, 254–259.
- Guterstam, A., Björnsdotter, M., Bergouignan, L., Gentile, G., Li, T.-Q., & Ehrsson, H. H. (2015). Decoding illusory self-location from activity in the human hippocampus. *Frontiers in Human Neuroscience*, 9. <https://doi.org/10.3389/fnhum.2015.00412>

- Guterstam, A., Björnsdotter, M., Gentile, G., & Ehrsson, H. H. (2015). Posterior cingulate cortex integrates the senses of self-location and body ownership. *Current Biology*, 25, 1416–1425. <https://doi.org/10.1016/j.cub.2015.03.059>
- Guterstam, A., Collins, K. L., Cronin, J. A., Zeberg, H., Darvas, F., Weaver, K. E., Ojemann, J. G., & Ehrsson, H. H. (2019). Direct electrophysiological correlates of body ownership in human cerebral cortex. *Cerebral Cortex*, 29, 1328–1341. <https://doi.org/10.1093/cercor/bhy285>
- Haggard, P., & Chambon, V. (2012). Sense of agency. *Current Biology: CB*, 22, R390–R392. <https://doi.org/10.1016/j.cub.2012.02.040>
- Halgren, E., Walter, R. D., Cherlow, D. G., & Crandall, P. H. (1978). Mental phenomena evoked by electrical stimulation of the human hippocampal formation and amygdala. *Brain: A Journal of Neurology*, 101, 83–117. <https://doi.org/10.1093/brain/101.1.83>
- Hao, G., Wang, X., Yan, H., He, L., Ni, D., Qiao, L., Zhang, X., Yu, K., & Yu, T. (2022). Limb loss experience evoked by electric cortical stimulation. *Epileptic Disorders*, 24, 67–74. <https://doi.org/10.1016/epd.2021.1364>
- Hari, R., Hänninen, R., Mäkinen, T., Jousmäki, V., Forss, N., Seppä, M., & Salonen, O. (1998). Three hands: Fragmentation of human bodily awareness. *Neuroscience Letters*, 240, 131–134.
- Herbet, G. (2021). Should Complex Cognitive Functions Be Mapped With Direct Electrostimulation in Wide-Awake Surgery? A network perspective. *Frontiers in Neurology*, 12, 635439. <https://doi.org/10.3389/fneur.2021.635439>
- Herbet, G., Lafargue, G., & Duffau, H. (2016). The dorsal cingulate cortex as a critical gateway in the network supporting conscious awareness. *Brain*, 139, e23. <https://doi.org/10.1093/brain/awv381>
- Heydrich, L., & Blanke, O. (2013). Distinct illusory own-body perceptions caused by damage to posterior insula and extrastriate cortex. *Brain*, 136, 790–803. <https://doi.org/10.1093/brain/aws364>
- Heydrich, L., Dieguez, S., Grunwald, T., Seeck, M., & Blanke, O. (2010). Illusory own body perceptions: case reports and relevance for bodily self-consciousness. *Consciousness and Cognition*, 19, 702–710. <https://doi.org/10.1016/j.concog.2010.04.010>
- Hitier, M., Zhang, Y.-F., Sato, G., Besnard, S., Zheng, Y., & Smith, P. F. (2021). Stratification of hippocampal electrophysiological activation evoked by selective electrical stimulation of different angular and linear acceleration sensors in the rat peripheral vestibular system. *Hearing Research*, 403, 108173. <https://doi.org/10.1016/j.heares.2021.108173>
- Horii, A., Russell, N. A., Smith, P. F., Darlington, C. L., & Bilkey, D. K. (2004). Vestibular influences on CA1 neurons in the rat hippocampus: an electrophysiological study in vivo. *Experimental Brain Research*, 155, 245–250.
- Hsu, T.-Y., Zhou, J.-F., Yeh, S.-L., Northoff, G., & Lane, T. J. (2022). Intrinsic neural activity predisposes susceptibility to a body illusion. *Cereb. Cortex Commun.*, 3, tgac012. <https://doi.org/10.1093/texcom/tgac012>
- Ionta, S., Heydrich, L., Lenggenhager, B., Mouthon, M., Fornari, E., Chapuis, D., Gassert, R., & Blanke, O. (2011). Multisensory mechanisms in temporo-parietal cortex support self-location and first-person perspective. *Neuron*, 70, 363–374. <https://doi.org/10.1016/j.neuron.2011.03.009>
- Isnard, J., Guénot, M., Sindou, M., & Mauguière, F. (2004). Clinical manifestations of insular lobe seizures: A stereo-electroencephalographic study. *Epilepsia*, 45, 1079–1090. <https://doi.org/10.1111/j.0013-9580.2004.68903.x>
- Isnard, J., Taussig, D., Bartolomei, F., Bourdillon, P., Catenoux, H., Chassoux, F., Chipaux, M., Clémenceau, S., Colnat-Coulbois, S., Denuelle, M., Derrey, S., Devaux, B., Dorfmueller, G., Gilard, V., Guenot, M., Job-Chapron, A.-S., Landré, E., Lebas, A., Maillard, L., ... Sauleau, P. (2018). French guidelines on stereoelectroencephalography (SEEG). *Neurophysiologie Clinique*, 48, 5–13. [10.1016/j.neucli.2017.11.005](https://doi.org/10.1016/j.neucli.2017.11.005)
- Janzen, J., Schlindwein, P., Bense, S., Bauermann, T., Vucurevic, G., Stoeter, P., & Dieterich, M. (2008). Neural correlates of hemispheric dominance and ipsilaterality within the vestibular system. *NeuroImage*, 42, 1508–1518. <https://doi.org/10.1016/j.neuroimage.2008.06.026>
- Jeannerod, M. (2006). *Motor cognition: What actions tell to the self*. Oxford University Press.
- Kahane, P., Hoffmann, D., Minotti, L., & Berthoz, A. (2003). Reappraisal of the human vestibular cortex by cortical electrical stimulation study. *Annals of Neurology*, 54, 615–624.
- Kannape, O. A., & Blanke, O. (2013). Self in motion: Sensorimotor and cognitive mechanisms in gait agency. *Journal of Neurophysiology*, 110, 1837–1847. <https://doi.org/10.1152/jn.01042.2012>
- Kannape, O. A., Smith, E. J. T., Moseley, P., Roy, M. P., & Lenggenhager, B. (2019). Experimentally induced limb-disownership in mixed reality. *Neuropsychologia*, 124, 161–170. <https://doi.org/10.1016/j.neuropsychologia.2018.12.014>
- Kavounoudias, A., Roll, J. P., Anton, J. L., Nazarian, B., Roth, M., & Roll, R. (2008). Proprio-tactile integration for kinesthetic perception: an fMRI study. *Neuropsychologia*, 46, 567–575. <https://doi.org/10.1016/j.neuropsychologia.2007.10.002>
- Keenan, J. P., Rubio, J., Racioppi, C., Johnson, A., & Barnacz, A. (2005). The right hemisphere and the dark side of consciousness. *Cortex*, 41, 695–704; discussion 731–734. [https://doi.org/10.1016/s0010-9452\(08\)70286-7](https://doi.org/10.1016/s0010-9452(08)70286-7)
- Khateb, A., Simon, S. R., Dieguez, S., Lazeyras, F., Momjian-Mayor, I., Blanke, O., Landis, T., Pegna, A. J., & Annoni, J. M. (2009). Seeing the phantom: a functional magnetic resonance imaging study of a supernumerary phantom limb. *Annals of Neurology*, 65, 698–705. <https://doi.org/10.1002/ana.21647>
- Kircher, T. T., Senior, C., Phillips, M. L., Benson, P. J., Bullmore, E. T., Brammer, M., Simmons, A., Williams, S. C., Bartels, M., & David, A. S. (2000). Towards a functional neuroanatomy of self processing: effects of faces and words. *Brain Research. Cognitive Brain Research*, 10, 133–144. [https://doi.org/10.1016/s0926-6410\(00\)00036-7](https://doi.org/10.1016/s0926-6410(00)00036-7)
- Kirsch, V., Boegle, R., Keeser, D., Kierig, E., Ertl-Wagner, B., Brandt, T., & Dieterich, M. (2018). Handedness-dependent functional organizational patterns within the bilateral vestibular cortical network revealed by fMRI connectivity based parcellation. *NeuroImage*, 178, 224–237. <https://doi.org/10.1016/j.neuroimage.2018.05.018>
- Kirsch, V., Keeser, D., Hergenroeder, T., Erat, O., Ertl-Wagner, B., Brandt, T., & Dieterich, M. (2016). Structural and functional connectivity mapping of the vestibular circuitry from human brainstem to cortex. *Brain Structure & Function*, 221, 1291–1308. <https://doi.org/10.1007/s00429-014-0971-x>
- Koch, C. (2018). What is consciousness? *Nature*, 557, S8–S12. <https://doi.org/10.1038/d41586-018-05097-x>
- Kremer, S., Chassagnon, S., Hoffmann, D., Benabid, A., & Kahane, P. (2001). The cingulate hidden hand. *Journal of Neurology, Neurosurgery, and Psychiatry*, 70, 264–265. <https://doi.org/10.1136/jnnp.70.2.264>
- Kurth, F., Eickhoff, S. B., Schleicher, A., Hoemke, L., Zilles, K., & Amunts, K. (2010). Cytoarchitecture and probabilistic maps of the human posterior insular cortex. *Cerebral Cortex*, 20, 1448–1461. <https://doi.org/10.1093/cercor/bhp208>
- Kurth, F., Zilles, K., Fox, P. T., Laird, A. R., & Eickhoff, S. B. (2010). A link between the systems: functional differentiation and integration within the human insula revealed by meta-analysis. *Brain Structure & Function*, 214, 519–534. <https://doi.org/10.1007/s00429-010-0255-z>
- Lackner, J. R. (1988). Some proprioceptive influences on the perceptual representation of body shape and orientation. *Brain*, 111(Pt 2), 281–297.
- Lagarde, S., Bénar, C.-G., Wendling, F., & Bartolomei, F. (2022). Interictal functional connectivity in focal refractory epilepsies investigated by intracranial EEG. *Brain Connectivity*, 12(10), 850–869. <https://doi.org/10.1089/brain.2021.0190>

- Lambrey, S., Doeller, C., Berthoz, A., & Burgess, N. (2012). Imagining being somewhere else: neural basis of changing perspective in space. *Cerebral Cortex*, 22, 166–174. <https://doi.org/10.1093/cercor/bhr101>
- Lanteaume, L., Khalifa, S., Regis, J., Marquis, P., Chauvel, P., & Bartolomei, F. (2007). Emotion induction after direct intracerebral stimulations of human amygdala. *Cerebral Cortex*, 17, 1307–1313. <https://doi.org/10.1093/cercor/bhl041>
- Lenggenhager, B., & Lopez, C. (2015). Vestibular contributions to the sense of body, self, and others. In T. Metzinger & J. M. Windt (Eds.), *Open MIND* (pp. 1–38). MIND-Group.
- Lenggenhager, B., Tadi, T., Metzinger, T., & Blanke, O. (2007). Video ergo sum: manipulating bodily self-consciousness. *Science*, 317, 1096–1099.
- Lesser, R. P., Crone, N. E., & Webber, W. R. S. (2010). Subdural electrodes. *Clinical Neurophysiology*, 121, 1376–1392. <https://doi.org/10.1016/j.clinph.2010.04.037>
- Limanowski, J., & Blankenburg, F. (2015). Network activity underlying the illusory self-attribution of a dummy arm. *Human Brain Mapping*, 36, 2284–2304. <https://doi.org/10.1002/hbm.22770>
- Limanowski, J., Lutti, A., & Blankenburg, F. (2014). The extrastriate body area is involved in illusory limb ownership. *NeuroImage*, 86, 514–524. <https://doi.org/10.1016/j.neuroimage.2013.10.035>
- Linkenauger, S. A., Leyrer, M., Bühlhoff, H. H., & Mohler, B. J. (2013). Welcome to wonderland: the influence of the size and shape of a virtual hand on the perceived size and shape of virtual objects. *PLoS One*, 8, e68594. <https://doi.org/10.1371/journal.pone.0068594>
- Lira, M., Pantaleão, F. N., de Souza Ramos, C. G., & Boggio, P. S. (2018). Anodal transcranial direct current stimulation over the posterior parietal cortex reduces the onset time to the rubber hand illusion and increases the body ownership. *Experimental Brain Research*, 236, 2935–2943. <https://doi.org/10.1007/s00221-018-5353-9>
- Lopez, C. (2013). A neuroscientific account of how vestibular disorders impair bodily self-consciousness. *Frontiers in Integrative Neuroscience*, 7. <https://doi.org/10.3389/fnint.2013.00091>
- Lopez, C. (2016). The vestibular system: balancing more than just the body. *Current Opinion in Neurology*, 29, 74–83. <https://doi.org/10.1097/WCO.0000000000000286>
- Lopez, C., Blanke, O., & Mast, F. W. (2012). The vestibular cortex in the human brain revealed by coordinate-based activation likelihood estimation meta-analysis. *Neuroscience*, 212, 159–179.
- Lopez, C., & Elzière, M. (2018). Out-of-body experience in vestibular disorders - A prospective study of 210 patients with dizziness. *Cortex*, 104, 193–206. <https://doi.org/10.1016/j.cortex.2017.05.026>
- Lopez, C., Schreyer, H. M., Preuss, N., & Mast, F. W. (2012). Vestibular stimulation modifies the body schema. *Neuropsychologia*, 50, 1830–1837. <https://doi.org/10.1016/j.neuropsychologia.2012.04.008>
- Makris, N., Papadimitriou, G. M., Sorg, S., Kennedy, D. N., Caviness, V. S., & Pandya, D. N. (2007). The occipitofrontal fascicle in humans: a quantitative, in vivo, DT-MRI study. *NeuroImage*, 37, 1100–1111. <https://doi.org/10.1016/j.neuroimage.2007.05.042>
- Mandonnet, E. (2021). Should Complex Cognitive Functions Be Mapped With Direct Electrostimulation in Wide-Awake Surgery? A commentary. *Frontiers in Neurology*, 12, 721038. <https://doi.org/10.3389/fneur.2021.721038>
- Mandonnet, E., Margulies, D., Stengel, C., Dali, M., Rheault, F., Toba, M. N., Bonnetblanc, F., & Valero-Cabre, A. (2020). “I do not feel my hand where I see it”: causal mapping of visuo-proprioceptive integration network in a surgical glioma patient. *Acta Neurochirurgica*, 162, 1949–1955. <https://doi.org/10.1007/s00701-020-04399-2>
- Mandonnet, E., Winkler, P. A., & Duffau, H. (2010). Direct electrical stimulation as an input gate into brain functional networks: principles, advantages and limitations. *Acta Neurochirurgica*, 152, 185–193. <https://doi.org/10.1007/s00701-009-0469-0>
- Martinaud, O., Besharati, S., Jenkinson, P. M., & Fotopoulou, A. (2017). Ownership illusions in patients with body delusions: Different neural profiles of visual capture and disownership. *Cortex, Confabulation and Related Disorders*, 87, 174–185. <https://doi.org/10.1016/j.cortex.2016.09.025>
- Mazzola, L., Isnard, J., Peyron, R., Guénot, M., & Mauguière, F. (2009). Somatotopic organization of pain responses to direct electrical stimulation of the human insular cortex. *Pain*, 146, 99–104. <https://doi.org/10.1016/j.pain.2009.07.014>
- Mazzola, L., Lopez, C., Faillenot, I., Chouchou, F., Mauguière, F., & Isnard, J. (2014). Vestibular responses to direct stimulation of the human insular cortex. *Annals of Neurology*, 76, 609–619. <https://doi.org/10.1002/ana.24252>
- Mazzola, L., Mauguière, F., & Isnard, J. (2019). Functional mapping of the human insula: Data from electrical stimulations. *Revue Neurologique (Paris)*, 175, 150–156. <https://doi.org/10.1016/j.neuro.2018.12.003>
- McGonigal, A., Bartolomei, F., & Chauvel, P. (2021). On seizure semiology. *Epilepsia*, 62, 2019–2035. <https://doi.org/10.1111/epi.16994>
- Medina Villalon, S., Paz, R., Roehri, N., Lagarde, S., Pizzo, F., Colombet, B., Bartolomei, F., Carron, R., & Bénar, C.-G. (2018). EpiTools, A software suite for presurgical brain mapping in epilepsy: Intracerebral EEG. *Journal of Neuroscience Methods*, 303, 7–15. <https://doi.org/10.1016/j.jneumeth.2018.03.018>
- Mercier, M. R., Dubarry, A.-S., Tadel, F., Avanzini, P., Axmacher, N., Cellier, D., Vecchio, M. D., Hamilton, L. S., Hermes, D., Kahana, M. J., Knight, R. T., Llorens, A., Megevand, P., Melloni, L., Miller, K. J., Piai, V., Puce, A., Ramsey, N. F., Schwiedrzik, C. M., ... Oostenveld, R. (2022). Advances in human intracranial electroencephalography research, guidelines and good practices. *NeuroImage*, 260, 119438. <https://doi.org/10.1016/j.neuroimage.2022.119438>
- Miller, J. F., Neufang, M., Solway, A., Brandt, A., Trippel, M., Mader, I., Hefft, S., Merkow, M., Polyn, S. M., Jacobs, J., Kahana, M. J., & Schulze-Bonhage, A. (2013). Neural activity in human hippocampal formation reveals the spatial context of retrieved memories. *Science*, 342, 1111–1114. <https://doi.org/10.1126/science.1244056>
- Miyazawa, N., Hayashi, M., Komiya, K., & Akiyama, I. (2004). Supernumerary phantom limbs associated with left hemispheric stroke: Case report and review of the literature. *Neurosurgery*, 54, 228–231. <https://doi.org/10.1227/01.NEU.0000097558.01639.F5>
- Mohr, C., Blanke, O., & Brugger, P. (2006). Perceptual aberrations impair mental own-body transformations. *Behavioral Neuroscience*, 120, 528–534.
- Moore, J. W., Ruge, D., Wenke, D., Rothwell, J., & Haggard, P. (2010). Disrupting the experience of control in the human brain: pre-supplementary motor area contributes to the sense of agency. *Proceedings of the Royal Society B: Biological Sciences*, 277, 2503–2509. <https://doi.org/10.1098/rspb.2010.0404>
- Mulak, A., Kahane, P., Hoffmann, D., Minotti, L., & Bonaz, B. (2008). Brain mapping of digestive sensations elicited by cortical electrical stimulations. *Neurogastroenterology and Motility*, 20, 588–596. <https://doi.org/10.1111/j.1365-2982.2007.01066.x>
- Mullan, S., & Penfield, W. (1959). Illusions of comparative interpretation and emotion; production by epileptic discharge and by electrical stimulation in the temporal cortex. *A.M.A. Archives of Neurology and Psychiatry*, 81, 269–284.
- Naito, E., Morita, T., Saito, D. N., Ban, M., Shimada, K., Okamoto, Y., Kosaka, H., Okazawa, H., & Asada, M. (2017). Development of right-hemispheric dominance of inferior parietal lobule in proprioceptive illusion task. *Cerebral Cortex*, 27, 5385–5397. <https://doi.org/10.1093/cercor/bhx223>
- Nakul, E., Bartolomei, F., & Lopez, C. (2021). Vestibular-evoked cerebral potentials. *Frontiers in Neurology*, 12, 674100. <https://doi.org/10.3389/fneur.2021.674100>
- Nakul, E., Dabard, C., Toupet, M., Hautefort, C., van Nechel, C., Lenggenhager, B., & Lopez, C. (2020). Interoception and embodiment in patients with bilateral vestibulopathy. *Journal of Neurology*, 267, 109–117. <https://doi.org/10.1007/s00415-020-10221-x>

- Nakul, E., & Lopez, C. (2017). Commentary: out-of-body experience during awake craniotomy. *Frontiers in Human Neuroscience*, 11, 417. <https://doi.org/10.3389/fnhum.2017.00417>
- Nakul, E., Orlando-Dessaints, N., Lenggenhager, B., & Lopez, C. (2020). Measuring perceived self-location in virtual reality. *Scientific Reports*, 10, 6802. <https://doi.org/10.1038/s41598-020-63643-y>
- O'Keefe, J., & Conway, D. H. (1978). Hippocampal place units in the freely moving rat: why they fire where they fire. *Experimental Brain Research*, 31, 573–590.
- O'Mara, S. M., Rolls, E. T., Berthoz, A., & Kesner, R. P. (1994). Neurons responding to whole-body motion in the primate hippocampus. *The Journal of Neuroscience*, 14, 6511–6523.
- Oane, I., Barborica, A., Chetan, F., Donos, C., Maliia, M. D., Arbune, A. A., Daneasa, A., Pistol, C., Nica, A. E., Bajenaru, O. A., & Mindruta, I. (2020). Cingulate cortex function and multi-modal connectivity mapped using intracranial stimulation. *NeuroImage*, 220, 117059. <https://doi.org/10.1016/j.neuroimage.2020.117059>
- Odegaard, B., Knight, R. T., & Lau, H. (2017). Should a Few Null Findings Falsify Prefrontal Theories of Conscious Perception? *The Journal of Neuroscience*, 37, 9593–9602. <https://doi.org/10.1523/JNEUROSCI.3217-16.2017>
- Ostrowsky, K., Magnin, M., Rylvlin, P., Isnard, J., Guenot, M., & Mauguiere, F. (2002). Representation of pain and somatic sensation in the human insula: a study of responses to direct electrical cortical stimulation. *Cerebral Cortex*, 12, 376–385.
- Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E., Chou, R., Glanville, J., Grimshaw, J. M., Hróbjartsson, A., Lalu, M. M., Li, T., Loder, E. W., Mayo-Wilson, E., McDonald, S., ... Moher, D. (2021). The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*, 372, n71. <https://doi.org/10.1136/bmj.n71>
- Park, H.-D., & Blanke, O. (2019). Coupling inner and outer body for self-consciousness. *Trends in Cognitive Sciences*, 23, 377–388. <https://doi.org/10.1016/j.tics.2019.02.002>
- Parvizi, J. (2009). Corticocentric myopia: Old bias in new cognitive sciences. *Trends in Cognitive Sciences*, 13, 354–359. <https://doi.org/10.1016/j.tics.2009.04.008>
- Parvizi, J., Braga, R. M., Kucyi, A., Veit, M. J., Pinheiro-Chagas, P., Perry, C., Sava-Segal, C., Zeineh, M., van Staalduinen, E. K., Henderson, J. M., & Markert, M. (2021). Altered sense of self during seizures in the posteromedial cortex. *PNAS*, 118(29), e2100522118. <https://doi.org/10.1073/pnas.2100522118>
- Parvizi, J., & Kastner, S. (2018). Promises and limitations of human intracranial electroencephalography. *Nature Neuroscience*, 21, 474–483. <https://doi.org/10.1038/s41593-018-0108-2>
- Penfield, W. (1947). Some observations on the cerebral cortex of man. *Proceedings of the Royal Society B: Biological Sciences*, 134, 329–347.
- Penfield, W. (1955). The twenty-ninth Maudsley lecture: the role of the temporal cortex in certain psychical phenomena. *The Journal of Mental Science*, 101, 451–465.
- Penfield, W. (1957). Vestibular sensation and the cerebral cortex. *The Annals of Otolaryngology, Rhinology, and Laryngology*, 66, 691–698. <https://doi.org/10.1177/000348945706600307>
- Penfield, W. (1958). Some mechanisms of consciousness discovered during electrical stimulation of the brain. *Proceedings of the National Academy of Sciences of the United States of America*, 44, 51–66.
- Penfield, W., & Boldrey, E. (1937). Somatic motor and sensory representation in the cerebral cortex of man as studied by electric stimulation. *Brain*, 60, 389–443. <https://doi.org/10.1093/brain/60.4.389>
- Penfield, W., & Perot, P. (1963). The brain's record of auditory and visual experience. A final summary and discussion. *Brain: A Journal of Neurology*, 86, 595–696. <https://doi.org/10.1093/brain/86.4.595>
- Penfield, W., & Rasmussen, T. (1950). *The cerebral cortex of man. A clinical study of localization of function*. Macmillan, Oxford.
- Petkova, V. I., Bjornsdotter, M., Gentile, G., Jonsson, T., Li, T. Q., & Ehrsson, H. H. (2011). From part- to whole-body ownership in the multisensory brain. *Current Biology*, 21, 1118–1122. <https://doi.org/10.1016/j.cub.2011.05.022>
- Pfeiffer, C., Lopez, C., Schmutz, V., Duenas, J. A., Martuzzi, R., & Blanke, O. (2013). Multisensory origin of the subjective first-person perspective: visual, tactile, and vestibular mechanisms. *PLoS One*, 8, e61751. <https://doi.org/10.1371/journal.pone.0061751>
- Piryankova, I. V., Wong, H. Y., Linkenauger, S. A., Stinson, C., Longo, M. R., Bühlhoff, H. H., & Mohler, B. J. (2014). Owning an overweight or underweight body: distinguishing the physical, experienced and virtual body. *PLoS One*, 9, e103428. <https://doi.org/10.1371/journal.pone.0103428>
- Popa, I., Barborica, A., Scholly, J., Donos, C., Bartolomei, F., Lagarde, S., Hirsch, E., Valenti-Hirsch, M.-P., Maliia, M. D., Arbune, A. A., Daneasa, A., Ciurea, J., Bajenaru, O.-A., & Mindruta, I. (2019). Illusory own body perceptions mapped in the cingulate cortex—An intracranial stimulation study. *Human Brain Mapping*, 40, 2813–2826. <https://doi.org/10.1002/hbm.24563>
- Poucet, B., Lenck-Santini, P. P., Paz-Villagran, V., & Save, E. (2003). Place cells, neocortex and spatial navigation: A short review. *Journal of Physiology, Paris*, 97, 537–546.
- Procyk, E., Wilson, C. R. E., Stoll, F. M., Faraut, M. C. M., Petrides, M., & Amiez, C. (2016). Midcingulate motor map and feedback detection: Converging data from humans and monkeys. *Cerebral Cortex*, 26, 467–476. <https://doi.org/10.1093/cercor/bhu213>
- Racah, O., Block, N., & Fox, K. C. R. (2021). Does the Prefrontal Cortex Play an Essential Role in Consciousness? Insights from Intracranial Electrical Stimulation of the Human Brain. *The Journal of Neuroscience*, 41, 2076–2087. <https://doi.org/10.1523/JNEUROSCI.1141-20.2020>
- Raiser, T. M., Flanagan, V. L., Duering, M., van Ombergen, A., Ruehl, R. M., & Zu Eulenburg, P. (2020). The human corticocortical vestibular network. *NeuroImage*, 223, 117362. <https://doi.org/10.1016/j.neuroimage.2020.117362>
- Richer, F., Martinez, M., Robert, M., Bouvier, G., & Saint-Hilaire, J. M. (1993). Stimulation of human somatosensory cortex: Tactile and body displacement perceptions in medial regions. *Experimental Brain Research*, 93, 173–176. <https://doi.org/10.1007/BF00227792>
- Ritaccio, A. L., Brunner, P., & Schalk, G. (2018). Electrical stimulation mapping of the brain: Basic principles and emerging alternatives. *Journal of Clinical Neurophysiology*, 35, 86–97. <https://doi.org/10.1097/WNP.0000000000000440>
- Ruby, P., & Decety, J. (2001). Effect of subjective perspective taking during simulation of action: A PET investigation of agency. *Nature Neuroscience*, 4, 546–550.
- Rushworth, M. F. S., Behrens, T. E. J., & Johansen-Berg, H. (2006). Connection patterns distinguish 3 regions of human parietal cortex. *Cerebral Cortex*, 16, 1418–1430. <https://doi.org/10.1093/cercor/bhj079>
- Salanova, V., Andermann, F., Rasmussen, T., Olivier, A., & Quesney, L. F. (1995a). Tumoural parietal lobe epilepsy. Clinical manifestations and outcome in 34 patients treated between 1934 and 1988. *Brain*, 118(Pt 5), 1289–1304. <https://doi.org/10.1093/brain/118.5.1289>
- Salanova, V., Andermann, F., Rasmussen, T., Olivier, A., & Quesney, L. F. (1995b). Parietal lobe epilepsy. Clinical manifestations and outcome in 82 patients treated surgically between 1929 and 1988. *Brain*, 118(Pt 3), 607–627. <https://doi.org/10.1093/brain/118.3.607>
- Schaller, K., Iannotti, G. R., Orepic, P., Betka, S., Haemmerli, J., Boex, C., Alcoba-Banqueri, S., Garin, D. F. A., Herbelin, B., Park, H.-D., Michel, C. M., & Blanke, O. (2021). The perspectives of mapping and monitoring of the sense of self in neurosurgical patients. *Acta Neurochirurgica*, 163, 1213–1226. <https://doi.org/10.1007/s00701-021-04778-3>
- Schneider, R. J., Friedman, D. P., & Mishkin, M. (1993). A modality-specific somatosensory area within the insula of the rhesus monkey. *Brain Research*, 621, 116–120.

- Seghier, M. L. (2013). The Angular Gyrus. *The Neuroscientist*, 19, 43–61. <https://doi.org/10.1177/1073858412440596>
- Selimbeyoglu, A., & Parvizi, J. (2010). Electrical stimulation of the human brain: perceptual and behavioral phenomena reported in the old and new literature. *Frontiers in Human Neuroscience*, 4, 46. <https://doi.org/10.3389/fnhum.2010.00046>
- Serino, A., Alsmith, A., Costantini, M., Mandrigin, A., Tajadura-Jimenez, A., & Lopez, C. (2013). Bodily ownership and self-location: Components of bodily self-consciousness. *Consciousness and Cognition*, 22, 1239–1252. <https://doi.org/10.1016/j.concog.2013.08.013>
- Seth, A. K. (2013). Interoceptive inference, emotion, and the embodied self. *Trends in Cognitive Sciences*, 17, 565–573. <https://doi.org/10.1016/j.tics.2013.09.007>
- Simeon, D., & Abugle, J. (2006). *Feeling unreal. Depersonalization disorder and the loss of the self*. Oxford University Press.
- Smith, A. T., Wall, M. B., & Thilo, K. V. (2011). Vestibular inputs to human motion-sensitive visual cortex. *Cerebral Cortex*, 22(5), 1068–1077. <https://doi.org/10.1093/cercor/bhr179>
- Smith, P. F. (1997). Vestibular-hippocampal interactions. *Hippocampus*, 7, 465–471.
- So, E. L., & Schaüble, B. S. (2004). Ictal asomatognosia as a cause of epileptic falls: simultaneous video, EMG, and invasive EEG. *Neurology*, 63, 2153–2154. <https://doi.org/10.1212/01.wnl.0000145628.38030.3e>
- Sun, F., Zhang, G., Ren, L., Yu, T., Ren, Z., Gao, R., & Zhang, X. (2021). Functional organization of the human primary somatosensory cortex: A stereo-electroencephalography study. *Clinical Neurophysiology*, 132, 487–497. <https://doi.org/10.1016/j.clinph.2020.11.032>
- Sun, F., Zhang, G., Yu, T., Zhang, X., Wang, X., Yan, X., Qiao, L., Ma, K., & Zhang, X. (2021). Functional characteristics of the human primary somatosensory cortex: An electrostimulation study. *Epilepsy & Behavior*, 118, 107920. <https://doi.org/10.1016/j.yebeh.2021.107920>
- Suzuki, M., Kitano, H., Ito, R., Kitaniishi, T., Yazawa, Y., Ogawa, T., Shiino, A., & Kitajima, K. (2001). Cortical and subcortical vestibular response to caloric stimulation detected by functional magnetic resonance imaging. *Brain Research. Cognitive Brain Research*, 12, 441–449.
- Trébuchon, A., & Chauvel, P. (2016). Electrical stimulation for seizure induction and functional mapping in stereoelectroencephalography. *Journal of Clinical Neurophysiology*, 33, 511–521. <https://doi.org/10.1097/WNP.0000000000000313>
- Trebuchon, A., Racila, R., Cardinale, F., Lagarde, S., McGonigal, A., Lo Russo, G., Scavarda, D., Carron, R., Mai, R., Chauvel, P., Bartolomei, F., & Francione, S. (2020). Electrical stimulation for seizure induction during SEEG exploration: a useful predictor of postoperative seizure recurrence? *Journal of Neurology, Neurosurgery, and Psychiatry*, 92, 22–26. <https://doi.org/10.1136/jnnp-2019-322469>
- Tsakiris, M. (2010). My body in the brain: A neurocognitive model of body-ownership. *Neuropsychologia*, 48, 703–712. <https://doi.org/10.1016/j.neuropsychologia.2009.09.034>
- Tsakiris, M., Schutz-Bosbach, S., & Gallagher, S. (2007). On agency and body-ownership: Phenomenological and neurocognitive reflections. *Consciousness and Cognition*, 16, 645–660.
- Tsakiris, M., Tajadura-Jiménez, A., & Costantini, M. (2011). Just a heartbeat away from one's body: Interoceptive sensitivity predicts malleability of body-representations. *Proceedings of the Biological Sciences*, 278, 2470–2476. <https://doi.org/10.1098/rspb.2010.2547>
- Uddin, L. Q., Supekar, K., Amin, H., Rykhlevskaia, E., Nguyen, D. A., Greicius, M. D., & Menon, V. (2010). Dissociable connectivity within human angular gyrus and intraparietal sulcus: evidence from functional and structural connectivity. *Cerebral Cortex*, 20, 2636–2646. <https://doi.org/10.1093/cercor/bhq011>
- van Elk, M., Duizer, M., Sligte, I., & van Schie, H. (2017). Transcranial direct current stimulation of the right temporoparietal junction impairs third-person perspective taking. *Cognitive, Affective, & Behavioral Neuroscience*, 17, 9–23. <https://doi.org/10.3758/s13415-016-0462-z>
- Vignal, J.-P., Maillard, L., McGonigal, A., & Chauvel, P. (2007). The dreamy state: hallucinations of autobiographic memory evoked by temporal lobe stimulations and seizures. *Brain*, 130, 88–99. <https://doi.org/10.1093/brain/awl329>
- Vitte, E., Derosier, C., Caritu, Y., Berthoz, A., Hasboun, D., & Soulie, D. (1996). Activation of the hippocampal formation by vestibular stimulation: a functional magnetic resonance imaging study. *Experimental Brain Research*, 112, 523–526.
- Vogeley, K., & Fink, G. R. (2003). Neural correlates of the first-person-perspective. *Trends in Cognitive Sciences*, 7, 38–42.
- Vogeley, K., May, M., Ritzl, A., Falkai, P., Zilles, K., & Fink, G. R. (2004). Neural correlates of first-person perspective as one constituent of human self-consciousness. *Journal of Cognitive Neuroscience*, 16, 817–827.
- Wiener, S. I., Berthoz, A., & Zugaro, M. B. (2002). Multisensory processing in the elaboration of place and head direction responses by limbic system neurons. *Brain Research. Cognitive Brain Research*, 14, 75–90.
- Wiest, G., Zimprich, F., Prayer, D., Czech, T., Serles, W., & Baumgartner, C. (2004). Vestibular processing in human paramedian precuneus as shown by electrical cortical stimulation. *Neurology*, 62, 473–475. <https://doi.org/10.1212/01.wnl.0000106948.17561.55>
- Windt, J. M. (2015). *Dreaming. A conceptual framework for philosophy of mind and empirical research*. MIT Press.
- Wirsich, J., Perry, A., Ridley, B., Proix, T., Golos, M., Benar, C., Ranjeva, J.-P., Bartolomei, F., Breakspear, M., Jirsa, V., & Guye, M. (2016). Whole-brain analytic measures of network communication reveal increased structure-function correlation in right temporal lobe epilepsy. *NeuroImage Clin.*, 11, 707–718. <https://doi.org/10.1016/j.nicl.2016.05.010>
- Yomogida, Y., Sugiura, M., Sassa, Y., Wakusawa, K., Sekiguchi, A., Fukushima, A., Takeuchi, H., Horie, K., Sato, S., & Kawashima, R. (2010). The neural basis of agency: An fMRI study. *NeuroImage*, 50, 198–207. <https://doi.org/10.1016/j.neuroimage.2009.12.054>
- Yu, K., Liu, C., Yu, T., Wang, X., Xu, C., Ni, D., & Li, Y. (2018). Out-of-body experience in the anterior insular cortex during the intracranial electrodes stimulation in an epileptic child. *Journal of Clinical Neuroscience*, 54, 122–125. <https://doi.org/10.1016/j.jocn.2018.04.050>
- Yu, K., Yu, T., Qiao, L., Liu, C., Wang, X., Zhou, X., Ni, D., Zhang, G., & Li, Y. (2018). Electrical stimulation of the insulo-opercular region: Visual phenomena and altered body-ownership symptoms. *Epilepsy Research*, 148, 96–106. <https://doi.org/10.1016/j.eplepsyres.2018.09.014>
- Zu Eulenburg, P., Caspers, S., Roski, C., & Eickhoff, S. B. (2012). Meta-analytical definition and functional connectivity of the human vestibular cortex. *NeuroImage*, 60, 162–169. <https://doi.org/10.1016/j.neuroimage.2011.12.032>

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Dary, Z., Lenggenhager, B., Lagarde, S., Medina Villalon, S., Bartolomei, F., & Lopez, C. (2023). Neural bases of the bodily self as revealed by electrical brain stimulation: A systematic review. *Human Brain Mapping*, 1–24. <https://doi.org/10.1002/hbm.26253>

Supplementary Table 1. PRIMSA Main Checklist

TITLE			
Title	1	Identify the report as a systematic review.	Page 1
ABSTRACT			
Abstract	2	See the PRISMA 2020 for Abstracts checklist	
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of existing knowledge.	Pages 3 and 5
Objectives	4	Provide an explicit statement of the objective(s) or question(s) the review addresses.	Page 5
METHODS			
Eligibility criteria	5	Specify the inclusion and exclusion criteria for the review and how studies were grouped for the syntheses.	Pages 6 and 7
Information sources	6	Specify all databases, registers, websites, organisations, reference lists and other sources searched or consulted to identify studies. Specify the date when each source was last searched or consulted.	Page 6 and Figure 2
Search strategy	7	Present the full search strategies for all databases, registers and websites, including any filters and limits used.	Page 6 and figure 2, and supplementary table 3
Selection process	8	Specify the methods used to decide whether a study met the inclusion criteria of the review, including how many reviewers screened each record and each report retrieved, whether they worked independently, and if applicable, details of automation tools used in the process.	Pages 6 and 7 and figure 2
Data collection process	9	Specify the methods used to collect data from reports, including how many reviewers collected data from each report, whether they worked independently, any processes for obtaining or confirming data from study investigators, and if applicable, details of automation tools used in the process.	Pages 6 and 7 and figure 2
Data items	10a	List and define all outcomes for which data were sought. Specify whether all results that were compatible with each outcome domain in each study were sought (e.g. for all measures, time points, analyses), and if not, the methods used to decide which results to collect.	Page 7
	10b	List and define all other variables for which data were sought (e.g. participant and intervention characteristics, funding sources). Describe any assumptions made about any missing or unclear information.	Page 7
Study risk of bias assessment	11	Specify the methods used to assess risk of bias in the included studies, including details of the tool(s) used, how many reviewers assessed each study and whether they worked independently, and if applicable, details of automation tools used in the process.	No registration
Effect measures	12	Specify for each outcome the effect measure(s) (e.g. risk ratio, mean difference) used in the synthesis or presentation of results.	No registration
Synthesis methods	13a	Describe the processes used to decide which studies were eligible for each synthesis (e.g. tabulating the study intervention characteristics and comparing against the planned groups for each synthesis (item 5)).	No registration
	13b	Describe any methods required to prepare the data for presentation or synthesis, such as handling of missing summary statistics, or data conversions.	Pages 7 and 8
	13c	Describe any methods used to tabulate or visually display results of individual studies and syntheses.	Pages 7 and 8
	13d	Describe any methods used to synthesize results and provide a rationale for the choice(s). If meta-analysis was performed, describe the model(s), method(s) to identify the presence and extent of statistical heterogeneity, and software package(s) used.	Pages 7 and 8
	13e	Describe any methods used to explore possible causes of heterogeneity among study results (e.g. subgroup analysis, meta-regression).	No registration
	13f	Describe any sensitivity analyses conducted to assess robustness of the synthesized results.	No registration

Reporting bias assessment	14	Describe any methods used to assess risk of bias due to missing results in a synthesis (arising from reporting biases).	No registration
Certainty assessment	15	Describe any methods used to assess certainty (or confidence) in the body of evidence for an outcome.	No registration
RESULTS			
Study selection	16a	Describe the results of the search and selection process, from the number of records identified in the search to the number of studies included in the review, ideally using a flow diagram.	Figure 2
	16b	Cite studies that might appear to meet the inclusion criteria, but which were excluded, and explain why they were excluded.	Figure 2
Study characteristics	17	Cite each included study and present its characteristics.	Table 2 and supplementary table 4
Risk of bias in studies	18	Present assessments of risk of bias for each included study.	No registration
Results of individual studies	19	For all outcomes, present, for each study: (a) summary statistics for each group (where appropriate) and (b) an effect estimate and its precision (e.g. confidence/credible interval), ideally using structured tables or plots.	Page 8 and table 2
Results of syntheses	20a	For each synthesis, briefly summarise the characteristics and risk of bias among contributing studies.	No registration
	20b	Present results of all statistical syntheses conducted. If meta-analysis was done, present for each the summary estimate and its precision (e.g. confidence/credible interval) and measures of statistical heterogeneity. If comparing groups, describe the direction of the effect.	Page 12
	20c	Present results of all investigations of possible causes of heterogeneity among study results.	No registration
	20d	Present results of all sensitivity analyses conducted to assess the robustness of the synthesized results.	No registration
Reporting biases	21	Present assessments of risk of bias due to missing results (arising from reporting biases) for each synthesis assessed.	No registration
Certainty of evidence	22	Present assessments of certainty (or confidence) in the body of evidence for each outcome assessed.	No registration
DISCUSSION			
	23a	Provide a general interpretation of the results in the context of other evidence.	Pages 13 and 20
	23b	Discuss any limitations of the evidence included in the review.	Pages 20 and 21
	23c	Discuss any limitations of the review processes used.	No registration
	23d	Discuss implications of the results for practice, policy, and future research.	Pages 21 and 22
OTHER INFORMATION			
Registration and protocol	24a	Provide registration information for the review, including register name and registration number, or state that the review was not registered.	No registration
	24b	Indicate where the review protocol can be accessed, or state that a protocol was not prepared.	No registration
	24c	Describe and explain any amendments to information provided at registration or in the protocol.	No registration
Support	25	Describe sources of financial or non-financial support for the review, and the role of the funders or sponsors in the review.	Page 22
Competing interests	26	Declare any competing interests of review authors.	Page 22
Availability of data, code and other materials	27	Report which of the following are publicly available and where they can be found: template data collection forms; data extracted from included studies; data used for all analyses; analytic code; any other materials used in the review.	Data extracted from included studies available in supplementary table 4

From: Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *MetaArXiv*. 2020, September 14. DOI: 10.31222/osf.io/v7gm2. For more information, visit: www.prisma-statement.org

Supplementary Table 2. PRISMA Abstract Checklist

TITLE			
Title	1	Identify the report as a systematic review.	Yes
BACKGROUND			
Objectives	2	Provide an explicit statement of the main objective(s) or question(s) the review addresses.	Yes
METHODS			
Eligibility criteria	3	Specify the inclusion and exclusion criteria for the review.	Yes
Information sources	4	Specify the information sources (e.g. databases, registers) used to identify studies and the date when each was last searched.	No
Risk of bias	5	Specify the methods used to assess risk of bias in the included studies.	No
Synthesis of results	6	Specify the methods used to present and synthesize results.	No
RESULTS			
Included studies	7	Give the total number of included studies and participants and summarise relevant characteristics of studies.	Yes
Synthesis of results	8	Present results for main outcomes, preferably indicating the number of included studies and participants for each. If meta-analysis was done, report the summary estimate and confidence/credible interval. If comparing groups, indicate the direction of the effect (i.e. which group is favoured).	Yes
DISCUSSION			
Limitations of evidence	9	Provide a brief summary of the limitations of the evidence included in the review (e.g. study risk of bias, inconsistency and imprecision).	No
Interpretation	10	Provide a general interpretation of the results and important implications.	No
OTHER			
Funding	11	Specify the primary source of funding for the review.	No
Registration	12	Provide the register name and registration number.	No

From: Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *MetaArXiv*. 2020, September 14. DOI: 10.31222/osf.io/v7gm2. For more information, visit: www.prisma-statement.org

Search #	Query in PubMed	Results
1	((("SEEG"[All Fields] OR "stereo-EEG" [All Fields] OR "electrical cortical stimulation*" [All Fields] OR "stereoencephalograph*" [All Fields] OR "electrical stim*" [All Fields] OR "direct electrical stimulation*" [All Fields] OR "direct stimulat*" [All Fields] OR "stereo EEG" [All Fields] OR "intracranial electroencephalog*" [All Fields] OR "stereo-electro-encephalograph*" [All Fields] OR "stimulation" [All Fields]) AND ("epilep*" [All Fields] OR "awake" [All Fields] OR "craniotom*" [All Fields] OR "glioma" [All Fields]))	23 035
2	("embodi*" [All Fields] OR "disembodi*" [All Fields] OR "doppelgänger" [All Fields] OR "feeling of a presence*" [All Fields] OR "heautoscop*" [All Fields] OR "depersonal*" [All Fields] OR "self-location" [All Fields] OR "first-person" [All Fields] OR "dereal*" [All Fields] OR "autoscop*" [All Fields] OR "bodily self" [All Fields] OR "out of body experience" [All Fields] OR "self-consciousness" [All Fields] OR "out-of-body illusion*" [All Fields] OR "body schema*" [All Fields] OR "body image*" [All Fields] OR "body illusion*" [All Fields] OR "body representation*" [All Fields] OR "body ownership" [All Fields] OR "sense of agenc*" [All Fields] OR "agency" [All Fields] OR "sense of ownership" [All Fields] OR "ownership" OR "body perception*" [All Fields] OR "vestibular" [All Fields] OR "posture" OR "proprioceptive" OR "proprioception")	386 617
3 = 1 AND 2	((("SEEG"[All Fields] OR "stereo-EEG" [All Fields] OR "electrical cortical stimulation*" [All Fields] OR "stereoencephalograph*" [All Fields] OR "electrical stim*" [All Fields] OR "direct electrical stimulation*" [All Fields] OR "direct stimulat*" [All Fields] OR "stereo EEG" [All Fields] OR "intracranial electroencephalog*" [All Fields] OR "stereo-electro-encephalograph*" [All Fields] OR "stimulation" [All Fields]) AND ("epilep*" [All Fields] OR "awake" [All Fields] OR "craniotom*" [All Fields] OR "glioma" [All Fields])) AND ("embodi*" [All Fields] OR "disembodi*" [All Fields] OR "doppelgänger" [All Fields] OR "feeling of a presence*" [All Fields] OR "heautoscop*" [All Fields] OR "depersonal*" [All Fields] OR "self-location" [All Fields] OR "first-person" [All Fields] OR "dereal*" [All Fields] OR "autoscop*" [All Fields] OR "bodily self" [All Fields] OR "out of body experience" [All Fields] OR "self-consciousness" [All Fields] OR "out-of-body illusion*" [All Fields] OR "body schema*" [All Fields] OR "body image*" [All Fields] OR "body illusion*" [All Fields] OR "body representation*" [All Fields] OR "body ownership" [All Fields] OR "sense of agenc*" [All Fields] OR "agency" [All Fields] OR "sense of ownership" [All Fields] OR "ownership" OR "body perception*" [All Fields] OR "vestibular" [All Fields] OR "posture" OR "proprioceptive" OR "proprioception")	393

Supplementary Table 3. Search terms used in PubMed (research done until March 23, 2022).

Supplementary Table 4. Cases of bodily self disturbances evoked by electrical brain stimulation sorted by year of publication. Components of bodily self disturbances evoked by electrical brain stimulation (EBS): SL (vest): self-location (vestibular); SL (OBE): self-location (out-of-body experience); SL (prop): self-location (proprioceptive); Ag.: agency; BI: body image; IPP: first-person perspective; Own.: ownership; Other (feeling of a presence, depersonalization, derealization). Stimulation parameters: F: frequency, I: intensity (or voltage), D: duration. S1: primary somatosensory cortex. BA: Brodmann area. ¹All coordinates are reported in the MNI space, unless otherwise specified. ²Coordinates in the Talairach space. ³Additional information provided by the authors who were contacted personally. ⁴Individual EBS evoking bodily self disturbances were reported as different cases when the number of patients was not indicated in the article.

Publication	Sample size	Patients population	EBS procedure	Category of bodily self disturbances	Brain area explored	Cases included in our review and phenomenological description	Stimulation parameters	Site of EBS evoking a response and EBS coordinates ¹		
								x	y	z
Penfield and Boldrey (1937)	163	Epilepsy	Awake brain surgery	SL (prop)	Frontal and parietal cortex	Case 1: “The patient realized he had not moved but observed that it felt as though his finger, arm, foot, etc., had moved”.		Fissure of Rolando		
				Ag.	Frontal, parietal, and temporal cortex	Case 2: “Desire to move”. “The patient said the tongue did not actually move, but as the mouth was closed this could not be verified objectively”.		Postcentral gyrus		
				Ag.	Frontal, parietal, and temporal cortex	Case 3: “Desire to move”. “She wanted to move her opposite hand up and down”.		Middle temporal gyrus		
Penfield (1947)	190	Epilepsy	Awake brain surgery	SL (vest), Other	Temporal cortex	Case 4: “I feel queer, as though I were floating away.” “I have a queer sensation as if I am not here”. “As though I were half and half here.” (Patient GA)		Right superior temporal gyrus		
Penfield (1955)	190	Epilepsy	Awake brain surgery	SL (OBE)	Temporal cortex	Case 5: ““Oh God! I am leaving my body”, an altered relationship to his own person as though he were outside of his body”. (Patient VF)		Right temporal region near the insula		
Penfield (1957)	108	Brain tumour, epilepsy	Awake brain surgery	SL (vest)	Temporal and central areas	Case 6: “Sinking feeling and sensation of full head” (Patient 5).		Right superior temporal gyrus		
Mullan and Penfield (1959)	217	Epilepsy	Awake brain surgery	Other	Temporal cortex	Case 7: “Queer feeling, as though everything were out of this world—the world is unreal; there is a strange feeling as though you weren’t the same person you were before—there are waves of peculiarity in things”. (Patient PB)		Left temporal cortex		
						Case 8: “As if I were not here.” (Patient HD)		Left middle temporal gyrus		
						Case 9: “Out of this world” (Patient DA)		Right Insula		
						Case 10: “There was a feeling as though I were somewhere else, as though you were all waiting here for me to come back. I have had that feeling before.” (Patient MM)		Right superior temporal gyrus		
Penfield and Perot (1963)	1132	Epilepsy	Awake brain surgery	SL (vest)	Temporal cortex	Case 10: “My whole body seemed to be moving back and forth, particularly my head” (Patient MM)		Right temporo-occipital junction		
Halgren et al. (1978)	36	Epilepsy	SEEG	SL (prop), Ag.	Temporal lobe	Cases 11–19: “Patient reports a movement of his body. Most commonly, this movement was a localized twitch, tremor, or small displacement”. “The patients perceived all the above movements as being induced by an agent outside themselves.” (14 stimulation)	F: 30 Hz I: 5–10 mA	Right and left anterior, middle, and posterior hippocampus, anterior and middle hippocampal gyrus		

				SL (vest)		Cases 20–25: “Spatial illusions: this included three types of responses: (i) an illusional translation of the entire body with respect to the environment; (ii) an illusional movement of the environment; and (iii) dizziness.” (9 stimulation)		Right and left amygdala, anterior and middle hippocampus
Fried et al. (1991)	13	Epilepsy	Subdural grid	Ag.	Fronto-parietal region	Case 26: “Urge to move”	F: 50 Hz D: 5 s	Supplementary motor area (side unknown)
						Case 27: “Urge to move”	F: 50 Hz D: 5 s	Supplementary motor area (side unknown)
						Case 28: “Urge to move”	F: 50 Hz D: 5 s	Supplementary motor area (side unknown)
						Case 29: “Urge to move right leg inward.” (Patient 12) (A3-B3)	F: 50 Hz I: 4 mA D: 5 s	Left supplementary motor area
						Case 29: “Urge to move right arm, slight extension right elbow.” (A4-B4) (Patient 12)	F: 50 Hz I: 5 mA D: 5 s	Left supplementary motor area
						Case 29: “Urge to move right arm” (A5-6) (Patient 12)	F: 50 Hz I: 4 mA D: 5 s	Left supplementary motor area
						Case 29: “Urge to move right arm away from midline”. (A5-6) (Patient 12)	F: 50 Hz I: 5 mA D: 5 s	Left supplementary motor area
						Case 29: “Urge to lift right elbow”. (A6-B6) (Patient 12)	F: 50 Hz I: 6 mA D: 5 s	Left supplementary motor area
						Case 29: “Urge to pronate right forearm, slight elbow flexion.” (A6-B6) (Patient 12)	F: 50 Hz I: 7 mA D: 5 s	Left supplementary motor area
						Case 29: “Sensation of tension at right wrist “urge” to move right hand and forearm”. (A6-B6) (Patient 12)	F: 50 Hz I: 8 mA D: 5 s	Left supplementary motor area
						Case 29: “Urge to move right arm” (A6-B6) (Patient 12)	F: 50 Hz I: 9 and 10 mA D: 5 s	Left supplementary motor area
				Case 29: “Strong urge to raise right elbow.” (A6-A7) (Patient 12)		F: 50 Hz I: 5 mA D: 5 s	Left supplementary motor area	
				SL (prop)		Case 30: “Subjective experience of movement in the absence of overt motor activity. (e.g. one patient reported “I fell my arm is moving”)”.	F: 50 Hz D: 5 s	Supplementary motor area (side unknown)
						Case 31: “Subjective experience of movement in the absence of overt motor activity.	F: 50 Hz D: 5 s	Supplementary motor area (side unknown)
	Case 32: “Subjective experience of movement in the absence of overt motor activity.	F: 50 Hz D: 5 s	Supplementary motor area (side unknown)					

Richer et al. (1993)	40	Epilepsy	SEEG	SL (prop)	Rolandic and parietal regions	Case 33: “Feeling his legs and seat floating above the bed”.	F: 50 Hz I: 1–10 mA D: 5 s	Sub-parietal sulcus: next to the horizontal portion of the sulcus, adjoining the cingulate sulcus (side unknown)			
				SL (vest)		Case 34: “Feeling their whole body floating just above the bed”.		Sub-parietal sulcus: next to the vertical portion of the sulcus, adjoining the cingulate sulcus (side unknown)			
						Case 35: “Feeling their whole body floating just above the bed”.		Sub-parietal sulcus: next to the horizontal portion of the sulcus, adjoining the cingulate sulcus (side unknown)			
Salanova et al. (1995a)	82	Epilepsy	Awake brain surgery	SL (vest)	Parietal lobe	Case 36: “Sensation of ‘rolling’ off the table”.	F: 60 Hz I: 4 V	Left superior parietal gyrus			
				BI		Case 37: Patient “felt a twisting sensation in the contralateral extremity”.		Left superior parietal, behind the postcentral gyrus			
				Other		Case 38: “A man behind”.		Left superior parietal lobule			
				Other		Case 39: “A patient described by Penfield (1938) reported a ‘far away sensation’, and ‘things seem distant and small’”.		Left superior parietal, just behind the sensory area			
				BI		Case 40: “Feeling that the leg is absent”.		Right parietal operculum and superior parietal behind the postcentral gyrus			
				BI		Case 41: “Limpness of right hand”.		Left parietal operculum behind the cyst			
Salanova et al. (1995b)	34	Epilepsy	Subdural grid, awake brain surgery	BI	Parietal lobe	Case 42: “Oh, I feel as though I were going to have an attack, because my hand feels as if it is going around like a screwdriver.”	F: 60 Hz I: 4V	Right inferior parietal lobe			
Blanke et al. (2000)	1	Epilepsy	Subdural grid	SL (vest)	Frontal, temporal, and parietal cortex around the central sulcus	Case 43: “Sliding towards the lower end of the bed”. “I’m rolling to the right and falling out of the bed”.	I: 4.5–5.5 mA D: 2 s	Left inferior parietal lobule at the anterior part of the intraparietal sulcus			
						Case 43: “I feel the urge to hold on to something in order to prevent myself from falling out of the bed”.		-62	-46	-21 ²	
						Case 43: “My whole body moves”. “He felt as if he were lying in a swinging hammock”.		-62	-59	23 ²	
Blanke et al. (2002)	1	Epilepsy	Subdural grid	SL (vest)	Frontal, temporal, and parietal cortex around the lateral sulcus	Case 45: “The patient reported that she was “sinking into the bed” or “falling from a height”.”	F: 50 Hz I: 2–3 mA D: 2 s 3 stim.	Right angular gyrus			
				BI		Case 45: “The patient was then asked to watch her (real) legs during the electrical stimulation. As before, she was lying down (upper body supported at an angle of 45°, legs outstretched). This time, she reported seeing her legs “becoming shorter”.”					F: 50 Hz I: 4.0, 4.5 mA D: 2 s 2 stim.
Kremer et al. (2001)	1	Epilepsy	SEEG	Ag.	Cingulate cortex	Case 44: “Urge to grasp”.	F: 50 Hz I: 1.2 mA D: 5 s	Right ventral bank of the anterior cingulate sulcus			

				BI		Case 45: “If the patient’s legs were bent before the stimulation, she reported that her legs appeared to be moving quickly towards her face and took evasive action”.	F: 50 Hz I: 4.0–5.0 mA D: 2 s 2 stim.	
				BI		Case 45: “When asked to look at her outstretched arms during the electrical stimulation, the patient felt as though her left arm was shortened”.	F: 50 Hz I: 4.5–5.0 mA D: 2 s 2 stim.	
				BI		Case 45: “If both arms were in the same position but bent by 90° at the elbow, she felt that her left lower arm and hand were moving towards her face”.	F: 50 Hz I: 4.5–5 mA D: 2s 2 stim.	
				BI		Case 45: “When her eyes were shut, she felt that her upper body was moving towards her legs, which were stable.”	F: 50 Hz I: 4.0–5.0 mA D: 2 s 2 stim.	
				1PP, SL (OBE)		Case 45: “I see myself lying in bed, from above, but I only see my legs and lower trunk” (i.e., autoscopia). “Feeling of “lightness” and “floating” about two metres above the bed, close to the ceiling”.	F: 50 Hz I: 3.5 mA D: 2 s	
Ostrowsky et al. (2002)	30	Epilepsy	SEEG	BI	Temporal cortex	Case 46: “An impression of “heaviness” in half the body.”	F: 1 Hz I: 1–5 mA D: 5–10 s and F: 50 Hz I: 0.8–6 mA D: 5 s	Temporo-polar cortex
						Case 47: “A bloating of the upper limb”.		
Kahane et al. (2003)	260	Epilepsy	SEEG	SL (vest)	Temporal, occipital, parietal, frontal and insular lobes	Case 48: “Levitation, lightness”.	F: 1 Hz I: 3 mA D: 40 s	Right temporal lobe: amygdala 27 -12 -17 ²
						Case 48: “Levitation, lightness”.	F: 1 Hz I: 3 mA D: 40 s	Right 2 nd temporal gyrus midpart 47 -12 -17 ²
						Case 49: “Sensation of flying”.	F: 50 Hz I: 1 mA D: 5 s	Left temporal lobe: superior longitudinal fasciculus -32 -34 28 ²
						Case 50: “Sensation of sinking into the bed, of being heavier”.	F: 1 Hz I: 3 mA D: 40 s	Left occipital lobe: cuneus -5 -79 26 ²
						Case 51: “Sensation to be tilted toward the right”.	F: 50 Hz I: 1 mA D: 5 s	Right parietal operculum 62 -13 15 ²
						Case 52: “Sensation of body oscillations, forward backward”.	F: 1 Hz I: 3 mA D: 40 s	Right parietal operculum 51 -17 21 ²
						Case 53: “Sensation to be thrust forward”.	F: 50 Hz I: 1 mA D: 5 s	Right parietal operculum 38 -24 22 ²

						Case 54: “Sensation to be attracted (head → body) toward the right”.	F: 50 Hz I: 1.5 mA D: 5 s	Right parietal operculum 41 -31 25 ²
						Case 55: “Falling backward”.	F: 50 Hz I: 3 mA D: 5 s	Right parietal operculum 47 -37 29 ²
						Case 56: “Falling backward (head and trunk)”.	F: 50 Hz I: 3 mA D: 5 s	Right parietal operculum 44 -45 28 ²
						Case 57: “Sensation of falling flat”.	F: 50 Hz I: 2 mA D: 5 s	Right precuneus 3 -38 48 ²
						Case 58: “Sensation to be pushed (head) from the left toward the right”.	F: 50 Hz I: 1.6 mA D: 5 s	Right precuneus 6 -44 54 ²
						Case 58: “Sensation of rocking motion (head → trunk), backward”.	F: 50 Hz I: 1.2 mA D: 5 s	Right anterior cingulate gyrus 9 3 31 ²
						Case 59: “Body oscillations toward the right and left.”	F: 50 Hz I: 3 mA D: 5 s	Right 3 rd frontal gyrus 56 24 15 ²
						Case 60: “Sensation of falling flat”.	F: 50 Hz I: 1 mA D: 5 s	Right deepness of cingulate sulcus (BA 7/5) 15 -36 50 ²
Wiest et al. (2004)	1	Epilepsy	Subdural grid	SL (vest)	Parietal cortex	Case 61: “Rocking and alternating tilting sensations of his body and the environment”.	I: 1 mA	Right paramedian precuneus
So and Schäuble (2004)	1	Epilepsy	Subdural grid	Own.	Frontocentral region	Case 62: “Electrocortical stimulation of the area of seizure discharge induced his ictal symptom of sudden estrangement of the left lower extremity from the rest of his body”.	I: 10–11 mA	Right posterior parietal region (superior parietal lobule)
Isnard et al. (2004)	50	Epilepsy	SEEG	Other	Temporal lobe and insula	Case 63–68: “Sensation of unreality”.		Insula
				SL (vest)		Case 69–74: “A sudden sensation of displacement of their body in space, such as a brisk forward projection, a vertical or horizontal rotation of their body, or a sensation of levitation”.		Insula
Arzy et al. (2006)	1	Epilepsy	Subdural grid	Other	Frontal, temporal, and parietal cortex around the lateral sulcus	Case 75: “The patient had the impression that somebody was behind her”. “The patient describing the “person” as young and of indeterminate sex, a “shadow” who did not speak or move, and whose position beneath her back was identical to her own”. “He is behind me, almost at my body, but I do not feel it”.	I: 10 mA 3 stim.	Left temporo-parietal junction -57.15 42.08 15.63 ³
						Case 75: “She again reported the presence of the sitting “person”, this time displaced behind her to her right and attempting to interfere with the execution of her task (“He wants to take the card”; “He doesn’t want me to read”)”.	I: 11 mA 1 stim.	-57.15 31.96 23.00 ³
Vignal et al. (2007)	180	Epilepsy	SEEG	Other	Temporal lobe	Case 76: “Impression of being elsewhere” (patient 5).	F: 50 Hz I: 3 mA D: 5 s	Right hippocampus

				Other		Case 77: “Impression of being elsewhere” (patient 9).	F: 50 Hz I: 2 mA D: 5 s	Left amygdala		
				SL (OBE)		Case 77: “Impression of leaving his body” (patient 9).	F: 50 Hz I: 2.5 mA D: 5 s	Left amygdala		
				Other		Case 78: “Impression of being elsewhere” (patient 12).	F: 50 Hz I: 1.5 mA D: 5 s	Left hippocampus		
De Ridder et al. (2007)	1	Tinnitus	Paddle electrode	SL (OBE)	Temporo-parietal junction	Case 79: “His perception of disembodiment always involved a location about 50 cm behind his body and off to the left. There was no autoscopia and no voluntary control of movements of the disembodied perception. The environment was visually perceived from his real person perspective, not from the disembodied perspective. Stimulation at these specific settings had similar effects whether the patient was in a sitting or lying position”.	F: 40 Hz I: 3.7 V	Right posterior part of the superior temporal gyrus at the junction of the angular gyrus		
Mulak et al. (2008)	339	Epilepsy	SEEG	Other	Different lobes and amygdala	Case 80: “Feeling of unreality”.	F: 1 Hz (2 stim.) F: 50 Hz (3 stim.) D: 5 or 40 s	Left parahippocampal gyrus ⁴		
						Case 81: “Feeling of unreality”.		Right parahippocampal gyrus ⁴		
						Case 82: “Feeling of unreality”.		Right parahippocampal gyrus ⁴		
						Case 83: “Feeling of unreality”.		Right parahippocampal gyrus ⁴		
						Case 84: “Feeling of unreality”.		Right parahippocampal gyrus ⁴		
Desmurget et al. (2009)	7	Brain tumour	Awake brain surgery	Ag., SL (prop)	Parietal and premotor cortex	Case 85: Stimulation produced “a pure intention, that is, a felt desire to move without any overt movement being produced or EMG activity recorded in the concerned muscles. (...) These patients experienced awareness of an illusory movement”. (e.g., “I felt a desire to lick my lips”, “I moved my mouth, I talked, what did I say.”) (Patient PP3)	I: 5 and 8 mA D: 4 s	Left supramarginal gyrus (BA 40)		
						Case 86: Experience awareness of an illusory movement (hand and foot). (Patient PP1)			I: 8 mA D: 4 s	Right angular gyrus (BA 39) and supramarginal gyrus (BA 40)
						Case 87: Experience awareness of an illusory movement (e.g., “she felt “like a will to move” her chest”). (Patient PP2)			I: 8 mA D: 4 s	Right angular gyrus (BA 39) and supramarginal gyrus (BA 40)
Mazzola et al. (2014)	219	Epilepsy	SEEG	SL (vest)	Insula	Case 88: “Feeling of flying”.	F: 50 Hz I: 0.2–3.5 mA D: 5 s	Right insula		
								35	–10	14
						Case 89: “Feeling of levitation, floating up in the air”.		Right insula		
								40	–18	3
								41	–11	–7
						Case 90: “Feeling of levitation”.		Right insula		
	39	–18	–5							
Case 91: “Feeling of levitation, floating on water”.	Left insula									
	–34	–13	–9							
Case 92: “Feeling of falling backward to left side”.	Right insula									
	40	–10	–1							

						<p>Case 93: “Feeling of body rising”.</p> <p>Case 94: “Feeling of body rising (like a bubble)”.</p> <p>Case 95: “Feeling of falling laterally on right side of bed”.</p> <p>Case 96: “Feeling of limb rising (as if she were flying away); associated with electrical sensation in both hands”.</p> <p>Case 97: “Feeling of being projected, of falling backward; associated with electrical sensation moving from left hand to head”.</p> <p>Case 98: “Feeling of falling to right side associated with auditory illusion (whistling in right ear)”.</p> <p>Case 99: “Feeling of falling backward”.</p> <p>Case 100: “Feeling of falling backward; associated with paraesthesia in left hand”.</p>		<p>Right insula 40 -15 1</p> <p>Left insula -36 -9 -2</p> <p>Right insula 40 -12 8</p> <p>Right insula 33 -10 14</p> <p>Right insula 36 -13 17</p> <p>Right insula 34 -8 -4</p> <p>Right insula 36 -1 12</p> <p>Right insula 35 -16 8</p>
Blanke et al. (2014)	5	Epilepsy, brain tumour, brain lesion and abnormality	Subdural grid	Other	<p>Insula, fronto-parietal cortex</p> <p>Temporal et parietal cortex</p> <p>Temporal lobe</p>	<p>Case 101: “Presence of a man, behind to her right, in peripersonal space, fear and anxiety”. (Patient d)</p> <p>Case 102: “Presence of a male shadow, behind to the right, same position, echopraxia”. (Patient j)</p> <p>Case 103: “Presence behind to the right, strictly unilateral, unpleasant, no echopraxia”. (Patient k)</p>	F: 50 Hz I: 0.5–11 mA D: 2 s	Unknown location
Balestrini et al. (2015)	274	Epilepsy	SEEG	<p>BI, SL (vest)</p> <p>Other</p>	Parietal cortex	<p>Cases 104–117: “Body image alteration: including altered subjective perception of body image or movement” (e.g., "Patient reports that he feels his right hand larger and swollen", "Patient feels his body moving towards the left side")”.</p> <p>Case 118–119: “Psychic phenomena (e.g., “Patient describes to feel “like a doll”, “Patient reports the feeling of “being as in a parallel world”, Patient describes to feel a sensation of strangeness, unfamiliarity”).</p>	F: 1–50 Hz D: 5–30 mA	<p>Right precuneus</p> <p>Right inferior parietal lobule</p> <p>Right post-central gyrus</p> <p>Right precuneus</p> <p>Right inferior parietal lobule</p> <p>Right posterior cingulum</p> <p>Right inferior parietal lobule</p> <p>Right inferior parietal lobule</p> <p>Right inferior parietal lobule</p> <p>Right precuneus</p> <p>Right superior parietal lobule</p> <p>Right inferior parietal lobule</p> <p>Right superior parietal lobule</p> <p>Right superior parietal lobule</p> <p>Right precuneus</p> <p>Left posterior cingulum</p>

Bos et al. (2016)	1	Low grade glioma	Awake brain surgery	SL (prop)	Parietal lobe, temporal lobe, and the sylvian fissure	Case 120: “The patient felt her right leg being drawn towards the opposite wall of the operating theatre”.	I: 8–12 mA	Left subcortical white matter in the temporo-parietal junction
				SL (OBE), IPP		Case 120: “She experienced a complete OBE with autoscopia, in which she felt as if she floated just below the ceiling and saw her own body lying on the operating table”.		Left subcortical white matter in the temporo-parietal junction
Caruana et al. (2018)	329	Epilepsy	SEEG	SL (vest)	Cingulate cortex	Case 121–145: e.g., “Feeling of falling into a void” (and dizziness, vertigo)”. (19 patients stimulated on the right side and 6 patients on the left side)	F: 50 Hz I: 0.4–5 mA D: 5 s	Left ventral posterior middle cingulate cortex
								-3.3 23.7 41.1 ³
								Right ventral posterior middle cingulate cortex
								3.9 -20.8 41.2 ³
								Left ventral anterior middle cingulate cortex
								-3 0.1 35.3 ³
Yu et al. (2018a)	43	Epilepsy	SEEG	BI	Operculo-insular cortex	Case 146: “The patient reported that he could not feel the existence of his right hand when we stimulated the contralateral contacts”.	F: 50 Hz I: 1.0 mA D: 3 s	Left superior part of the anterior insular long gyrus
				SL (vest)		Case 147: “Changes in the perception of the body’s location, such as the feeling of body elevation or movement to one side”.	F: 50 Hz I: 1–4 mA D: 3 s	Parietal operculum (side unknown)
				BI		Case 148: “Sensations of deficiency of the contralateral limb or trunk”.	F: 50 Hz I: 1–2 mA D: 3 s	Parietal operculum (side unknown)
Yu et al. (2018b)	1	Epilepsy	SEEG	SL (OBE), IPP	Frontal, temporal, and parietal areas	Case 149: “She saw herself elevating to the left upper side 2 m high under the ceiling without any vestibular sensations like floating, flying, rotation and vertigo. She felt the virtual body was real herself and saw her own body entirely lying in bed being electrode stimulated by the doctor and nurses walking around in the sickroom”.	F: 50 Hz I: 0.5 mA D: 3 s	Left anterior insula
Popa et al. (2019)	110	Epilepsy	SEEG	BI	Cingulate cortex	Case 150: “Sensation that the lower limb detaches from the body, loses it”. (Patient 2)	F: 50 Hz I: 0.75 mA D: 5 s	Right middle cingulate cortex
				BI		Case 151: “Right upper limb feels lighter”. (Patient 5)	F: 50 Hz I: 0.25 mA D: 5 s	Left middle cingulate cortex
				BI		Case 151: “Sensation that the upper right limb is heavier”. (Patient 5)	F: 50 Hz I: 1 mA D: 5 s	Left middle cingulate cortex
				SL (prop)		Case 152: “Sensation of floating of the left hemibody”. (Patient 1)	F: 50 Hz I: 1 mA D: 5 s	Right middle cingulate cortex
				BI, IPP		Case 153: “Sensation that the head turns to the right side, that the head will explode, will detach from the neck, sees himself from outside” (i.e., autoscopia). (Patient 7)	F: 50 Hz I: 2 mA D: 5 s	Left middle cingulate cortex
				BI		Case 154: “Sensation that the left upper limb becomes heavier”. (Patient 8)	F: 50 Hz I: 1 mA D: 5 s	Right middle cingulate cortex

				BI		Case 155: "Sensation that the upper right limb becomes heavier associated to pain". (Patient 11)	F: 50 Hz I: 1 mA D: 5 s	Left middle cingulate cortex
				BI		Case 156: "Sensation that the right hemibody and all the head become heavier". (Patient 10)	F: 50 Hz I: 1 mA D: 5 s	Left middle cingulate cortex
				SL (vest)		Case 157: "Sensation that the body is being pushed to the left". (Patient 3)	F: 50 Hz I: 1 mA D: 5 s	Left middle cingulate cortex
				SL (prop)		Case 158: "Feels that the right hand is pulled from behind". (Patient 6)	F: 50 Hz I: 0.75 mA D: 5 s	Right middle cingulate cortex
				SL (prop)		Case 159: "Sensation that the upper part of the body moves upwards". (Patient 9)	F: 50 Hz I: 2 mA D: 5 s	Right middle cingulate cortex
				SL (prop)		Case 160: "Sensation that upper right limb moves to the left side". (Patient 12)	F: 50 Hz I: 1mA D: 5 s	Right middle cingulate cortex
Sun et al. (2020)	20	Epilepsy	SEEG	BI	Anterior parietal lobe involving S1	Case 161: "An illusion that the hand was absent".	F: 50 Hz I: 0.1–6 mA D: 3 s	Right superior parietal lobule and post central gyrus 41.05 ± 6.26 -29.46 ± 8.66 52.3 ± 7.29
Mandonnet et al. (2020)	1	Glioblastoma	Awake brain surgery	BI	Superior parietal lobule	Case 162: "Feeling that there was a discrepancy between the visual and proprioceptive perceptions of her right hand: "I do not feel my (right) hand where I see it"."	F: 60 Hz I: 3 mA D: 3 s	Left superior parietal lobule
Fornia et al. (2020)	12	Low grade glioma	Awake brain surgery	Ag.	Premotor cortex and somatosensory cortex (S1)	Cases 163–170: EBS "was effective in interfering with motor execution when applied to both premotor cortex and S1, but, crucially, it dramatically altered the patients' motor awareness only when applied to the premotor cortex."	F: 60 Hz I: 2–4.5 mA	Left premotor cortex -60 9.2 35.6 ³ -58.4 8 38.8 ³ -60.6 5.8 38 ³ -61.1 3.3 38.9 ³ -61.7 1.9 38.6 ³ -61 -0.4 40.7 ³ -60.1 -1.5 42.1 ³ -55.3 1.7 42.5 ³ -57.3 -0.1 43.1 ³ -58.3 -0.7 45.4 ³ -47.3 -6.2 56.9 ³
Andelman-Gur et al. (2020)	62	Epilepsy	Subdural grid	Ag.	Frontal, parietal, temporal, occipital lobes	Cases 171–180⁴: Change in will/urge, e.g., "Feeling a strong resistance to performing the task said despite his conscious will to continue with the task something blocks him". See : case report (Andelman-Gur et al., 2019) ³ "I feel resistance to anything I am told to do [...] the possibility of continuing in any action is blocked".	F: 50 Hz I: 1–10 mA for subdural electrodes I: 1–4 mA for depth electrodes D: 5 s	Left precentral gyrus -2 -22 56 ³ -2 -32 56 ³ Left superior frontal gyrus -14 14 66 ³ -18 14 66 ³ -20 14 64 ³ -22 14 60 ³ -22 14 60 ³ -20 6 66 ³ -16 4 68 ³

								Left planum temporale
								-58 -36 18 ³
								Right supramarginal gyrus
								60 -42 26 ³
								Right superior temporal gyrus
								66 -40 16 ³
								Right middle temporal gyrus
								62 -12 -26 ³
								Right inferior temporal gyrus
								54 -42 -26 ³
								Left precentral gyrus
								-2 -22 56 ³
								Left superior frontal gyrus
								-18 2 70 ³
								Right inferior occipital gyrus
								30 -94 -4 ³
								48 -68 12 ³
								-36 -90 -14 ³
								Right middle frontal gyrus
								40 38 26 ³
								Right inferior temporal gyrus
								56 -62 -8 ³
								Left middle occipital gyrus
								-32 -82 30 ³
								Left middle cingulate cortex
								-12 -7.605 44.662 ³
								-12 -4.933 48.603 ³
								-14 12 35 ³
								-13.302 13.973 38.698 ³
								-14 12 35 ³
								-13.302 13.973 38.698 ³
								Right middle cingulate cortex
								8.413 7.374 30.351 ³
								12.018 6.838 34.01 ³
								4.982 21.97 34.989 ³
								11.236 21.426 37.687 ³
								11.236 21.426 37.687 ³
								8.301 22.96 38.805 ³
								Right anterior cingulate cortex
								3.731 11.152 22.828 ³
								6.142 8.843 26.096 ³
								Right posterior cingulate cortex
								6.888 -13.141 35.794 ³
								10.884 -10.899 37 ³
Oane et al. (2020)	47	Epilepsy	SEEG	SL (vest)	Cingulate cortex	<p>Cases 181–189⁴: Levitation, e.g., “Felt she was floating in the air”.</p> <p>Cases 190–192⁴: OBE: e.g.: “Felt she is getting out of her body”.</p> <p>Cases 193–195: “Altered perception related to location, gravity or displacement of whole-body or body-part”.</p> <p>Cases 196–198: “Altered perception related to location, gravity or displacement of whole-body or body-part”.</p> <p>Case 199: “Altered perception related to location, gravity or displacement of whole-body or body-part”.</p> <p>Case 200: “Altered perception related to location, gravity or displacement of whole-body or body-part”.</p>	F: 50 Hz I: 0.1–3 mA	

Fox et al. (2020)	67	Epilepsy	SEEG or/and subdural strip of electrodes	SL (vest)	Frontal, temporal, parietal and cingulate cortices	Case 201: “Vestibular effects, such as feelings of rotation, flotation and acceleration”.	F: 50 Hz I: 4 mA	- 62.792	- 35.335	- 6.726 ³
						Case 202: “Vestibular effects, such as feelings of rotation, flotation and acceleration”.	F: 50 Hz I: 3 mA	- 45.859	- 65.760	8.723 ³
						Cases 203: “Vestibular effects, such as feelings of rotation, flotation and acceleration”.	F: 50 Hz I: 10 mA	- 37.649	- 65.592	49.965 ³
							F: 50 Hz I: 10 mA	- 24.163	- 60.957	52.125 ³
						Case 204: “Vestibular effects, such as feelings of rotation, flotation and acceleration”.	F: 50 Hz I: 4 mA	- 53.879	0.564	43.596 ³
						Case 205: “Vestibular effects, such as feelings of rotation, flotation and acceleration”.	F: 50 Hz I: 6 mA	- 2.452	- 39.961	44.367 ³
						Case 206: “Vestibular effects, such as feelings of rotation, flotation and acceleration”.	F: 50 Hz I: 8 mA	- 16.193	- 34.338	75.675 ³
Cases 207: “Vestibular effects, such as feelings of rotation, flotation and acceleration”.	F: 50 Hz I: 8 mA	- 8.598	46.356	11.650 ³						
	F: 50 Hz I: 8 mA	- 10.517	46.903	0.683 ³						
Sun et al. (2021b)	33	Epilepsy	SEEG	Ag.	Primary somatosensory cortex	Cases 208–210: “Urge to move”. ⁴	F: 50 Hz I: 0.1–5.0 mA	BA 3 (side unknown)		
						Case 211: “Urge to move”. ⁴		BA 1 (side unknown)		
						Cases 212–216: “Urge to move”. ⁴		BA 2 (side unknown)		
Parvizi et al. (2021)	1	Epilepsy	SEEG	Other	Posteromedial cortex, medial frontal cortex, lateral posterior parietal cortex, mesial and lateral temporal lobes and anterior and left posterior insula	Case 217: “Okay, I know. Maybe my depersonalization? This feeling of being disconnected from something. (...) I don't want to say weightlessness, or.... It's like being weightless in your own mind as a personality”.	F: 50 Hz (electrodes LJ3–LJ4) I: 6 mA	Left posteromedial cortex (precuneus)		
						Case 217: “My personality was more attached to my body. That was less like floating in myself as a personality. That was less like who I am was cut off from the other parts of my body. That was more like, it was more like imagine if 40% of who you are becomes what you are and 60% becomes diffuse. So, the dissociation you become 40% of. You know how people will describe consciousness as difficult to describe. You ever hear that, I don't know how to put it.”	F: 50 Hz (electrode LJ4) I: 8 mA			
						Case 217: “Absolutely, so spatial coordinates are just geometry right. So, you have the plane and that theta angle, and you have 3D space. My sense of self does not associate with 3D space at all. Even day to day I would be pretty impressed either of your three selves did. Your sense of self is associated more with, my sense of self is associated more with the characteristics and qualities of who I am and the way I think, right? And the way I process information. So, my sense of self can be removed from 3D space. This could be because of epilepsy, this could easily be because of that. But if I can't move my right leg that does not make me different. Um, I would almost describe this as an ongoing thing maybe because I've had so many episodes of my epilepsy, sorry seizures. But I no longer, for a very long time I would	F: 50 Hz (electrode LJ5) I: 8 mA			

					say, those were two separate things. So, what you just kind of put me, <i>what you created was that separation. But not of my control over my body. Maybe in the same way a pilot can lose control over a plane. They can be forced out of the cockpit but still, or forced to not control it, but still see what is happening to the whole plane. That's kind of what just happened. I got pulled out of the cockpit, but, or I got pulled out of the chair, the pilot's chair, but I could still see all the gauges, and all the.... You know how I always try to. (...) No, not passive.... This is what I mean by a lot of the things I've told you guys is that there are ways to make whoever is in control of the plane snap back, snap everything back into place or for the plane to realize I can sit in this chair again. What you did was maybe move me out of that chair a little bit. But I'm very used to being out of that chair. (...) No, I was floating more in myself.</i>			
					Case 217: "My confidence in my ability to hold on to my body and my sensors really for lack of a better phrase. I'm losing confidence in it and I know where it's going so I can still reach a chair. I guess I can wrestle control, I can still push myself into a chair and into a position where I won't fall".	F: 50 Hz (electrodes L4-5) I: 8 mA		
Bratu et al. (2021)	1	Epilepsy	SEEG	SL (OBE), IPP	Temporal, occipital, frontal, cingulate, parietal, insular cortex	Case 218: "The patient mentioned seeing his face, mouth from outside his body, from above like in a mirror with no additional sensation, no other visual symptoms."	F: 43.3 Hz I: 6 mA	Right anterior part of the periventricular nodular heterotopia
Hao et al. (2022)	376	Epilepsy	SEEG	BI	Temporal, occipital, frontal, cingulate, parietal, insular cortex	Case 219: "Loss of the right hand". (Patient 1)	F: 50 Hz I: 3 mA D: 3 s	Left posterior short gyrus of the insula
						Case 220: "Loss of the right hand". (Patient 2)	F: 50 Hz I: 4 mA D: 3 s	Left posterior dorsal cingulate
						Case 221: "Loss of the right hand". (Patient 3)	F: 50 Hz I: 1 mA D: 3 s	Left posterior part of the anterior long gyrus of the insula
						Case 221: "Loss of the right upper limb". (Patient 3)	F: 50 Hz I: 2 mA D: 3 s	Left parietal operculum
						Case 221: "Sensation of loss of the right side of the body". (Patient 3)	F: 50 Hz I: 2 mA D: 3 s	Left parietal operculum

References

- Andelman-Gur, M.M., Gazit, T., Andelman, F., Kipervasser, S., Kramer, U., Neufeld, M.Y., Fried, I., Fahoum, F., 2020. Spatial distribution and hemispheric asymmetry of electrically evoked experiential phenomena in the human brain. *J Neurosurg* 133, 54–62. <https://doi.org/10.3171/2019.3.JNS183429>
- Andelman-Gur, M.M., Gazit, T., Fahoum, F., Andelman, F., Fried, I., 2019. Negative and positive volitional responses induced by stimulating the superior frontal gyrus: A case study. *Brain Stimul* 12, 1614–1616. <https://doi.org/10.1016/j.brs.2019.08.011>
- Arzy, S., Seeck, M., Ortigue, S., Spinelli, L., Blanke, O., 2006. Induction of an illusory shadow person. *Nature* 443, 287. <https://doi.org/10.1038/443287a>
- Balestrini, S., Francione, S., Mai, R., Castana, L., Casaceli, G., Marino, D., Provinciali, L., Cardinale, F., Tassi, L., 2015. Multimodal responses induced by cortical stimulation of the parietal lobe: a stereo-electroencephalography study. *Brain* 138, 2596–2607. <https://doi.org/10.1093/brain/awv187>
- Blanke, O., Ortigue, S., Landis, T., Seeck, M., 2002. Stimulating illusory own-body perceptions. *Nature* 419, 269–270. <https://doi.org/10.1038/419269a>
- Blanke, O., Perrig, S., Thut, G., Landis, T., Seeck, M., 2000. Simple and complex vestibular responses induced by electrical cortical stimulation of the parietal cortex in humans. *J Neurol Neurosurg Psychiatry* 69, 553–556. <https://doi.org/10.1136/jnnp.69.4.553>
- Blanke, O., Pozeg, P., Hara, M., Heydrich, L., Serino, A., Yamamoto, A., Higuchi, T., Salomon, R., Seeck, M., Landis, T., Arzy, S., Herbelin, B., Bleuler, H., Rognini, G., 2014. Neurological and robot-controlled induction of an apparition. *Curr Biol* 24, 2681–2686. <https://doi.org/10.1016/j.cub.2014.09.049>
- Bos, E.M., Spoor, J.K.H., Smits, M., Schouten, J.W., Vincent, A.J.P.E., 2016. Out-of-Body Experience During Awake Craniotomy. *World Neurosurgery* 92, 586.e9-586.e13. <https://doi.org/10.1016/j.wneu.2016.05.002>
- Caruana, F., Gerbella, M., Avanzini, P., Gozzo, F., Pelliccia, V., Mai, R., Abdollahi, R.O., Cardinale, F., Sartori, I., Lo Russo, G., Rizzolatti, G., 2018. Motor and emotional behaviours elicited by electrical stimulation of the human cingulate cortex. *Brain* 141, 3035–3051. <https://doi.org/10.1093/brain/awy219>
- De Ridder, D., Van Laere, K., Dupont, P., Menovsky, T., Van de Heyning, P., 2007. Visualizing out-of-body experience in the brain. *N Engl J Med* 357, 1829–1833. <https://doi.org/10.1056/NEJMoa070010>
- Desmurget, M., Reilly, K.T., Richard, N., Szathmari, A., Mottolese, C., Sirigu, A., 2009. Movement intention after parietal cortex stimulation in humans. *Science* 324, 811–813. <https://doi.org/10.1126/science.1169896>
- Fornia, L., Puglisi, G., Leonetti, A., Bello, L., Berti, A., Cerri, G., Garbarini, F., 2020. Direct electrical stimulation of the premotor cortex shuts down awareness of voluntary actions. *Nature Communications* 11, 705. <https://doi.org/10.1038/s41467-020-14517-4>
- Fox, K.C.R., Shi, L., Baek, S., Raccach, O., Foster, B.L., Saha, S., Margulies, D.S., Kucyi, A., Parvizi, J., 2020. Intrinsic network architecture predicts the effects elicited by intracranial electrical stimulation of the human brain. *Nat Hum Behav* 4, 1039–1052. <https://doi.org/10.1038/s41562-020-0910-1>
- Fried, I., Katz, A., McCarthy, G., Sass, K., Williamson, P., Spencer, S., Spencer, D., 1991. Functional organization of human supplementary motor cortex studied by electrical stimulation. *J Neurosci* 11, 3656–3666. <https://doi.org/10.1523/JNEUROSCI.11-11-03656.1991>
- Halgren, E., Walter, R.D., Cherlow, D.G., Crandall, P.H., 1978. Mental phenomena evoked by electrical stimulation of the human hippocampal formation and amygdala. *Brain* 101, 83–117. <https://doi.org/10.1093/brain/101.1.83>
- Hao, G., Wang, X., Yan, H., He, L., Ni, D., Qiao, L., Zhang, X., Yu, K., Yu, T., 2022. Limb loss experience evoked by electric cortical stimulation. *Epileptic Disord* 24, 67–74. <https://doi.org/10.1684/epd.2021.1364>
- Isnard, J., Guénot, M., Sindou, M., Mauguère, F., 2004. Clinical Manifestations of Insular Lobe Seizures: A Stereo-electroencephalographic Study. *Epilepsia* 45, 1079–1090. <https://doi.org/10.1111/j.0013-9580.2004.68903.x>
- Kahane, P., Hoffmann, D., Minotti, L., Berthoz, A., 2003. Reappraisal of the human vestibular cortex by cortical electrical stimulation study. *Ann Neurol* 54, 615–624. <https://doi.org/10.1002/ana.10726>
- Kremer, S., Chassagnon, S., Hoffmann, D., Benabid, A., Kahane, P., 2001. The cingulate hidden hand. *J Neurol Neurosurg Psychiatry* 70, 264–265. <https://doi.org/10.1136/jnnp.70.2.264>
- Mandonnet, E., Margulies, D., Stengel, C., Dali, M., Rheault, F., Toba, M.N., Bonnetblanc, F., Valero-Cabre, A., 2020. “I do not feel my hand where I see it”: causal mapping of visuo-proprioceptive integration network in a surgical glioma patient. *Acta Neurochir (Wien)* 162, 1949–1955. <https://doi.org/10.1007/s00701-020-04399-2>
- Mazzola, L., Lopez, C., Faillenot, I., Chouchou, F., Mauguère, F., Isnard, J., 2014. Vestibular responses to direct stimulation of the human insular cortex. *Ann Neurol* 76, 609–619. <https://doi.org/10.1002/ana.24252>

- Mulak, A., Kahane, P., Hoffmann, D., Minotti, L., Bonaz, B., 2008. Brain mapping of digestive sensations elicited by cortical electrical stimulations. *Neurogastroenterology & Motility* 20, 588–596. <https://doi.org/10.1111/j.1365-2982.2007.01066.x>
- Mullan, S., Penfield, W., 1959. Illusions of comparative interpretation and emotion; production by epileptic discharge and by electrical stimulation in the temporal cortex. *AMA Arch Neurol Psychiatry* 81, 269–284.
- Oane, I., Barborica, A., Chetan, F., Donos, C., Maliia, M.D., Arbune, A.A., Daneasa, A., Pistol, C., Nica, A.E., Bajenaru, O.A., Mindruta, I., 2020. Cingulate cortex function and multi-modal connectivity mapped using intracranial stimulation. *NeuroImage* 220, 117059. <https://doi.org/10.1016/j.neuroimage.2020.117059>
- Ostrowsky, K., Desestret, V., Ryvlin, P., Coste, S., Mauguière, F., 2002. Direct electrical stimulations of the temporal pole in human. *Epileptic Disord* 4 Suppl 1, S23-27.
- Parvizi, J., Braga, R.M., Kucyi, A., Veit, M.J., Pinheiro-Chagas, P., Perry, C., Sava-Segal, C., Zeineh, M., Staalduinen, E.K. van, Henderson, J.M., Markert, M., 2021. Altered sense of self during seizures in the posteromedial cortex. *PNAS* 118. <https://doi.org/10.1073/pnas.2100522118>
- Penfield, W., 1957. Vestibular sensation and the cerebral cortex. *Ann Otol Rhinol Laryngol* 66, 691–698. <https://doi.org/10.1177/000348945706600307>
- Penfield, W., 1955. The twenty-ninth Maudsley lecture: the role of the temporal cortex in certain psychical phenomena. *J Ment Sci* 101, 451–465. <https://doi.org/10.1192/bjp.101.424.451>
- Penfield, W., 1947. Some observations on the cerebral cortex of man. *Proc R Soc Lond B Biol Sci* 134, 329–347. <https://doi.org/10.1098/rspb.1947.0017>
- Penfield, W., Boldrey, E., 1937. Somatic motor and sensory representation in the cerebral cortex of man as studied by electrical stimulation. *Brain* 60, 389–443. <https://doi.org/10.1093/brain/60.4.389>
- Penfield, W., Perot, P., 1963. The brain's record of auditory and visual experience: a final summary and discussion. *Brain* 86, 595–696. <https://doi.org/10.1093/brain/86.4.595>
- Penfield, W., 1938. The cerebral cortex in man: the cerebral cortex and consciousness. *Archives of Neurology & Psychiatry* 40, 417–442. <https://doi.org/10.1001/archneurpsyc.1938.02270090011001>
- Popa, I., Barborica, A., Scholly, J., Donos, C., Bartolomei, F., Lagarde, S., Hirsch, E., Valenti-Hirsch, M.-P., Maliia, M.D., Arbune, A.A., Daneasa, A., Ciurea, J., Bajenaru, O.-A., Mindruta, I., 2019. Illusory own body perceptions mapped in the cingulate cortex-An intracranial stimulation study. *Hum Brain Mapp* 40, 2813–2826. <https://doi.org/10.1002/hbm.24563>
- Richer, F., Martinez, M., Robert, M., Bouvier, G., Saint-Hilaire, J.M., 1993. Stimulation of human somatosensory cortex: tactile and body displacement perceptions in medial regions. *Exp Brain Res* 93, 173–176. <https://doi.org/10.1007/BF00227792>
- Salanova, V., Andermann, F., Rasmussen, T., Olivier, A., Quesney, L.F., 1995a. Parietal lobe epilepsy. Clinical manifestations and outcome in 82 patients treated surgically between 1929 and 1988. *Brain* 118 (Pt 3), 607–627. <https://doi.org/10.1093/brain/118.3.607>
- Salanova, V., Andermann, F., Rasmussen, T., Olivier, A., Quesney, L.F., 1995b. Tumoural parietal lobe epilepsy. Clinical manifestations and outcome in 34 patients treated between 1934 and 1988. *Brain* 118 (Pt 5), 1289–1304. <https://doi.org/10.1093/brain/118.5.1289>
- So, E.L., Schaüble, B.S., 2004. Ictal asomatognosia as a cause of epileptic falls: simultaneous video, EMG, and invasive EEG. *Neurology* 63, 2153–2154. <https://doi.org/10.1212/01.wnl.0000145628.38030.3e>
- Sun, F., Zhang, G., Ren, L., Yu, T., Ren, Z., Gao, R., Zhang, X., 2021a. Functional organization of the human primary somatosensory cortex: A stereo-electroencephalography study. *Clin Neurophysiol* 132, 487–497. <https://doi.org/10.1016/j.clinph.2020.11.032>
- Sun, F., Zhang, G., Yu, T., Zhang, Xiaohua, Wang, X., Yan, X., Qiao, L., Ma, K., Zhang, Xi, 2021b. Functional characteristics of the human primary somatosensory cortex: An electrostimulation study. *Epilepsy & Behavior* 118, 107920. <https://doi.org/10.1016/j.yebeh.2021.107920>
- Vignal, J.-P., Maillard, L., McGonigal, A., Chauvel, P., 2007. The dreamy state: hallucinations of autobiographic memory evoked by temporal lobe stimulations and seizures. *Brain* 130, 88–99. <https://doi.org/10.1093/brain/awl329>
- Wiest, G., Zimprich, F., Prayer, D., Czech, T., Serles, W., Baumgartner, C., 2004. Vestibular processing in human paramedian precuneus as shown by electrical cortical stimulation. *Neurology* 62, 473–475. <https://doi.org/10.1212/01.wnl.0000106948.17561.55>
- Yu, K., Liu, C., Yu, T., Wang, X., Xu, C., Ni, D., Li, Y., 2018a. Out-of-body experience in the anterior insular cortex during the intracranial electrodes stimulation in an epileptic child. *J Clin Neurosci* 54, 122–125. <https://doi.org/10.1016/j.jocn.2018.04.050>
- Yu, K., Yu, T., Qiao, L., Liu, C., Wang, X., Zhou, X., Ni, D., Zhang, G., Li, Y., 2018b. Electrical stimulation of the insulo-opercular region: visual phenomena and altered body-ownership symptoms. *Epilepsy Res* 148, 96–106. <https://doi.org/10.1016/j.eplepsyres.2018.09.014>